

Guidelines for the selection of biological Areas of Special Scientific Interest (ASSIs) on the Isle of Man

Volume 1: Background, guiding principles and Priority Sites Criteria



**Isle of Man
Government**

Reiltys Ellan Vannin

Guidelines for the selection of biological Areas of Special Scientific Interest (ASSIs) on the Isle of Man

This document sets out the background, rationale, guiding principles and minimum criteria for the selection of Areas of Special Scientific Interest (ASSIs) on the Isle of Man. It is one of a series of documents intended to provide a sound scientific basis for the selection of legally protected areas on and around the Island.

The Wildlife Act 1990 provides the basis for the designation of ASSIs on the Isle of Man. Reasons for designation may include terrestrial, marine or geological importance of a site, or a combination of features. Please note that the criteria detailed in this document are for the selection of land-based ASSIs for biological reasons. These may include coastal and intertidal zones down to the astronomical low tide limit, but the selection of strictly Marine Nature Reserves is covered by a separate, companion document. Criteria for ASSIs selected primarily for their earth science importance (geological and physiographical features) will also be treated separately. An ASSI may contain a mix of features of importance from any or all of the above categories, which often occur in association. In such cases, more than one set of criteria may be applicable.

The purpose of the ASSI system is to safeguard a series of sites which are individually of high natural heritage importance and which collectively represent the diversity of habitats, species and geological and geomorphological features on the Isle of Man. The aim of the Criteria is to provide a logical and consistent basis for site selection, using recognised nature conservation principles. The Criteria also enable Manx sites to be placed in a wider context, so that internationally threatened and vulnerable biodiversity, for which the Island has a shared responsibility, can be taken into account.



Bee orchid *Ophrys apifera*, a protected species restricted to two sites on the Island

Volume 1 – General guidelines

Contents

Part 1: Background and rationale.....	3
1.1 Statutory wildlife protection on the Isle of Man.....	3
Table 1: summary of nature conservation site protection measures on the Isle of Man.....	4
1.2 Aims of the Manx biological ASSI system	4
1.3 The Manx context.....	5
1.4 ASSI designation - the reasons.....	7
1.5 ASSI designation - the process.....	8
1.6 Types of site selection criteria	9
Part 2: General principles	11
2.1 Naturalness and scientific importance	11
2.2 Dates of records and use of population data.....	11
2.3 Boundaries.....	11
2.4 Habitat assemblages.....	12
2.5 Fragmented sites	13
2.6 Significance of species.....	13
2.7 Marine species and habitats	13
2.8 Geological and physiographical sites	13
2.9 Monitoring	13
2.10 Revision of site status.....	14
Part 3: Priority Sites Criteria	15
3.1 Checklist of Priority Sites Criteria.....	15
3.2 Using the Priority Sites Criteria	17

Part 4: Detailed selection guidelines, and Part 5: Bibliography and references are contained within **Volume 2**.

Part 1: Background and rationale

1.1 Statutory wildlife protection on the Isle of Man

The Wildlife Act 1990 enables habitats and species to be protected by law. Section 27 of the Act includes the following provision:

(1) *Where the Department, after consultation with the Wildlife Committee is of the opinion that any area of land is of special interest by reason of any of its flora, fauna, or geological or physiographical features, it may notify that fact to-*

- (a) the Department of Local Government and the Environment; and*
- (b) to every owner and occupier of any of that land.*


Such areas of land include Areas of Special Scientific Interest (ASSIs). The Wildlife Act 1990 also provides for the designation of Areas of Special Protection (e.g. for birds), Marine Nature Reserves, National Nature Reserves, and a range of specific measures designed to protect individual bird, animal and plant species. There are a range of other designations applicable to the Isle of Man, including locally important Wildlife Sites and internationally recognised Ramsar Sites. ASSIs therefore fit into a range of protective measures, based on the level of importance of the site as follows:

- **Undesignated areas.** The abundance of improved farmland habitats in the wider countryside, whilst important for some species, makes it usually the lowest priority for designation as a protected area. Likewise, semi-improved habitats along the coast and in the wider countryside which don't meet the criteria for designation as a Wildlife Site are unlikely to receive special protection on the grounds of nature conservation. However, these habitats may occasionally fall within areas under the protection of Manx National Heritage (see below), and are all protected to a certain extent by agricultural codes of practice and planning regulations. It should also be remembered that some important wildlife species, such as Chough, depend on productive farmland which is subject to ordinary farming practices. These species would not normally be protected through designating their habitat but by conservation management agreements (under Section 30 of the Wildlife Act 1990).
- **Wildlife Sites.** This is a non-statutory designation, used widely throughout the British Isles as a site protection system which is generally operated by local planning authorities and/or the Wildlife Trusts. On the Isle of Man, Wildlife Sites are a voluntary designation, selected using criteria designed by the Manx Wildlife Trust and agreed amongst relevant organisations;
- **Bird Sanctuaries.** Designated under the Protection of Birds Acts 1932 & 1955, and remain protected although this legislation has now been superseded by the Wildlife Act 1990;
- **Areas of Special Protection.** May be designated under Section 13 of the Wildlife Act 1990 in order to extend the provision of the Wildlife Act for certain species in certain areas, e.g. the Area of Special Protection for Birds at the Ayres Gravel Pit;
- **Areas of Special Scientific Interest (ASSIs).** Detailed in this document;
- **National Nature Reserves (NNRs).** Designated under Section 31 of the Wildlife Act 1990, may be on private or public land, and may be subject to protective byelaws. NNRs are likely to be of ASSI quality or higher, i.e. of national importance;
- **Marine Nature Reserves.** Designated under Section 32, are also of national importance for nature conservation, and may also be protected with byelaws;
- **Land in the ownership of the Manx National Trust** (administered as Manx National Heritage). This land is protected under the Manx Museum and National Trust Act (1959-1982), which provides for the protection of places 'of natural interest or beauty' and places of historic interest. This includes wildlife, historic buildings and also the landscape setting and physical features within it. Whilst land owned by Manx National Heritage

will not always be acquired on the basis of its wildlife alone (since cultural heritage and landscape setting may also be important factors), the fauna and flora of that land is protected by byelaws whether they are of local, Manx national, or international value.

- Similarly, some **Manx Wildlife Trust reserves** may meet the criteria for ASSI designation while other areas may meet Wildlife Site selection criteria;
- **Ramsar Sites** are internationally important wetland sites chosen according to criteria agreed by parties to the Ramsar Convention on Wetlands (1971). The Isle of Man is a signatory to this convention, and thus has an obligation to investigate potential Ramsar Sites on the Island. Such sites will be of ASSI or higher status;
- **Emerald Network** sites are internationally important wildlife areas comparable with the European Union's Natura 2000 network. Sites consist of Areas of Special Conservation Interest designated under a recommendation of the Bern Convention (to which the Isle of Man is signatory). Thus, although not part of the EU and its Natura 2000 network (which includes Special Areas of Conservation in the UK), the Island may still designate internationally important sites at the same level. If designated, these sites are likely to include the most important ASSIs on the Island.

Table 1: summary of nature conservation site protection measures on the Isle of Man

Scale of priority for protection Greater priority  Lesser priority				
Ramsar Sites	National Nature Reserves (NNRs)	Wildlife Sites	Habitats in farmland and wider countryside	Improved farmland
Emerald Network Sites	Areas of Special Scientific Interest (ASSIs)			
	Marine Nature Reserves (MNRs)			
	Areas of Special Protection (ASPs)			
	Bird Sanctuaries			
Manx National Trust land – could apply to any of the above.				

1.2 Aims of the Manx biological ASSI system

•

1.3 The Manx context

An Island-based approach

The Selection Criteria have been written to fulfil the needs of wildlife conservation in the context of the Isle of Man as a Crown Dependency, with its own Wildlife Act. Sites will be selected based on what is important for the Island as a nation rather than as if it were a county of the UK.

The reasons for an Island-based approach, rather than simply using the JNCC criteria for SSSIs in the UK, are:

- the need to retain areas of habitat and populations of species that are rare on the Island, even if they are commoner elsewhere, to maintain the biodiversity of the Island;
- the isolation of the Island, resulting in limited prospects for natural recolonisation from surrounding land masses if species are lost and become extinct here;
- because small populations are more vulnerable to extinction, therefore wider habitat protection may be needed in order to maintain a viable population;
- to ensure the protection of any species or habitats which show a distinctly Manx characteristic not found elsewhere.

The objective is to maximise the conservation of species biodiversity through protection of their habitats. Sites will therefore be considered on the basis of their Island-wide/national, British Isles or international/European importance. The aim is for the ASSI series to include at least one good example of each main semi-natural habitat type and assemblage. For very scarce and important habitats, such as saltmarsh, the majority of examples will require protection.

The Island's origins and ecology

The Isle of Man was joined to Britain by a land bridge until around nine thousand years ago (Allen 1978). The melt water from the ice sheet led to the rise in sea levels which cut off the various British islands and separated them from the Continent. The present day native flora and much of the fauna are descended from the species that colonised after the retreat of the last ice sheet. During the last Ice Age the Island was covered by an ice sheet and no evidence of glacial refuges (small pockets of land not covered by ice) has been found. It became separated from the rest of the British Isles by the formation of the Irish Sea during the melting of the ice. Isolated islands which have been cut off for many thousands of years usually have endemic species. Although no Manx endemic species have been identified there are likely to be sub-species or genotypes specific to the Island.

Some species moving north with the warming climate would have reached the Island before others. The formation of the Irish Sea cut the Island off from further terrestrial invasion, but some animals would have arrived before then. Later immigrants would have arrived naturally by air or sea. Not all the Manx fauna have clear origins however: for example, one of the mysteries of migration and colonisation is how the lesser mottled grasshopper came to the Isle of Man and yet is found nowhere else in the British Isles.

The Isle of Man is much smaller than mainland Britain and we know, through the study of island biogeography, that the number of fauna species on an island is related to its size. Small islands also have a high turnover of species because small, isolated populations are prone to extinction, both from strong climatic events and man-induced changes.

Island biodiversity is normally lower than that of the adjacent larger landmasses. The Isle of Man follows this pattern. *"In every plant and animal order (except the wholly freshwater ones) that has so far been adequately worked, with striking consistency Man proves to have two thirds of the Irish total and two-fifths of the British"* (Allen 1984). There is thus a noticeable difference between the Manx fauna and that of mainland Britain. A number of mammal species are absent, including the badger, otter, voles and the British weasel. Likewise some birds which are resident in Britain are rarely seen on the Island, such as woodpeckers and buzzards.

Mann is closest to Galloway (16 miles), Cumbria and Ulster in that order. Wales is rather further off. Although the Manx ecology generally reflects this position, the situation is tempered by the mild oceanic climate, which has led to a low frequency and severity of frost and more hospitable conditions for some species than would normally be expected given the northerly latitude.

The Island is 227 square miles in area (58,793 hectares) and within this landmass the geology affects the landform and soil type, which in turn affect the development of habitats.

Woodland animals might have been commoner in the past before most of the trees were cleared by early settlers. Most of the present woodland on the Island has been planted within the last hundred and fifty years. A few sites have older plantations and some very small remnants of natural woodland are thought to survive; the woodland fauna on the Island is correspondingly restricted.

Of the Island's birds, the dipper and corn bunting seem to have been lost as breeding species, corncrakes continue to breed but only in extremely low numbers, but the fulmar established a breeding population from 1936, and the hen harrier, which has been a non-breeding visitor since 1900, established a significant breeding population in the last quarter of the twentieth century. However, the buzzard, although expanding in distribution in the UK, is seen only rarely, passing through. The merlin, another raptor breeding on moorlands, is also rarely seen in the breeding season.

The Island has a good variety of habitats, but some of them are of very limited extent, and are very remote from similar areas that could act as source populations. Hence they are vulnerable to loss of rarer and more specialised fauna which would have little chance of re-colonising from similar habitats nearby.

Despite having less overall species richness than the UK, the Island is home to several species of special interest. It is particularly noted for the lesser mottled grasshopper, which is found nowhere else in the British Isles. Although the great auk became extinct from Mann as elsewhere, the Island still remains a stronghold for species of the rocky west coast of Europe, such as the chough and peregrine falcon. Some coastal species are found here at the northern edge of their range, such as the centipede *Schendyla peyerimhoffi*, found on beaches.

Climatic factors and vegetation communities

Climatic factors have had a marked effect on the distribution and variety of vegetation communities on the Isle of Man. The Island lies at the same latitude as the north of England and Northern Ireland, and has a correspondingly short growing season for plants. However, it benefits from the warming effects of North Atlantic Drift, and tends not to receive the sharp frosts and heavy snows found at the same latitude further east. Thus it has a generally wet and windy maritime climate, with a longer growing season and less precipitation than the Hebridean islands, but colder conditions than islands off the coast of Wales and Cornwall.

Conditions are extremely variable from one part of the Island to another: the highest point of Snaefell barely falls within the range of British arctic-alpine vegetation, with permanently cold and exposed conditions, whilst the south coast can provide very warm and sunny situations, almost comparable with the level of warmth and light found on south-facing coasts in the south west of the UK. Precipitation levels in drier parts of the Island, such as Port St Mary, are, on average, less than half the levels in the wettest areas, such as Laxey. This wide variety of climatic conditions has led to a situation where hardly any two areas of semi-natural vegetation are exactly the same in terms of biodiversity.

The National Vegetation Classification (NVC), as detailed in *British Plant Communities* (Rodwell *et al*, JNCC 1995), provides valuable comparative descriptions of plant communities and will be consulted during selection of ASSI habitats. However, some vegetation communities on the Isle of Man only partially match National Vegetation Classification types. This may be due to the "in-between" position in relation to other parts of the British Isles, the low level of sampling on the Island whilst the NVC was being researched, or the naturally low biodiversity typical of this and other islands. For example, the few areas of calcareous grassland (CG in the NVC) on the Island may show characteristics of CG2, CG10 and maritime swards all within the same vegetation community. Likewise the Manx "curragh", today dominated by mature tallows and with a few larger trees (once much more open wetland), does not quite fit the usual woodland or wet scrub ("willow carr") NVC description.

The poor match which sometimes occurs between Manx habitats and NVC types does not necessarily mean that these habitats are degraded examples of the type; they may be characteristic, diverse Manx habitats of high

wildlife importance that have developed a slightly different species composition to UK vegetation communities identified during the NVC research. In view of this, NVC types alone are not considered sufficient to provide a set of criteria for selecting a series of ASSIs representative of all the Island's characteristic semi-natural habitats. Other factors, for example presence or absence of rare species with a restricted distribution on the Island, will also be taken into account. The relationship between the Selection Criteria and the NVC is detailed in Part 3 of this document.

1.4 ASSI designation - the reasons

Conserving biodiversity

The importance of conserving the richness and abundance of wildlife is widely recognised. A range of international conventions exist in order to further the protection of biodiversity for the benefit of future generations. In 1992 the Convention on Biological Diversity met and highlighted the importance of maintaining biodiversity for the health of the earth and its people. The Convention has over one hundred and fifty signatories who are committed to conserving habitats and species and managing biodiversity sustainably. Other international conventions include the Berne Convention on the Conservation of European Wildlife and Natural Habitats, which urges contracting parties (which include the Isle of Man) to establish a network of protected sites and identify those of European importance (the "Emerald Network"). The Bonn Convention on the Conservation of Migratory Species of Wildlife Animals, and related agreements on bats and migratory water birds, require the conservation of habitats for such species, as the loss of a site used for one part of the life cycle of a species (See 1.1 above) may jeopardise its survival by breaking the links.

The chief aim of international wildlife conventions is to prevent further extinctions of species in the wild and ensure their sustainability for future generations. The Isle of Man is not immune to the threat of extinctions; they can happen at the global scale down to the local scale as people change their environment. On the Manx scale some species have gone extinct in the last 100 years (eg ringlet butterfly and dense-flowered orchid), but other more mobile or adaptive species have arrived naturally (eg collared doves).

The process of designating biological ASSIs on the Isle of Man has one overriding aim:

To conserve the Island's biodiversity by ensuring that good examples of all the Island's native species populations, and the characteristic habitats they rely upon, are protected by law from activities that are known to jeopardize, damage, or destroy them.

This aim will be achieved, not just by identifying and designating a suitable series of ASSIs as "areas on a map", but also by working with landowners and users to help them undertake the best management for the Island's most important wildlife sites.



Round-leaved sundew Drosera rotundifolia, on a site managed for its moorland and bog

Early surveys

You can't conserve something until you know what you have, how many there are and where they are. Over the centuries various enthusiastic individuals and groups have recorded the flora and fauna of the Island, mostly concentrating on certain plant or animal groups. The first comprehensive habitat survey was carried out in the early 1970s, when the government commissioned an ecological survey from the Nature Conservancy Council and the Institute of Terrestrial Ecology. In 1975 they published their report, which identified potential national nature reserves and smaller "Sites of Ecological Importance". The latter were envisaged as having comparable statutory status to the UK's Sites of Special Scientific interest and (in Northern Ireland) Areas of Special Scientific Interest, (SSSIs/ASSIs). These 34 sites, and the criteria used to select them (see below), are listed in Garrad (1990).

It wasn't until 1991 that a systematic land use and habitat survey was undertaken by Government. Shortly after the passing of the Wildlife Act in 1990 the Department of Agriculture Fisheries and Forestry set up the Ecological Survey team to map the Island's habitats and land uses, using Phase 1 survey techniques developed by the Nature Conservancy Council in the UK. The largest areas of lowland semi-natural vegetation thought to have the best potential wildlife conservation value were then mapped at Phase 2 level. Meanwhile the Biological Records Centre was established at the Manx Museum to hold all the biological information. These surveys provide the basis for the list of sites to be evaluated as ASSIs for their habitat importance - a list which includes and enlarges on the 34 sites in NCC (1975). Other sites may also be considered, either on the basis of species interest or where new information indicates high importance.

What proportion of the land should be designated ?

In 1998 a programme for achieving full coverage of ASSIs was initiated. The target has been estimated to be a minimum of **10%** of the area of the Island (5,650 ha), until a proper evaluation of biodiversity has been undertaken. This figure was based on a comparison with the situation in the UK, where in Scotland the percentage is 12.8% (Scottish Natural Heritage figure for 2002) and in England 7% (English Nature, 2002).

Site protection is only one part of the process of conserving the Island's biodiversity. Many species depend on farmed land and other habitats in the wider countryside, therefore the ASSI network must be complimented by other measures such as the Agri-environment Scheme and the assignment of Areas of Special Protection (ASP) – see 1.1 above.

1.5 ASSI designation - the process

Consultation and decision-making

The Wildlife Committee is a statutory body established by the Wildlife Act 1990 to act as a scientific advisory committee for all matters arising from the Act. The Committee consists of advisers from DAFF and other government bodies as well as academic and research organisations. This is the forum for considering designation proposals.

Sites will be selected on the basis of the criteria contained within this document, and on the professional judgement of the DAFF Wildlife Office staff in conjunction with the Wildlife Committee. The legal relationship and implications of ASSI designation on land already protected by other legislation, such as the Manx Museum and National Trust Act (1959-1982), will be defined. The recommendations arising from this process will then be referred to the Department for agreement and ratification.

A leaflet explaining the designation process based on the procedures laid down in the Wildlife Act 1990 is available from DAFF.

1.6 Types of site selection criteria

RSNC's Wildlife Sites Handbook and the Manx Wildlife Sites Handbook

Published in 1977, Version 2 of the Royal Society for Nature Conservation (RSNC – now the Royal Society for Wildlife Trusts) *Wildlife Sites Handbook* describes Wildlife Sites as having “substantive nature conservation value”. According to the recommendations of the Handbook, Wildlife Sites must reach the specified thresholds for the county or equivalent area for which they are written (e.g. the Isle of Man).

The Manx Wildlife Site selection criteria are couched in similar terms to the criteria in the RSNC Handbook. Whilst the Ratcliffe criteria are recognised, the need for ease of use has condensed the Wildlife Sites criteria into statements. For example “All H1 (NVC *Calluna vulgaris-Festuca ovina*) grasslands greater than 2.0 ha should be Wildlife Sites”. This effectively incorporates an assessment of botanical diversity, typicalness, size and habitat rarity.

The ASSI and Wildlife Site systems (statutory and voluntary) operate side by side, so that sites not meeting the ASSI criteria but still having substantive wildlife value may qualify as Wildlife Sites.

The “Ratcliffe criteria” methodology

Ratcliffe (1977) was the first to publish criteria by which sites should be selected. As Ratcliffe explained these are criteria which have “*by general agreement and established practice, become accepted as a means of judging the nature conservation value of a defined area of land*”. The Ratcliffe approach was to become the main rationale for the UK Nature Conservancy Council (now Joint Nature Conservation Committee or JNCC) “Guidelines for selection of biological SSSIs” (see **1.6.3**, below) .

Some of the Ratcliffe criteria might be termed primary criteria:

- Size (extent)
- Naturalness
- Typicalness
- Rarity
- Fragility
- Diversity (of habitat, structure, plant community and species)
- Position in the ecological/geographical unit

Others are secondary criteria:

- Recorded history
- Potential value (if restored)
- Intrinsic appeal

Manx first evaluation

The criteria applied to the Isle of Man by the Nature Conservancy Council (NCC/ITE 1975; Garrad *per* Robinson & McCarroll, 1990) are generally referred to as the Garrad criteria. These are more specific versions of Ratcliffe's criteria. They are heavily biased towards botanical and vegetation characteristics of sites, making little use of fauna, probably because it is less well known (especially the invertebrates) and often dependent on the flora and habitats in any case. Under these criteria, Manx sites are considered on the basis of the:

- Presence of plants not found elsewhere on the Island
- Presence of plants rare on Mann (fewer than 10 sites)
- Presence of plants rare in this area of the Island
- Number of plant species present.
- Degree to which plant assemblage is representative of the habitat
- Degree to which plant assemblage is markedly unusual for the habitat
- Variety of plant species present
- Presence of unusual plant communities
- Presence of representative plant communities
- Presence of assemblages or habitats that are

- a) rare in Mann
 - b) rare in this area of the Island
 - c) rare in British Isles as a whole
 - d) rare in northwest Europe
- Diversity of habitats
- Major bird uses
 - a) passage on migration
 - b) wintering
 - c) breeding

Both the Ratcliffe and Garrad criteria are broadly relevant to all site selection, whether for habitat or species interest. In view of this, a “checklist” for assessing the quality of potential ASSIs, based on the Ratcliffe approach, is contained in Part 3 of this document, entitled Priority Sites Criteria.

Guidelines for Selection of Biological SSSIs in the UK

In 1989 the then Nature Conservancy Council (now the Joint Nature Conservation Committee) published a detailed set of criteria for selecting Sites of Special Scientific Interest in the UK. With a lengthy introduction and rationale based on the “Ratcliffe criteria” (see above), the detailed guidelines cover all the habitat types and species groups which at the time were considered relevant to the selection of protected sites. Each habitat and species group is given its own chapter, with a description, classification information and specific guidance on the minimum criteria for site selection. These chapters are periodically updated as new information becomes available. This widely-used approach has been followed in **Parts 3 and 4** of this document, entitled **Priority Sites Criteria** and **Detailed Selection Guidelines**.



Surveying ponds at Rosehill Quarry, Billown, prior to ASSI assessment; the site was subsequently selected as an ASSI, and designated in December 2005.

Part 2: General principles

The criteria by which Manx biological ASSIs are to be selected have been written according to the following general principles. These are applicable throughout, regardless of the habitats or species in question.

2.1 Naturalness and scientific importance

The ASSI criteria are designed to take into account the fact that much of the Island's native biodiversity exists within semi-natural habitats that have been shaped by human activities. However, as is the practice elsewhere, site protection is more likely to be considered a priority if the habitats involved are considered to be unusually pristine examples, exceptionally diverse, a recognised locally distinctive type, or impossible to restore once degraded or lost.

There has been much discussion amongst conservationists about the relative importance of natural, semi-natural and recently-established habitats for biodiversity. This is chiefly due to the fact that a high proportion of native British Isles species have survived through colonising man-made environments during and following the removal of their original ecological niches. The ASSI criteria reflect the prevailing scientific opinion that all habitats with a high complement of native species, in communities which form an interdependent ecosystem, should be valued for their contribution to biodiversity conservation.

Semi-natural habitats are important both for the communities they support, and for the individual rare species that may occur there (especially species that don't occur anywhere else on the Island). Longstanding semi-natural habitats that have resulted from traditional land management practices tend to be more diverse and contain more rare species than farmland where regular rotations and modern fertilising systems are practiced. However, some more recent activities, such as quarrying, may also result in important areas of wildlife value. In the interests of biodiversity conservation, such sites will not be ruled out of the ASSI selection process just because they are not strictly "natural".

2.2 Dates of records and use of population data.

Sites will be assessed using the most recent species and habitat data available. A potential ASSI will not be disqualified from selection on the grounds of deficient data; however, in such cases efforts will be made to confirm the importance of the site prior to designation as a matter of priority.

The acceptable dates for species records which are used in order to assess sites will vary with the species or group. In general, invertebrate and lower plant surveys are less frequent than bird and vascular plant surveys, therefore older records sometimes have to be accepted for use with invertebrate or lower plant-based criteria. Rarity evaluations will make use of the best species distribution information available at the time. If new information comes to light indicating that a species is significantly more common or more rare than previously thought, a site proposed for designation may require a re-evaluation.

2.3 Boundaries

Each ASSI should be large enough to provide adequate site-based protection for the feature(s) for which it has been designated. Site boundaries will therefore be defined with the aim of encompassing all the key ecological features for which the ASSI has been selected, plus any fringing or linking habitats without which the ecology of the site could not be protected, even if those fringing habitats do not in themselves contain key features.

Well-defined boundaries are crucial to the success of any network of protected sites. In defining ASSI boundaries there are two potentially conflicting considerations:

1. The boundary should be the minimum necessary -
 - a) to avoid unnecessary restrictions on land owners;
 - b) to maintain the site integrity and high standards of the ASSI system;
 - c) to reduce unnecessary work and expenditure by DAFF in processing consents and negotiating management agreements.
2. At the same time the boundary must be large enough to protect the site adequately, by ensuring that all scientifically important features are included and that an appropriate conservation management regime can be maintained.

In defining boundaries it should be possible to explain to individual landowners why their land is included in the site while other lands are not. DAFF should therefore be able to demonstrate that a consistent, logical and methodical approach to defining site boundaries has been employed. Providing effective control of management could involve designating all land within the site up to the nearest stock-proof boundary, especially on hill land. This may not necessarily be practical or desirable where large areas of land with minimal nature conservation value are included. A common-sense approach is therefore required to balance the considerations above.

The main principle in defining ASSI boundaries is that all land of prime scientific interest must be included. Boundaries must be clearly indicated, showing any hedges, streams or other linear features which are considered to be part of the designation. Other semi-natural habitats will also be included where they are considered vital for the survival of the key ecological features for which the site has been selected. This might include:

- crucial feeding areas for a rare species, linking habitats without which the overall interest on the site would be threatened by fragmentation, hydrological features on which the habitat depends, or
- other less-diverse habitats without which the most important habitats and species on an ASSI could not be adequately protected.

In cases where less diverse areas are included within the ASSI boundary, they will be subject to the same level of protection as the rest of the site; they are not separate or expendable. (This is **not** the same as "protection zones", in which a landowner may additionally enter into a voluntary agreement on areas outside of the ASSI to further enhance its protection, e.g. by not using sprays close to an ASSI boundary.)

Where semi-natural vegetation is very extensive (e.g. upland moorland), boundaries can be difficult to define, in which case man-made or natural landmarks may be chosen as all or part of an ASSI boundary for ease of identifying and managing the area.

Minor roads, tracks and parking areas may be included within sites, as may un-inhabited buildings and ruins (including some farm buildings), masts, aërials, poles and other minor man-made features. However, where houses and cultivated gardens are surrounded by land meeting the designation criteria they will be excluded from the ASSI boundary.

2.4 Habitat assemblages

The Criteria are generally arranged into common habitat types for ease of reference. However the ASSI selection process takes into account the importance of mosaic habitat communities, including examples of natural zonation of successional stages in vegetation development, and valuable, often mixed habitat corridors, as well as good examples of individual habitats.

Whilst ASSIs can be selected on the basis of one criterion, they can also contain several important habitats and features. This is vital for the conservation of species which are dependent on the presence of several interrelated habitat types, e.g. invertebrates which have both aquatic and terrestrial life stages, and species using marginal habitats. It is also important in the Manx context, where large, homogeneous sites are rare, and the majority of semi-natural vegetation consists of a mix of different but interrelated habitats. Therefore, although most habitat types have special-case selection criteria which are enough to justify ASSI selection in their own right (eg. "best example of species-rich wet grassland on the whole Island"), in general each habitat will be considered in the light of other, complementary site features. The selection criteria have been designed to reflect this.

2.5 Fragmented sites

Some ASSIs may be chosen which consist of several small, non-contiguous sites.

This is appropriate for such habitats as ponds, native woodland fragments, and hay meadows where they, as a group, contribute jointly to the survival of more mobile species in the area. It is recognised that these fragments can be more vulnerable than larger sites, and that linking "corridors" may form an important part of their conservation. Because of this, some fringing and/or linking habitats may need to be included within an ASSI boundary, as outlined in 2.4 above.

2.6 Significance of species.

Selection of sites for the species they support will require an evaluation of species rarity and vulnerability. The presence of species on relevant schedules of the Wildlife Act will be a prime consideration when assessing potential ASSIs.

Wildlife Act schedules will be subject to periodic review. Species protected under the Wildlife Act 1990 are scheduled on the basis of rarity, level of threat to survival, vulnerability to exploitation, "red data book" status, and existing obligations under international conventions. They therefore represent the species of highest nature conservation importance on the Island.

It is recognised that some significant species are not and cannot be protected by designation of their habitat alone. For example, it is not intended to designate bat roosts in people's homes as ASSIs.

2.7 Marine species and habitats

ASSI designation may extend to land covered temporarily or permanently by water. In practice, ASSI boundaries will extend as far as the astronomical low tide line but not beyond, incorporating intertidal habitats but not open sea.

Wider marine site protection will be covered by the selection of Marine Nature Reserves, for which a different set of site selection criteria will be applied. See the companion document: *Guidelines for the Selection of Marine Nature Reserves*.

2.8 Geological and physiographical sites

ASSIs may be selected on the grounds of geological and/or geomorphological importance, in which case separate criteria will be used. In many cases overlap with biological sites is likely. ASSI citations will make clear the main reasons for designation, so that site conservation management can be prioritised accordingly.

See the companion document: *Guidelines for the Selection of Geological and Physiographical ASSIs*.


2.9 Monitoring

A basic principle of management on the majority of ASSIs is that the scientific interest of an area is largely the result of the sympathetic management practices employed by the past and present land managers, and that this interest is often dependent upon the management practices continuing. The success of the ASSI system therefore depends on the co-operation of the owners and occupiers and continuing appropriate land management practices. As land managers they need to know what makes their land of scientific interest. Therefore citations for sites require to be concise, clear and comprehensive statements of the scientific interest of the site. However, because of the complexity of the ecological system described, the citation cannot describe the area in its entirety, nor provide the justification for every single feature or species which is present.

Full information upon which to base site selection decisions is not always available and conservation priorities change with time and wider European ecological change. It is therefore important to continue gathering information, monitoring sites and reviewing the site network. This is an integral part of the site evaluation and selection procedure.

2.10 Revision of site status

Ecological systems are in a state of flux and species' arrival and disappearance usually cannot be managed. Hence, inclusion of new sites as well as de-notification of sites which no longer meet the criteria are options which are open to DAFF. Very careful consideration will be given to the removal of protected status from a site or part of it. There will be a presumption against de-notification, in favour of remediation and management to regain the wildlife interest for which it was originally designated. If this proves impossible, denotification will only be undertaken on sound scientific reasons, after a thorough survey has been carried out to re-find species or other features for which the site was originally designated.



The newly-unveiled board commemorating the designation of Ballaugh Curragh ASSI as a Ramsar Wetland of International Importance – September 2006.

Part 3: Priority Sites Criteria

All ASSIs will be chosen because they have special scientific interest which contributes to the aims and principles for ASSI selection as described in Parts 1 and 2. For example, they might be the only example of a very rare habitat without which the Island's biodiversity would be diminished, or they might represent a priority site for conservation of key habitats and species. Once a site has been shown to support one or more habitats and/or species that merit inclusion within the Island's ASSIs, the next stage is to assess the quality of the site compared with other examples, so that the sites which best meet the aims of the ASSI selection process are identified.

Section 1.6 above lists the Ratcliffe criteria that have long been used for the assessment and selection of protected sites throughout the British Isles. These criteria have been used as a basis for the following Checklist for assessing the quality of a potential ASSI:

3.1 Checklist of Priority Sites Criteria

Size

priority sites:

- the site is an exceptionally large area of an important natural or semi-natural habitat e.g. the largest on the Island or the largest within a distinct region of the Island;
- the site supports an exceptionally large and/or thriving population of an important species (as defined in the Species Criteria);
- the site supports a high proportion of the total area of an important habitat or the total numbers of an important species on the Island and/or in a wider international context.

Location

priority sites:

- the scientific interest of the site is dependent on a rare or unique combination of site-related factors such as geology, aspect, soil type, microclimate, hydrology or altitude. Consequently if the site was damaged or destroyed, the habitat and species communities present would be irreplaceable;
- the site supports habitats or species which are on the very edge of their natural range.

Diversity

priority sites:

- the site contains a high proportion of species typical of the habitat as it is found on the Island in its most favourable condition;
- the site contains a range of semi-natural habitats in close proximity;
- a range of successional stages of habitat development are present on the site;
- the habitats present exhibit a wide range of natural structural diversity.

Rarity, fragility and vulnerability

priority sites:

- the habitats and/or species present are internationally rare or vulnerable to loss, damage or exploitation;
- the habitats and/or species present are rare and/or fragile or otherwise vulnerable to loss, damage or exploitation, on the Isle of Man.

Naturalness and Representativeness

priority sites:

- the site contains all the typical species and assemblages - including stages of succession, subtypes and variations - for which a habitat type is considered scientifically important on the Island;
- the site is an excellent representative of a habitat or species population that forms a distinctive element of Manx biodiversity;
- the site is the only example of a particular habitat sub-type or variation that cannot be protected elsewhere;
- the site represents an excellent example of a mosaic of associated habitats typical of the Manx situation, e.g.

coastal, upland;

- compared with other examples on the Island, the habitat present is notable for its lack of human disturbance, introduced plant or animal species, mechanical damage, litter, agricultural spray drift or other factors which could adversely affect the vegetation structure and/or species composition or community.

Ecological position

priority sites:

- the site forms part of an important, larger ecological unit which would be reduced in value as a whole if the site was damaged or destroyed;
- the site forms a vital part of a sequence of habitats all of which are required in order to conserve a key population of an important species (e.g. semi-aquatic invertebrates).



Patches of coastal grassland at Poyll Vaaish ASSI form part of a series of interrelated coastal habitats

History

priority sites:

- the nature conservation interest of the site is dependent on a rare or unique combination of historical factors such as long-term land use and management patterns;
- the habitats and species present have become established over a very long period of time and consequently represent a finite resource on the Island, as they could not be replaced or substituted in the short or medium term.

Research value

priority sites:

- the site provides the best or only example of a situation where a threatened or declining habitat or species of high nature conservation interest for which there is a research need may effectively be studied;
- the site has one or more features of nature conservation importance that would not ordinarily qualify for ASSI selection, but which are known to be declining or having to adapt due to factors which cannot be prevented, and for which research over the medium or long term is crucial for the success of conservation efforts elsewhere.

3.2 Using the Priority Sites Criteria

The checklist of Priority Sites Criteria is referred to throughout the detailed ASSI Selection Criteria in Part 4, and is applicable to each habitat type and species group. Where there is a choice to be made about which site to designate, the checklist should be used as a guide to site selection as follows:

- **For the very rarest habitats and species, simply fulfilling the minimum requirements as listed in the detailed selection criteria is likely to be enough to warrant selection.**

Example: species-rich neutral grassland is a rare and diverse habitat which is known to have suffered a severe decline throughout the British Isles, therefore all examples of this habitat over a certain minimum size may qualify for selection.

- **All other sites will be selected because they not only fulfil the minimum habitat or species requirements, but also meet at least one of the Priority Sites Criteria.**


Example: unimproved acid grassland is an important wildlife habitat on the Island but is much more widespread than neutral grassland. Examples of this habitat will only be considered if they not only meet the minimum size threshold, but also fit at least one of the Priority Sites Criteria as well, such as being an exceptionally good representative of the type.

- **A site will generally be considered to be of a higher quality than a similar site which meets fewer Priority Sites Criteria, although exceptional cases will be considered on their individual merit.**

Example: of two areas of unimproved acid grassland, both of which are above the minimum size threshold for the habitat, a site which has a high proportion of the total area of acid grassland on the Island is less likely to be selected as an ASSI than a similar site which not only has a high proportion of the total habitat area, but also fits several other Priority Sites Criteria -such as supporting species which are on the edge of their natural range, and forming part of an important, larger ecological unit.

- **Site selection may be subject to further prioritisation based on a range of factors depending on the species and habitats in question. Where relevant, this will be explained in the detailed selection criteria for each habitat type or species population.**

Example: for some grassland habitats, examples will be selected that fulfil minimum size criteria and fit at least one Priority Site Criterion, with further priority being given to sites that form part of a larger habitat mosaic.



Common blenny or Shanny, Liphophrys pholis, in rockpool at Dhoon Glen ASSI – a prime example of a priority site for its wide range of interrelated habitats



Cover pictures show protected species found on ASSIs in the Isle of Man:

Lesser mottled grasshopper *Stenobothrus stigmaticus*

Pyramidal orchid *Anacamptis pyramidalis*

Lapwing *Vanellus vanellus*

Isle of Man Cabbage *Coincya monensis* subsp. *monensis*

The information in this document can be provided in large print, on request.

DAFF

January 2008

©2003-2008 Isle of Man Government Department of Agriculture, Fisheries and Forestry
The contents of this document are the property of the Department of Agriculture, Fisheries and Forestry, and should not be copied without its permission.

Guidelines for the selection of biological Areas of Special Scientific Interest (ASSIs) on the Isle of Man

Volume 2: Detailed habitat and species criteria



**Isle of Man
Government**

Reiltpys Ellen Vaanin

Volume 2

Detailed habitat and species criteria

Introduction

The following chapters describe natural and semi-natural vegetation communities on the Isle of Man, their relationship to established methods of classification such as the National Vegetation Classification, characteristics that will be taken into consideration when assessing a potential ASSI, and minimum criteria for selection. Where a habitat is rare or locally distributed on the Island, the most important known sites may be listed individually.

The habitat selection criteria are followed by criteria for site selection based on the presence of important species from a range of plant and animal groups. It should be noted that in many cases a site will qualify on both habitat and species grounds.

Both the habitat and species chapters contain an indication of the current level of knowledge about the subject. Where knowledge is limited, e.g. for some invertebrate and lower plant groups, there is an assessment of research needs and priority level.

Where there is a choice of several sites with the same important habitats or species present and a choice needs to be made about which site to select, the relative quality of the sites will be assessed using the Priority Sites Criteria in Part 3 of Volume 1.

Chapter index to Detailed Guidelines

<i>Chapter</i>		<i>pages:</i>
4.1	Woodland and scrub	1
4.2	Lowland grasslands	22
4.3	Lowland heathland	43
4.4	Bogs, fens and flushes	54
4.5	Open water and swamp	65
4.6	Coastal habitats	78
4.7	Uplands	104
4.8	Flora	126
4.9	Avifauna (birds)	127
4.10	Fauna (all other animals)	133

Please note that in order to avoid duplication of habitat accounts some chapters – e.g. Upland habitats - may refer to sections of other chapters.

Contents

Introduction	i
Part 4: Detailed selection guidelines	1
4.1 Woodland and scrub	1
4.1.1 Semi-natural broadleaved woodland	6
Minimum selection criteria for semi-natural broadleaved woodland	7
Table 4.1.a: Broad categories of semi-natural woodland on the Isle of Man	8
4.1.2 Plantations	10
Table 4.1.b: Broad categories of plantation on the Isle of Man	11
Minimum selection criteria for plantations	12
4.1.3 Dry scrub	13
Table 4.1.c: Broad categories of dry scrub on the Isle of Man	14
Minimum selection criteria for dry scrub	16
4.1.4 Curragh	16
Minimum selection criteria for curragh	17
Appendices to Chapter 4.1: Diagnostic species lists	18
Appendix A: Tree and shrub species thought to be native to the Isle of Man	18
Appendix B: Woodland ground flora species associated with semi-natural woodland	19
Appendix C: Plants not of native origin which may indicate unfavourable condition	21
4.2 Lowland grasslands	22
4.2.1 Acid grasslands	24
Minimum selection criteria for lowland acid grassland	25
Table 4.2.a: Broad categories of acid grassland on the Isle of Man	26
4.2.2 Neutral grasslands	27
Minimum selection criteria for neutral grassland	28
Table 4.2.b: Broad categories of neutral grassland on the Isle of Man	29
4.2.3 Calcareous grasslands	31
Table 4.2.c: Broad categories of calcareous grassland on the Isle of Man	32
Minimum selection criteria for calcareous grassland	33
4.2.4 Coastal grassland	33
Table 4.2.d: Broad categories of coastal grassland on the Isle of Man	35
Minimum selection criteria for coastal grassland	36
Table 4.2e: Vascular plants characteristic of semi-natural grassland on the Isle of Man	37
4.3 Lowland heathland	43
4.3.1 Dry lowland heathland	48
Table 4.3.a: Broad categories of dry lowland heathland on the Isle of Man	49
Minimum selection criteria for dry lowland heathland	49
4.3.2 Wet lowland heathland	50
Table 4.3.b: Broad categories of wet lowland heathland on the Isle of Man	50
Minimum selection criteria for wet lowland heathland	51
4.3.3 Coastal heathland	51
Table 4.3.c: Broad categories of coastal heath on the Isle of Man	52
Minimum selection criteria for coastal heathland	53
4.4 Bogs, fens and flushes	54
4.4.1 Bog	56
Table 4.4a: Broad categories of bog on the Isle of Man	59
Minimum selection criteria for bogs	59
4.4.2 Valley mires and flushes	60
Minimum selection criteria for valley mires and flushes	61
4.4.3 Basin mires	61
Description and Distribution	61
Minimum selection criteria for basin mires	62
4.4.4 Fen meadows	62
Table 4.4c: Broad categories of fen meadow on the Isle of Man	63
Minimum selection criteria for fen meadows	64
4.5 Open water and swamp	65
4.5.1 Still water and swamp	68
Table 4.5.a: Broad categories of still water and swamp habitats on the Isle of Man	70
Minimum selection criteria for still water habitats	74

4.5.2 Ditch systems.....	75
Minimum selection criteria for ditch systems	76
4.5.3 Flowing water	76
Minimum selection criteria for flowing water	77
4.6 Coastal habitats	78
4.6.1 Intertidal habitats.....	80
4.6.2 Strandlines and shingle	80
Minimum selection criteria for strandlines and shingle	82
Table 4.6b: Broad categories of strandlines and shingle habitats on the Isle of Man	83
4.6.3 Dunes.....	85
Minimum selection criteria for dunes.....	86
Table 4.6c: Broad categories of dune habitats on the Isle of Man.....	87
4.6.4 Saltmarsh.....	89
Table 4.6.d: Broad categories of saltmarsh on the Isle of Man	92
Minimum selection criteria for saltmarsh	95
4.6.5 Brackish pools.....	95
Minimum criteria for selection	97
4.6.6 Sea cliffs, caves and slopes	97
Table 4.6.e: Broad categories of maritime cliff vegetation on the Isle of Man.....	100
Minimum criteria for selection	103
4.7 Uplands.....	104
Table 4.7.a – Manx upland conservation “hotspots”.....	108
4.7.1 Acid grassland.....	111
Minimum selection criteria for upland acid grassland	112
4.7.2 Upland heathland.....	112
Minimum selection criteria for upland heathland	116
4.7.3 Blanket bog.....	117
Minimum selection criteria for blanket bog.....	120
Table 4.7.b: Broad categories of upland habitat on the Isle of Man	121
Recommendations for selection of upland ASSIs	125
4.8 Flora – sites of exceptional importance for plant conservation	126
4.9 Avifauna – sites of exceptional importance for birds.....	127
Minimum selection criteria for sites of high importance for birds	129
Table 4.9a: Criterion 4 scores for habitats	131
4.10 Fauna – sites of exceptional importance for animals	133
Part 5: Bibliography and References	134

Parts 1-3: Background, principles and Priority Sites Criteria, are contained within **Volume 1.**



Six-spot burnet moths *Zygaena filipendulae*

Part 4: Detailed selection guidelines

4.1 Woodland and scrub

Description

As with many offshore islands around the British and Irish coasts, the Isle of Man has no natural tree cover for the majority of its land surface. It is thought that most of the Manx native mature oak woodland was already cut down and exploited by the late Iron Age (Garrad 1972; Robinson & McCarroll 1990). Whilst it is possible for young trees and shrubs to regenerate naturally on ungrazed sites, the high levels of exposure to wind and sea spray tend to militate against the redevelopment of mature semi-natural woodland in any but the most sheltered valley locations. However, research into the ancient pollen and seed deposits in Manx soils has revealed that at one time the Isle of Man was almost entirely covered by woodland. Archaeological evidence of birch, hazel, juniper, willows, sessile oak, wych elm, Scots pine, common alder and ash has been found at varying abundances in the soil strata, indicating that different species have tended to dominate at certain periods in time: hazel, for example, being apparently more prevalent in the past than at present (Allen 1984; Robinson & McCarroll 1990).

Whilst pollen records indicate which Manx trees can reasonably be thought of as **native species**, the usefulness of trees to the human inhabitants of the Island has led to much importation, of “native” trees as well as exotics. Hence it can be very hard to assess whether a population of oak, for example, is truly of native origin, rather than from elsewhere in the British Isles or even further afield. The most confusing example of this is the case of Scots pine, which is present on the Island only as an imported species, but which was clearly once native and is still quite capable of setting seed in the wild and spreading from planted areas to form part of a semi-natural, regenerating woodland environment. Likewise, the woodland **ground flora** on the Island is by no means free from human modification, with many of the more ornamental plants either introduced to the Glens for decoration, escaped from adjacent gardens, or “helped” to spread from their preferred native habitats (Allen 1984). In general, though, the more well-established wooded valleys, glens and plantations support a moderately diverse ground flora similar to some of the semi-natural oak and ash woodland types described in the National Vegetation Classification.

Woodland and scrub are here taken to include all communities dominated by woody vegetation, except for dwarf shrub heath (such as heather- and Manx gorse-dominated habitats), which will be covered in the chapter on Heathland. **Veteran tree and dead wood habitats** will generally be most relevant to site selection on species criteria (e.g. for rare beetles), but they add value to woodland habitats in their own right; hence they are also considered here.



The distinctive, drooping native bluebell *Hyacinthoides non-scripta*, at Glen Rushen ASSI

Assessing woodland and scrub

On the Isle of Man, woodland and scrub vegetation is a broad habitat category which has probably been more drastically affected by human activities, both ancient and modern, than any other (see NCC 1975; Sayle *et al*, 1995; Allen, 1984). This, coupled with the effects of the Island's climate and maritime influences, has led to difficulties in comparing Manx woodland habitats with those in neighbouring countries. Substantial areas of semi-natural ancient woodland such as those in Wales or Scotland do not occur on the Isle of Man, and in some situations scrub communities appear to form a successional climax here. For these reasons, ground flora composition is likely to play an even more important role in the site assessment process than elsewhere, hence the most diverse semi-natural scrub communities will be given a status on a par with that of woodland.

All the Priority Sites Criteria listed in Part 3 of this document are relevant to woodland assessment; of these, the following are especially important when comparing similar woodland vegetation communities with a view to site designation:

Size

The overriding aim is to protect the largest areas available of the major woodland and scrub habitat types so that the largest possible populations of the species associated with each type can be protected. This presumption takes into account the fact that woodland ground flora is particularly vulnerable to "edge effects" on small, narrow and fragmented sites. This aim may however need to be qualified if the larger habitat stands are also the most disturbed, and where smaller examples are needed to compliment the features found in the largest areas (some woodland types, for example, are naturally limited to small areas). Sites with a mixture of vegetation communities, each of which occupy only a small area, are important in a different way because they may demonstrate the relationship between species distribution and historic or ecological factors. They may also be particularly species rich. Hence, although larger sites are desirable, smaller woodlands should not be ruled out without considering all the ecological factors involved.

Diversity

Within any given woodland, clearings, rides, variable topographical features (such as steep ground, rock outcrops or wet ground), variations in drainage and presence of veteran trees and/or dead wood will all increase the importance of a site through providing greater diversity. Likewise, woods which adjoin or are in mosaic with other habitats tend to be more valuable than those sharply abutted to arable or improved grassland, and diffuse open or bushy edges are better than sharp boundaries.

British woodlands generally have fewer species of trees, shrubs and herbaceous plants than their continental counterparts and as a whole have suffered more loss and fragmentation through human activity (Ratcliffe, 1977). This situation is even more extreme on the Island, where native woodland would have been even more impoverished in species, and almost all the original habitat had been lost or fragmented by the sixteenth century (Allen, 1984). All references to species-richness should be tempered by consideration of these factors.

In all cases, special features such as populations of uncommon species, well-developed scrub communities, extensive moss carpets or fern assemblages, boulder scree, and woodland ponds will be considered positive attributes. This includes small meadows or bits of rough grassland, remnant heath or moorland and any other features which could be of high importance for the overall invertebrate diversity of a site. Invasive introduced species, many of which are capable of severely reducing the diversity of both plants and invertebrates, should be treated as negative attributes.

In order to help with assessing the diversity and naturalness of woodland and scrub, the **Appendices** at the end of this Chapter provide a list of tree and ground flora species thought to be indicative of Manx woodlands with a high nature conservation value, along with some common introduced/escaped alien species which may indicate a modified flora and reduced overall diversity.

Rarity, fragility and vulnerability

The minimum criteria listed in the relevant section of this chapter (sections 4.1.1-4.1.4) each take into account the fact that some types of woodland are naturally more fragile and vulnerable to damage than others. In the case of the rare semi-natural woodland communities, **all** examples will qualify for selection as ASSIs provided that they are of sufficient size and fulfil the minimum criteria. In most cases, however, candidate sites will be selected from the best good examples of each woodland and scrub type. Sites will be considered to be "good examples" where they not only fulfil all the minimum requirements, but also one or more of the Priority Sites Criteria listed in Part 3.

Where rare species occur in a woodland, the site may qualify on species grounds alone, or may be considered a particularly good example of the habitat type. This applies to rare or locally distinctive woodland habitat features as well as individual species, particularly where the Isle of Man has a high proportion of such features that lend a distinctive character.

Local rarity is not the only category: for some species and features, the Island may have plenty of examples of a type that is rare or threatened elsewhere. In this case there is wider responsibility for protection. In woodlands this applies particularly to the Island's elm population, which has largely escaped the ravages of Dutch elm disease that have decimated elm populations elsewhere in the British Isles, and consequently is now of special importance.

Naturalness

It should be noted that, as with grasslands, very few woodland and scrub samples were taken from the Isle of Man to contribute to the original research that formed the basis of the National Vegetation Classification. Perhaps because of this, Manx vegetation communities do not always fit well within NVC categories. This does not mean that the NVC is worthless in the Manx context, but it has led to a situation where the NVC alone is not usually sufficient for assessing the nature conservation importance of a site. This is even more the case with woodlands than with grasslands (see **Section 1.3.4**, above, regarding the use of the NVC in the Manx context). In view of this, proportions of species likely to be native will be as important a defining characteristic as NVC types. Broadly speaking the higher the proportion of native species then the greater the nature conservation value. For lists of tree and ground flora species thought to indicate naturalness in a woodland, see Appendices A and B at the end of this Chapter.

Alien species in woodlands may threaten the native flora by displacement and/or hybridisation, resulting in a less diverse ecosystem overall. Commonly-occurring introduced woodland species are listed in Appendix C at the end of this Chapter – although it should be noted that the overall list of potential introductions to Manx woodlands is much greater. Such species may not cause a problem in plantation settings – such as the Manx National Glens – but in general, abundant and spreading exotic species, along with rubbish tips, grossly polluted streams, heavy grazing and forestry operations which eliminate over-mature timber, should all be considered negative attributes, to be excluded from selected areas wherever possible. Large blocks of coniferous plantation should also be excluded, the exceptions being:

- compartments within a larger site that is being considered for selection, where the rides and/or ground flora strongly reflect the surrounding, more diverse, habitats;
- small conifer stands within larger, more diverse woodlands, where exclusion of the stands would lead to an impractical boundary on the ground or for subsequent management;
- any areas which are vital for the survival of important species and hence qualify under the species criteria.

Position in ecological unit

Because most woodland tends to have quite distinct boundaries, it is usual to notify more or less complete units (NCC, 1989). If boundaries within woods are required, they should follow well-defined permanent features such as paths or streams. Major tree species changes (such as the edge of a coniferous area) may also be used as designation boundaries providing they fall within well-defined compartments – in which case care should be taken to ensure that only long-term compartment boundaries are chosen. For old sites, boundary banks should be included as they may have ancient trees and special flora.

Consideration should be given to the inclusion of fringes of recently-developed semi-natural woodland or scrub outside the main area of woodland. Such features provide additional diversity as well as the opportunity to examine woodland processes such as species colonisation. Fringe habitats also allow for the possibility of natural extension of woodland by gradual regeneration along existing margins. This is of particular value in or adjacent to upland areas (see Chapter 4.7 for details of upland site selection).

Large blocks of semi-natural vegetation adjacent to a wood should normally be considered on their own criteria. The exceptions are mosaic sites with habitat mixtures where no individual element reaches the minimum standard for ASSI selection, but which together form a valuable wildlife site. Each site can only be treated on its own merits, and where an atypical site is chosen, full reasons for the choice of site and boundary will be given.

Where areas of woodland are much smaller than the minimum recommended area for ASSI selection in their own right, but are attached to larger grassland or heath habitats, they may still be designated under the relevant grassland or heathland criteria as a valuable mosaic element.



Hart's-tongue fern *Phyllitis scolopendrium*, at Glen Maye ASSI

Woodland and scrub types

The types of woodland stand structure typically associated with traditional management of UK and Irish woodlands - wood pasture, coppice and so forth - are not usually evident on the Isle of Man, and cannot be used as a method of categorisation. There are, however, a range of broad woodland and scrub types on the Island, according to age, structure, ground flora composition and degree of human influence. For example, the DAFF Phase 1 Habitat Survey recognises 10 different “woody” categories, divided according to whether the vegetation is woodland or scrub, dense or scattered, broadleaved or coniferous (Sayle *et al*, 1995). For the purpose of the Criteria, woodland and scrub on the Isle of Man have been grouped more broadly, as follows:

Semi-natural broadleaved woodland

This consists of fragments of woodland/groups of trees which are believed to have grown or regenerated from native (post- Ice Age) sources (distinguished if possible through DNA analysis), and areas of semi-natural woodland which may have regenerated from planted and native sources. Truly semi-natural woodland with negligible modification from introduced species and human activities is rare; such habitats tend to occur within the sheltered confines of steep river valleys, or in a “dwarf” form on sea cliffs.

Plantations

Well-established woodland most commonly occurs in the **Manx National Glens** – most of which result from the Victorian desire for attractive outdoor leisure areas, but which may also support a range of naturally-occurring tree, shrub and ground flora species, some of which are likely to be of native origin. All Manx woodland has some semi-natural vegetation, if only in the ground layer. The best examples have a high proportion of native species in tree, shrub and ground layers.

Other plantations that may be of value for wildlife include areas of conifer or mixed plantation which are sufficiently old and well-established to have developed a diverse woodland ecosystem. Even less diverse conifer plantations may sometimes provide valuable habitat for particular groups, notably birds of prey. Forest structure may be more important than species.

Dry scrub

Whilst there are few unmodified tree habitats on the Island, semi-natural scrub is plentiful and can form a diverse habitat that approaches semi-natural woodland in its structural characteristics and species composition. Examples include bracken- and gorse-dominated sites on the rare occasions when they are allowed to develop into regenerating woodland, and coastal cliffs, where exposed conditions can result in a mix of typically short woody species alongside trees growing at scrub height, such as stunted sessile oaks, birch and rowan.

Curragh

The most distinctive Manx scrub habitat is willow-dominated scrub on wet, peaty soils, known generally as curragh. The majority of curragh has a relatively low, scrubby canopy, but the most well-established examples may have a species composition similar to wet woodland, with taller trees such as ash, alder and aspen. This represents an internationally restricted habitat type. Curragh is strongly associated with other peatland and open water habitats, e.g. fen and ditch systems, and its value as part of a mosaic of related habitats should always be taken into account.

Several examples of each type should be selected in order to account for the geographical variation in structure, floristic composition, animal assemblages and habitat associations that characterise Manx woodland and scrub. The higher the variability of plant and animal communities of any woodland type on the Island, the more examples should be considered. The same applies where the type is nationally rare or particularly well-represented on the Island as opposed to elsewhere. The overall range of sites selected should also encompass all the important special features which occur in that type of woodland or scrub.

The description, classification, status and distribution of each of the main woodland and scrub types are each treated separately here for reasons of clarity and ease of reference. However, when assessing a site in the field it should be borne in mind that this is an artificial separation. Sites which do not “fit” the broad categories are not necessarily poor examples of species-rich communities, and may be as important for wildlife as more “typical” sites.

Sections 4.1.1-4.1.4, below, give detailed descriptions, distribution and status accounts for each of the main types of woodland and scrub on the Island, alongside the National Vegetation Classification (NVC) categories they most closely resemble, and minimum criteria for selection.

4.1.1 Semi-natural broadleaved woodland

Description and Distribution

The natural climax woodland of the Manx lowlands is hard to discern now that so few semi-natural woods remain, but it seems likely that for most of the drier soils of the Island sessile oakwood is the natural dominant. Ash, wych elm and hazel also feature strongly in lowland woodland, frequently alongside sycamore (a non-native and often invasive species thought to have been brought to the Island “some time before 1760” (Allen, 1984)). Examples include relict oak-and-hazel woods in lower Glen Maye and upper Glen Auldyn. On higher ground, rowan is a common pioneer and a strong element of semi-natural regenerating woodland. Wet ground typically supports willows (“sallies”), birch, ash and alder. Whilst native woodland regenerates well on the more sheltered areas of the Island, it is virtually absent from the more extreme environments – the wettest and most exposed – which tend to support a climax of curragh or coastal scrub rather than tall woodland. These habitats are covered described separately in sections 4.1.3 and 4.1.4, below.

The Island’s small remnants of semi-natural woodland are located mainly along the rivers and streams in small steep-sided valleys, often in comparatively inaccessible locations where surviving ancient woodland species could have persisted, out of reach of human clearance activities and most grazing animals. This woodland tends to be dominated by ash, wych elm and sycamore. There are also a number of isolated stands of sessile oak *Quercus petraea*, usually on steep coastal slopes beyond reach of most grazing animals. These have long been thought to be descended from native stock, and recent chloroplast DNA studies tend to confirm this hypothesis (Cottrell *et al*, 2002). It is also possible that “abbey land” oak woodland in Narradale, where evidence of large old oak stools can still be seen, has survived without ever being totally cleared (Garrahd, *pers comm*).

Semi-natural broadleaved woodland status

The 1994 Ecological Survey (based on Phase 1 Habitat Survey data) found 167ha of semi-natural broadleaved woodland on the Island – 3.6% of the Island’s total woodland, 1% of all the semi-natural habitats recorded, and less than 0.3% of the total Island land use. The largest semi-natural broadleaved woodland site found was just 9ha, and the mean average size was around half a hectare (Sayle *et al*, 1995). This makes it one of the rarest and most vulnerable semi-natural Manx habitats. For this reason, the Manx Wildlife Sites Handbook (Keehan, 1999) states that *all* relict **oak wood**, and the majority of all other semi-natural broadleaved woodland, should be considered for Wildlife Site selection.

In the European context, Manx native woodlands are examples of British woodland types which are themselves western outliers of the larger distribution pattern of continental European forest. Manx woodlands are particularly characteristic of Western Britain, with its mild, damp, oceanic climate. They are typified by species-rich and luxuriant communities of ferns, mosses and liverworts, with similarly oceanic lichen flora. This kind of woodland is very restricted in distribution, featuring in both the UK Biodiversity Action Plan (which lists Upland mixed ashwood, Wet woodland and Upland oak wood as Priority Habitats) and Annex 1 of the EU Habitats Directive (which includes both Dry oak-dominated woodland and Western acidic oak woodland).

Veteran trees and **dead wood** have both been recognised as having a high conservation status. A recent report states that European woodlands have around 5% of the volume of dead wood that might be expected under natural conditions, making it a critically low habitat resource – dead wood specialist species being the single largest group of threatened plants, animals and fungi in Europe. The report recommends that living veteran trees, standing dead trees, lying dead trees and dead-wood-rich litter are all elements of a healthy woodland which need to be encouraged and increased in volume in order to safeguard woodland biodiversity, especially in protected areas (WWF, October 2004).

Classification of semi-natural broadleaved woodland

Semi-natural broadleaved woodland on the Island may be classified broadly as outlined in **Table 4.1.a**.

Minimum selection criteria for semi-natural broadleaved woodland

Wet woodland

All examples of mature semi-natural wet woodland which fulfil at least one of the Priority Sites Criteria listed in Part 3 should be considered for selection, with special priority given to sites with the greatest size, naturalness and diversity, or best position within an ecological unit.

Lowland oak and ash woodland

- All examples of lowland oak and/or ash woodland over 2ha in extent should be considered for selection, even if their conservation condition is currently less than favourable.
- Smaller examples of lowland oak and/or ash woodland should *all* be considered for selection if they fulfil at least one of the Priority Sites Criteria listed in Part 3.

Upland oak and ash woodland

- All examples of upland oak woodland over 2ha in extent should be considered for selection, even if their conservation condition is currently less than favourable, whether or not they are part of a larger upland site designation.
- Smaller examples of upland oak woodland should *all* be considered for selection if they fulfil at least one of the Priority Sites Criteria listed in Part 3.

Upland woodland will usually be assessed according to the Upland Habitats criteria detailed in Chapter 4.7.



Dryad's saddle fungus *Polyporus squamosus* is one of the many organisms which benefit from veteran and dead trees; the large fruiting bodies form a micro-habitat for invertebrates

Table 4.1.a: Broad categories of semi-natural woodland on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (see Rodwell, 1992 ¹)	Rarities present?	Protected species present?
<p>Wet woodland Very restricted in distribution. Usually found in association with curragh, and containing some of the same elements and NVC types. Occasionally occurs on upland fringes. Birch-dominated examples are notable for the occasional presence of <i>B. pubescens</i> ssp. <i>tortuosa</i> – a small birch variety with fragrant foliage.</p>	Not always botanically diverse, but may support diverse bird and invertebrate populations. Very species rich when found in association with related habitat eg. Open water, mire.	<p>Mature W4 – <i>Betula pubescens</i>-<i>Molinia caerulea</i> woodland No Island NVC sample</p> <p>W7 – <i>Alnus glutinosa</i>-<i>Fraxinus excelsior</i>-<i>Lysimachia nemorum</i> woodland No Island NVC sample</p>	Rare birds and invertebrates a possibility; rare plants possible in diverse examples	Yes. Several Schedule 1 birds. Noted for providing winter hen harrier roost
<p>Lowland oak and ash woodland Oak and ash dominated woodland with a range of other native trees and shrubs, often indeterminate with upland woodland types and featuring sessile, rather than English, oak (although the two species are not always well-defined on the Island (Allen 1984)).</p>	Yes	<p>W8 – <i>Fraxinus excelsior</i>-<i>Acer campestre</i>-<i>Mercurialis perennis</i> (ash/field maple/dog's mercury) woodland Note that both <i>A. campestre</i> and <i>M. perennis</i> are absent from the Island as native spp. However, ash woodland on more base-rich Manx soils has other similarities to W8, mostly in the ground flora composition Two Island NVC samples.</p> <p>W9 – <i>Fraxinus excelsior</i>-<i>Sorbus aucuparia</i>-<i>Mercurialis perennis</i> (ash/rowan/dog's mercury) woodland Note: <i>M. perennis</i> is absent from the Island as a native species; however, other aspects of W9 woodlands are clearly evident in some of the ash/rowan/wych elm communities in steep stream valleys and on the edge of the uplands No Island NVC sample</p> <p>W10 – <i>Quercus robur</i>-<i>Pteridium aquilinum</i>-<i>Rubus fruticosus</i> (English oak/bracken/bramble) woodland One Island NVC sample</p>	Strong possibility of plant, fungi, invertebrate, mammal and bird rarities	Strong possibility of Schedule 1 birds, Schedule 7 plants and bats

Type and Description	Species rich?	Closest NVC categories (see Rodwell, 1992 ¹)	Rarities present?	Protected species present?
<p>Upland oak and birch woodland</p> <p>Diverse, often rather open woodland usually restricted to stream valleys and steep-sided gills. Often strongly associated with other upland habitats, and notable for lush fern and moss growth. All but the largest examples should be considered under the Upland habitats Criteria in Chapter 4.7. Note: There are places where birch will regenerate on valley sides when grazing is removed; this may be treated as a scrub community (see below) or as an immature upland oak and birch woodland, depending on the stage of development and the position in the surrounding ecological unit.</p>	Yes, although sometimes less diverse than lowland oak woodland	<p>W9 (see above)</p> <p>W11 – <i>Quercus petraea</i>-<i>Betula pubescens</i>-<i>Oxalis acetosella</i> (sessile oak/downy birch/wood sorrel) woodland One Island NVC sample</p> <p>W17 – <i>Quercus petraea</i>-<i>Betula pubescens</i>-<i>Dicranum majus</i> (sessile oak/downy birch/great fork moss) woodland No Island NVC sample</p>	Possibility of plant, fungi, invertebrate, mammal and bird rarities	<p>Possibility of Schedule 1 birds, bats and Schedule 7 plants eg. Beech fern <i>Phegopteris connectilis</i>, Wilson's filmy fern <i>Hymenophyllum wilsonii</i>.</p>

4.1.2 Plantations

Description and Distribution

Many of the largest areas of broadleaved woodland are **Manx National Glens** (17 in total). These were planted largely in the latter part of the 19th century and usually contain mature stands of trees. The process of re-developing the Island's woodland began in the 18th century when landowners, most notably Bishop Wilson, began planting in the many glens and in private gardens. This continued into the Victorian period and the heyday of Manx tourism, when glens were planted with a wider variety of ornamental trees and shrubs and opened to the public as "pleasure grounds". The most commonly planted species are beech, wych elm, sycamore, and some oak, sweet chestnut and horse chestnut (Garrahd 1972 & 2003). Such areas of broadleaved woodland can be of high value for wildlife.

Parkland is a special category of planted area that is not noted for its ground flora, but which can provide excellent invertebrate habitat due to the presence of large and over-mature specimen trees. This advantage is further increased by sympathetic management practices such as leaving large dead trees in situ, or keeping felled wood on site for the benefit of dead-wood invertebrates. There is very little classic parkland on the Isle of Man, although some estates have woodland landscaping which allows for the retention of large specimen trees. Any **veteran trees** are of value for wildlife, and the presence of such trees will be a material factor in ASSI selection for any site, whether woodland-dominated or not.

Well-established **beech woodland** is rare on the Island and not a native habitat, but it can have exceptional fungus diversity, and as such could fall within Priority Sites Criteria for particular species groups.

Much of the more recent broadleaved planting is composed of mixed broadleaved trees including wild cherry and other more decorative species. Such areas vary in ecological diversity depending on location and proximity of more diverse woodland. Large areas of young, native tree species adjacent to existing mature broadleaved woodland may offer bird and invertebrate territory even though the ground flora under the plantation is not particularly diverse.

Conifer plantation, mostly dating from the 20th century, accounts for well over half of all the woodland and scrub on the Island (*Sayle et al*, 1995). Dark, closely-spaced spruce plantations, which have little light reaching the field layer and a dense carpet of needles, tend to have the least ecological diversity. Plantations that have gaps or "failed" areas may however offer good localised heath or bog habitats, often sheltered by the surrounding trees. More mixed plantations, especially those with a proportion of larch and mixed scrub margins, tend to attract more birds and invertebrates, along with a few ground flora species that can take advantage of the slightly better light levels. Larch and pine plantations appear to attract better fungal communities than spruce (Hodgetts, 1996). All well-established conifer plantations can offer bird habitat, albeit for a more limited range of species than broadleaved plantations. In a few cases, conifer plantations may be significant nesting sites for rare species, notably birds of prey (Cullen & Jennings, 1996). Areas of young or failed conifer plantation can provide valuable bird habitat for hen harrier *Circus cyaneus* and short-eared owl *Asio flammeus*, where the canopy remains very open, providing shelter and perches.

Classification of plantations

Plantations are usually classified simply as broadleaved, mixed or coniferous. These categories do not relate closely to the National Vegetation Classification, but it should be noted that in the case of well-established broadleaved plantations the ground flora is likely to have features and species in common with semi-natural woodland types as listed in **Table 4.1.a**, above. For the purpose of the Criteria, species-rich broadleaved plantations are considered as a separate category to other broadleaved and mixed plantations, as detailed in **Table 4.1.b**.

Table 4.1.b: Broad categories of plantation on the Isle of Man

Type and Description	Species rich?	Closest NVC category (see Rodwell, 1992 ¹)	Rarities present?	Protected species present?
Species-rich broadleaved plantation Characterised by at least three of the following attributes: <ul style="list-style-type: none"> • mature canopy and varied understory structure consisting predominantly of native species (listed in Appendix A); • a range of Manx native ground flora species (listed in Appendix B); • absence or minimal presence of introduced species (listed in Appendix C); • other features indicative of a thriving woodland ecosystem, eg. good fungal assemblages 	Yes	May have ground flora species in common with any of the NVC categories listed in Table 4.1a	May support rare plants, invertebrates, bats and birds	Possibly, particularly bird and invertebrate species
Parkland Collections of large mature, over-mature and veteran trees dating from Victorian or earlier planting	Possibly rich in certain species groups		May support rare invertebrates, lichens and fungi	May support protected birds and invertebrates
Other broadleaved and mixed plantation Recently replanted conifer blocks, amenity planting areas etc.	Not usually		Not usually	May support protected birds
Conifer plantation	No		Not usually, although rare birds such as crossbill may use plantations	May support protected birds

Plantation status

The majority of plantations on the Isle of Man consist of structured compartments of commercial conifers such as larch, lodgepole pine, Scots pine, Sitka spruce and Western red cedar. The 1994 Ecological Survey also shows 1.11 % of the Island to have broadleaved and mixed plantation – a small proportion compared with the quantity of conifer plantation, but still more than three times the amount of semi-natural broadleaved woodland (Sayle *et al*, 1995).

Well-established broadleaved plantations such as the more diverse Manx National Glens do not attract the same attention from conservationists as semi-natural woodland, but are recognised nevertheless as providing a good habitat for woodland birds, mammals and invertebrates. They may also form a reservoir for native woodland ground flora species and regenerating native trees and shrubs, which, together with the more ornamental planted species, form a comparatively diverse woodland ecosystem. An example of this is the presence of good populations of woodland fungi in glens such as Glen Helen and Dhoon Glen, and plantations such as the Montpelier beech wood (Garrahd, 1972). Planted glens also have the great advantage of accessibility and public appeal, making them especially well-suited to field trips and environmental education. This combination of valuable features has led to the protection of Manx glens as a national asset, and as such they have a higher status in terms of local recognition and protection than most other woodland on the Island.

Parkland trees are often protected for their great visual importance by registering them under the Tree Preservation Act 1993. Other areas of broadleaved plantation attract less protection, and their conservation status will depend to a large extent on their position within a more diverse ecological unit. In the wider context, the significance of parkland trees for invertebrate communities in particular has been recognised in the UK Biodiversity Action Plan, where Lowland wood pasture and Parkland are Priority Habitats.

Stands of wych elm in both glen and parkland settings have a special importance, as the Isle of Man as a whole now has one of the best collections of elm in the British Isles, most of which is likely to be of native British stock (see <http://www.gov.im/daff/crops/trees/elm/welcome.xml>). This is chiefly due to the relatively low incidence of Dutch elm disease on the Island, compared with e.g. England, where native elm species have been devastated by the disease. We therefore have a special responsibility to ensure that this threatened reservoir of biodiversity is protected.

Conifer plantations similar to those on the Island are to be found throughout commercial forestry areas in the northern hemisphere, with large areas of Scotland, Wales and to a lesser extent England all having plantations of similar structure and composition. Birds such as siskins, redpolls and goldcrests benefit from conifer plantations, and some birds of prey may take advantage of the nesting vantage points they provide. Spruce and pine plantations may also attract rare crossbills (Cullen & Jennings, 1986). In general, though, these habitats are not noted for their biological diversity and their status as wildlife areas is usually linked only to the presence of particular rare species, or their useful accessibility for public access and educational visits.

Minimum selection criteria for plantations

Species-rich broad-leaved plantation (as defined in Table 4.1b)

Areas of species-rich broad-leaved plantation of 2 or more hectares in extent will be considered for selection where they fulfil one or more of the Priority Sites Criteria listed in Part 3; priority will be given to sites which fulfil several Priority Sites Criteria and form part of a larger woodland unit which includes semi-natural broadleaved woodland.

Parkland

There is currently insufficient data to assess the importance of parkland on the Island for the conservation of rare dead wood and mature tree species. Parkland will usually be selected only where it forms part of, or lies adjacent to, an area of habitat that merits designation as a whole, *or* on individual species criteria where it is known to be a vital habitat for rare invertebrates or other groups.

However, the designation of an exceptionally large and/or diverse parkland tree collection as a habitat in its own right should not be ruled out, if it can be shown to fulfil one or more of the Priority Sites Criteria listed in Part 3.

Other broadleaved and mixed plantation

Other broadleaved and mixed plantations will usually be selected only where they form part of a broadleaved woodland that merits designation as a whole, or on individual species criteria where they form a vital habitat for rare birds or other groups.

Conifer plantation

Conifer plantations will usually be selected only where they form part of a broadleaved woodland or other (e.g. heathland) habitat that merits designation as a whole (eg at the Ayres), or on individual species criteria where they form a vital habitat for rare birds or other groups.

4.1.3 Dry scrub

Description and Distribution

Scrub is a widespread habitat on the drier soils throughout the Isle of Man, providing extensive areas of cover for common bird and mammal species. Typically, dry scrub results from encroachment of hedging species, such as hawthorn, blackthorn and European gorse, onto ungrazed grassland and ruderal vegetation. Scrub communities may also spread naturally from woodland onto suitable adjacent land, or colonize naturally if ruderal vegetation such as dense bracken is left ungrazed.

At its most diverse, dry scrub offers good bird and invertebrate habitat, and a sheltered environment for species more typical of woodland ground flora, such as bluebells and common dog's-violet, which are revealed as a striking flush of colour on brooghs and coastal slopes in spring. Dense scrub also forms a useful habitat corridor where thick woody hedgerows link otherwise isolated semi-natural habitats. Typical species include hazel, holly, immature elm, downy birch and elder in addition to the ubiquitous gorse, hawthorn and blackthorn.

Naturally regenerating areas of scrub, such as bracken areas on which young ash, mountain ash/rowan and oak are starting to grow up, represent the start of a process of semi-natural woodland regeneration that will eventually result in mature woodland communities if allowed to grow up in the long-term absence of clearance or grazing pressure. This process is a crucial element of diverse woodland ecosystems, and for this reason semi-natural scrub communities adjacent to semi-natural broadleaved woodland have an important role to play in woodland conservation.

Classification

Dry scrub on the Island may be classified broadly as outlined in **Table 4.1.c**.

Table 4.1.c: Broad categories of dry scrub on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (see Rodwell, 1992 ¹)	Rarities present?	Protected species present?
<p>Hawthorn scrub This is mostly concentrated in the base-rich south of the Island Although very commonly planted in hedges and occasional in broadleaved woodland, extensive hawthorn scrub is uncommon, possibly because it favours base-rich soils. There is much doubt as to whether hawthorn has ever been truly native to the Island (see Allen, 1984).</p>	No	<p>W21 – <i>Crataegus monogyna</i>-<i>Hedera helix</i> scrub Note: scrub categories in the NVC are considered ubiquitous throughout, hence no sample maps were given</p>	Not usually	May support Schedule 1 birds
<p>Coastal scrub Characteristic of coastal slopes, where it may take a very low, wind-sculpted form. Usually dominated by blackthorn, although stunted oaks, birch, hawthorn, rowan and other scrub species may also occur. Bracken and/or bramble underscrub is common, and may have a striking bluebell-dominated “ground flora” in Spring.</p>	Variable; may be quite diverse, especially if woodland ground flora species are present in the underscrub.	<p>W22 – <i>Prunus spinosa</i>-<i>Rubus fruticosus</i> scrub W25 – <i>Pteridium aquilinum</i>-<i>Rubus fruticosus</i> underscrub</p>	Not usually; however, coastal bracken underscrub with common dog’s violet may support Dark green fritillary butterflies	May support Schedule 1 birds or bush crickets.
<p>European gorse scrub Generally regarded as a nuisance, but may provide nitrogen-rich litter and protection from grazing, both of which may offer cover for regenerating native woodland.</p>	No	<p>W23 – <i>Ulex europaeus</i>-<i>Rubus fruticosus</i> scrub W25 – <i>as above</i></p>	Not usually	May support Schedule 1 birds;

Type and Description	Species rich?	Closest NVC categories (see Rodwell, 1992 ¹)	Rarities present?	Protected species present?
Hedgerows Ancient hedgerows are not a feature of the Manx countryside, and sod hedges are more prevalent than woody hedges; however, there are some examples of well-established species-rich woody hedgerows, especially in the North of the Island, which have a diverse ground flora and offer excellent bird and invertebrate habitat.	Variable; not usually very diverse, but exceptional examples may have good ground flora and invertebrate diversity	Usually dominated by hawthorn, blackthorn or European gorse, so related to any of the above NVC categories. May also have a ground flora with some elements of semi-natural broadleaved woodland types listed in Table 4.1.a	Not usually	May support Schedule 1 birds, bush crickets in coastal sites, or, rarely, Schedule 7 plants

Note: dwarf heath shrub communities, including Manx gorse habitats, may also approach scrub height on suitable sites; these are covered in Chapter 4.3 Lowland heathland, and Chapter 4.7 Upland Habitats.



Hazel *Corylus avellana* at Glen Maye ASSI

Dry scrub status

The 1994 Ecological Survey indicates that around 6% of the Island's semi-natural habitat consists of scrub (over 1000ha) (Sayle *et al*, 1995). However, the Survey did not distinguish between dry and wet scrub; the exact proportion of curragh is therefore unclear. What is obvious is that dense scrub is far more prevalent on the Island than semi-natural broadleaved woodland, a fact which suggests, perhaps, that one way in which scrub should be valued is as a naturally regenerating precursor to more semi-natural woodlands in the future.

The importance of scrub as a diverse habitat has not historically received much recognition from conservationists pre-occupied with trying to save dwindling semi-natural woodland. However, the value of scrub as a habitat in its own right appears to be increasingly recognised, particularly in the case of ancient and/or species-rich hedgerows, which are now a UK Biodiversity Action Plan Priority Habitat. Scrub is also referred to in the Habitat Action Plan for another Priority Habitat, Maritime cliffs and slopes (covered in Chapter 4.6 Coastal Habitats).

Minimum selection criteria for dry scrub

Areas of **dense dry scrub** which support a range of native species as listed in Appendices A and B, and which are not adversely affected by introduced species (Appendix C), will be considered for selection where:

- The area of scrub concerned is larger than 4ha and fulfils at least one of the Priority Sites Criteria listed in Part 3; or
- The area of scrub is immediately adjacent to a semi-natural broadleaved woodland or species-rich broadleaved plantation which is also under consideration as an ASSI, regardless of the size of the area of scrub concerned; or
- The scrub forms part of an upland or coastal habitat group which is under consideration as an ASSI, regardless of the size or the area of scrub concerned.
- Sites where dry scrub is thought likely to form a viable precursor to naturally regenerating native woodland will usually be given the highest priority for selection.

Species-rich hedgerows which support a range of native species as listed in Appendices A and B, and which are not adversely affected by introduced species (Appendix C), will be considered for selection where:

- The hedgerow in question fulfils at least two of the Priority Sites Criteria listed in Part 3, one of which must be Size; or
- The hedgerow forms part of a larger designation area (eg. adjacent to a semi-natural broadleaved woodland) as for dense dry scrub (above).

4.1.4 Curragh

Description and Distribution

The term "curragh" is an old Manx word for bog; however, it has come to be used to describe the willow-dominated wet scrub which forms a very characteristic element of the Manx landscape on wet, peaty soils. It is restricted to lowland areas, principally the Ballaugh Curragh (an ancient basin mire) and the Central Valley Curraghs, which include Greeba Curragh and similar, smaller areas of wet willow scrub in the central lowlands. Small areas of curragh also form elsewhere where drainage is impeded, often in association with open water and swamp habitats, but not on the same scale as the Ballaugh or Greeba curraghs.

Curragh usually occurs within a mosaic of interrelated wetland habitats such as fen mire and bog pools. It may also contain ditch systems related to ancient and modern agricultural activities. These associated habitats may support rare and/or locally-distributed plant and invertebrate communities. In the case of invertebrates, research is incomplete, but mollusc, beetle and spider records all suggest that open water habitats in association with curragh are an important element of Manx biodiversity (BDIS). Curragh is perhaps most well-known for its value as a habitat for songbirds, water birds and birds of prey such as the hen harrier, which roosts on undisturbed wet ground in the winter. High numbers of bird species can be found in areas of curragh.

Classification

Manx curragh forms a single category in its own right. It is overwhelmingly dominated by one species – Grey willow, *Salix cinerea ssp oleifolia* ("sallies"). In this respect it most resembles NVC type W1 *Salix cinerea-Galium palustre* (grey sallow/marsh bedstraw) woodland, for which there are two Isle of Man type samples mapped in the British Plant Communities volume relating to woodland and scrub (Rodwell, CUP 1991). However, it also has affinities with wet ash, alder or birch woodland such as W4 *Betula pubescens-Molinia caerulea* (downy birch/purple moor-grass) woodland, or W8 *Alnus glutinosa-Fraxinus excelsior-Lysimachia nemorum* (alder/ash/yellow loosestrife) woodland (Rodwell, 1992¹).

Curragh status

Because of its association with diverse wetland habitats and its potential for abundant bird life, curragh has a well-documented place in Manx cultural history. Areas such as the Ballaugh Curragh provide an invaluable archaeological resource and important pollen records, in addition to its high importance for nature conservation. This importance has long been recognised in the case of the Ballaugh Curragh, the central part of which is under the protection of Manx National Heritage. Other areas of curragh have no such protection, and the relatively rich soils which underlie the willow scrub make these areas popular targets for agricultural reclamation. The area of mature curragh outside MNH land is therefore considered to be under threat. For these reasons, The Manx Wildlife Sites Handbook recommends that all areas of alder carr and all good examples of willow scrub over 2ha in extent should be considered for Wildlife Site selection (Keehan, 1999).

In the wider context, wet woodland, including wet willow habitat like Manx curragh, is recognised in the UK Biodiversity Action Plan as a Priority Habitat. Two other strongly related habitats, Floodplain alder woodland and Bog woodland, are listed in Annex 1 of the EU Habitats Directive. The diverse and complex mosaic of wetland habitats contained within large areas of Manx curragh, and its importance for birds, make it a potential candidate for Ramsar site selection as well as national designation.

Minimum selection criteria for curragh

- All areas of curragh over 10ha in extent should be considered for selection;
- All areas of curragh over 2ha in extent which occur in association with at least one other wetland habitat should be considered for selection where they are not adversely affected by negative factors such as invasive introduced species (see Appendix C), tipping, heavy stock poaching or other long-term damage;
- All other areas of curragh should be considered for selection where they fulfil at least one of the Priority Sites Criteria listed in Part 3 and form part of a larger habitat mosaic such as wet woodland and fen mire;
- Other curragh areas may be selected on the grounds of individual species criteria.

See also: semi-natural broadleaved woodland criteria (Section 4.1.1) and especially the criteria for wet woodlands.

Appendices to Chapter 4.1: Diagnostic species lists

Appendix A: Tree and shrub species thought to be native to the Isle of Man

Scientific name	English name	Manx name	Notes
<i>Alnus glutinosa</i>	common alder	farney	A good tree for wet places, except where the soil is very acid.
<i>Betula pendula</i>	silver birch	beith argid; billey veih	Prefers well-drained sites.
<i>Betula pubescens</i> <i>ssp pubescens</i>	downy birch		Tolerates wetter, more acid soils than <i>B. pendula</i> . The commonest birch on the Island.
<i>Betula pubescens</i> <i>ssp tortuosa</i>	downy birch		As above, but a more shrubby, upland subspecies. Has distinctive aromatic foliage.
<i>Corylus avellana</i>	hazel	couyll	Thrives in most habitats, esp. richer soils on sheltered sites.
<i>Fraxinus excelsior</i>	ash	unjin	Thrives in most habitats, esp. richer soils.
<i>Ilex aquifolium</i>	holly	cullyn; hollin	Thrives best on fairly dry sites.
<i>Lonicera periclymenum</i>	honeysuckle	ullaagagh	Hedges and scrub; tolerates coastal conditions.
<i>Malus sylvestris</i>	crab apple		Native trees (thought to be very rare) occur in sessile oakwood; otherwise much-planted.
<i>Populus tremula</i>	aspen	chengey ny mraane; cron craee	A good tree for wet or rocky places.
<i>Prunus spinosa</i>	blackthorn; sloe	drine airn	Tolerant of coastal conditions and native to most of the Island - hence more suitable than hawthorn for most native planting.
<i>Quercus petraea</i>	sessile oak	darrag	Sessile oak occurs on wetter, more upland areas and more acidic soils than English oak, and is the more typical oak on Man; however, hybridisation between the two species makes it difficult to find "100% sessile" trees.
<i>Quercus robur</i> , <i>Quercus robur x</i> <i>petraea</i> hybrids	pedunculate or English oak; oak hybrids	darrag	English oak is more typical of drier, lowland areas than sessile oak. It has been planted extensively, and many oaks on the Island are intermediate in characteristics between the two species. These hybrids are themselves variable.
<i>Rosa canina</i>	dog rose		A group of closely-related wild roses, of which the Isle of Man tends to have types which are typical of north-west England. In general these are downier than the dog-roses found in southern England.
<i>Rosa sherardii</i>	Sherard's downy-rose		A fairly common scrub and hedge species.
<i>Rubus fruticosus</i> <i>agg.</i>	brambles	dress smeyr	Very variable, with lots of varieties indigenous to Man. Bramble usually spreads by itself. If it is to be planted, the use of stock local to the area is strongly recommended (with cultivated varieties, many of which are over-vigorous, to be avoided).

<i>Rubus idaeus</i>	raspberry	berrish yu croaw gharey	Usually occurs on the edges of woodland and heath.
<i>Salix aurita</i>	eared willow		A bushy, low-growing willow of wet, acid moorland and curragh. Forms good bird habitat.
<i>Salix caprea</i> ssp <i>caprea</i>	goat willow; pussy willow		Uncommon on the Island. Prefers drier, more calcareous soils than the common grey "sallie".
<i>Salix cinerea</i> ssp. <i>oleifolia</i>	common willow	shellagh	Common in wetland habitats throughout the Island. Some doubt as to how native this species is on the Island.
<i>Sambucus nigra</i>	elder	tramman	Tolerant of coastal conditions; strong cultural significance.
<i>Sorbus aucuparia</i>	rowan	cuirn	strong Manx cultural significance
<i>Ulmus glabra</i>	wych elm	lhiauuan	Much-planted; possibly native (hard to tell). Suitable for sites with base-rich soils.

Appendix B: Woodland ground flora species associated with semi-natural woodland

Ferns

Scientific name	English name	Manx name	Distribution/Notes
<i>Athyrium filix-femina</i>	Lady fern		Shady places; very common
<i>Dryopteris aemula</i>	Hay-scented buckler-fern		Glens, shady banks; rare and virtually confined to the east
<i>Dryopteris affinis</i>	Scaly male fern		Glens and hedge banks; frequent
<i>Dryopteris carthusiana</i>	Narrow buckler-fern		Damp woods; very rare, mostly confined to glens along the west coast but found more recently further inland
<i>Dryopteris dilatata</i>	Broad buckler-fern		Shady places; common
<i>Dryopteris filix-mas</i>	Male fern		Shady places; common
<i>Hymenophyllum wilsonii</i>	Wilson's filmy fern		Boulders on moist upland glens; very rare
<i>Oreopteris limbosperma</i>	Lemon-scented fern		Glens, stream sides; common in hills
<i>Phegopteris connectilis</i>	Beech fern		Glens, damp shady rocks in hill areas; rare.
<i>Polystichum aculeatum</i>	Soft shield-fern		Hedge banks and glens esp. in wetter areas; rare and very scattered.
<i>Polystichum setiferum</i>	Hard shield-fern		Sheltered lanes, banks, glens; occasional, plentiful around Glen Maye

Herbs

Scientific name	English name	Manx name	Distribution/Notes
<i>Ajuga reptans</i>	Bugle	Lus ny Chayrn	Glens, roadsides etc; common
<i>Allium ursinum</i>	Wild garlic	Garleyd Feie	Distribution likely to have been modified by human activities
<i>Anemone nemorosa</i>	Wood anemone	Lus ny Geayee	(glens in south introduced – Summerhill, Colby and Silverdale – Allen) Locally abundant in Northern and Eastern Glens
<i>Arctium minus</i>	Lesser burdock	Bollan-dhoa	Woodland; very rare
<i>Chrysosplenium oppositifolium</i>	Opposite-leaved golden saxifrage		Damp shady places; common
<i>Conopodium majus</i>	Pignut	Curlan	Fields, shady places; common

<i>Digitalis purpurea</i>	Foxglove	Sleggan-sleeu	Hedges, glens; common
<i>Equisetum sylvaticum</i>	Wood horsetail		Glens, swamps and boggy fields; local mainly in east central area
<i>Fragaria vesca</i>	Wild strawberry	Soo Halloin	Glens, banks and other shady places; local
<i>Geranium robertianum</i>	Herb Robert	Crouw Yiarg	Hedge banks glens; common
<i>Geum urbanum</i>	Wood avens		Glens, plantations, shady roadsides and cliff faces; local
<i>Glechoma hederacea</i>	Ground ivy	Ard Losserey	Glens, hedges. Locally abundant esp. west and north coasts
<i>Hyacinthoides non-scriptus</i>	Bluebell	Gleih-vuc	Beware of confusion with Spanish and hybrid bluebells – see Appendix C
<i>Hypericum androsaemum</i>	Tutsan		Damp places in light shade, wet rocks; frequent - sometimes locally abundant in the moister districts
<i>Lathyrus linifolius</i>	Bitter-vetch		Sea cliffs, grassy banks, glens (normally within a mile of sea). Very local - mainly SW coast and Douglas bay area
<i>Lysimachia nemorum</i>	Yellow pimpernel		Glens, stream sides, wet woodland; frequent but locally distributed.
<i>Melampyrum pratense</i>	Common cow-wheat		Wet acid woodland below 600 ft. Rare
<i>Oxalis acetosella</i>	Wood sorrel	Bee Coog	glens and hedge banks locally abundant
<i>Potentilla sterilis</i>	Barren strawberry		Glens, dry banks; common
<i>Primula vulgaris</i>	Primrose	Sumark	Hedge banks; common
<i>Ranunculus ficaria</i> ssp. <i>ficaria</i>	Lesser celandine	Lus ny Mlainyn	Shady places, banks hillsides; very common (Note: the ssp. is without bulbils – in contrast to the garden ssp. <i>R. ficaria</i> ssp. <i>bulbifera</i>)
<i>Sanicula europaea</i>	Sanicle		Shady glens. Local mainly between Laxey and Ramsey
<i>Silene dioica</i>	Red campion	Lus ny Ferryghyn	Glens, hedges; common
<i>Stellaria holostea</i>	Wood stitchwort	Lien ny Ferrishyn	Hedge banks, glens; common
<i>Teucrium scorodonia</i>	Wood sage	Lys y Toar Vrein	Hedge banks, glens; common
<i>Veronica montana</i>	Wood speedwell		Very rare – one known site adjacent to Sulby Claddagh
<i>Vicia sylvatica</i>	Wood vetch		Damp, sheltered hollows opening onto the sea; rare and very local
<i>Viola riviniana</i>	Common dog's violet		Glens, hedge banks; common

Grasses and sedges

Scientific name	English name	Manx name	Notes
<i>Carex remota</i>	Remote sedge		Damp shady places; local. Southern streams and some East coast glens
<i>Luzula pilosa</i>	Hairy woodrush		Glens, shady places, open moors; occasional. The moorland occurrences (woodland relics ?) are all in the south west (also Glen Maye on open cliffs)
<i>Melica uniflora</i>	Wood melick		Glens, shady hedge banks; local. Perhaps accidentally introduced at some sites.
<i>Festuca altissima</i>	Wood fescue		Rocky woods; very rare
<i>Carex sylvatica</i>	Wood sedge		Glens, typically by waterfalls; rare

<i>Brachypodium sylvaticum</i>	Lesser wood-brome		Glens, hedge banks etc; locally abundant
<i>Luzula sylvatica</i>	Great woodrush		Shady places etc; locally abundant. A hardy survivor of deforestation
<i>Bromopsis ramosa</i>	Hairy-brome		Shady places on basic rocks; very rare
<i>Carex laevigata</i>	Smooth-stalked sedge		Moist, shady places esp. by waterfalls Locally distributed.

Appendix C: Plants not of native origin which may indicate unfavourable condition

Scientific name	English name	Manx name	Notes
<i>Montia sibirica</i>	Pink purslane		A rapidly increasing garden escape, capable of reducing native ground flora diversity by invading existing vegetation communities, esp. on damp sites.
<i>Fallopia japonica</i>	Japanese knotweed		A highly invasive notifiable weed; spread of this plant into the wild is an offence under the Wildlife Act 1990. Especially invasive along shady stream banks.
<i>Rhododendron ponticum</i>	Rhododendron		Well-established parkland escape known to reduce scrub layer and ground flora diversity by invading woodland on more acid soils
<i>Saxifraga spathularis</i> x <i>S. umbrosa</i>	London pride		A garden escape which may indicate a modified ground flora; can invade existing native ground flora in suitable sites
<i>Circaea lutetiana</i>	Enchanter's nightshade		A native of woodlands over much of the British Isles; not known to be invasive but may indicate a modified ground flora
<i>Symphoricarpum album</i>	Snowberry		Planted as game cover and as a parkland ornamental. May invade native scrub layer and reduce ground flora diversity; widespread on the Isle of Man but usually slow to invade new sites.
<i>Crocsmia aurea</i> x <i>C. pottsii</i>	Montbretia		A popular garden plant, increasingly spreading into open, rocky woodland sites and coastal broughs. May displace native ground flora.
<i>Fuchsia magellanica</i>	Fuchsia		A characteristic Manx escaped garden plant, not usually invasive. May indicate modified conditions, often found with Montbretia and other escapes.
<i>Hyacinthoides hispanica</i>	Spanish bluebell		A large-flowered, scentless bluebell often grown in gardens; hybridises with and displaces native bluebell <i>Hyacinthoides non-scripta</i>
<i>Hyacinthoides hispanica</i> x <i>non-scripta</i>	Hybrid bluebell		A vigorous, rapidly-spreading hybrid between native and Spanish bluebell. Known to displace native ground flora; lack of scent and different flower-shape mean that this plant cannot provide the same ecological niche as native bluebells, leading to effects on invertebrate diversity and abundance.
<i>Prunus</i> spp., especially <i>P. cerasus</i> , <i>P. cerasifera</i> and <i>P. laurocerasus</i>	Non-native ornamental cherries, especially dwarf cherry, cherry plum and cherry laurel		All of these species are capable of invading native scrub and woody hedgerows, and are regularly planted as ornamental trees throughout the Island.

List references: Allen, 1984; BDIS; Clement & Foster, 1994; Garrad, 1972; Preston et al, 2001.

4.2 Lowland grasslands

Description

Grasslands are here taken to mean low- to medium-height, herbaceous vegetation communities which are strongly characterised, if not dominated, by grasses. This includes patches within larger areas of habitat mosaic as well as more obviously-defined meadows and pastures. A site does not therefore have to have a homogeneous turf in order to be classified as grassland.

The very wide range of exposure levels, topography, hydrology, geology and farming methods on the Isle of Man has led to a correspondingly wide range of grassland types, often occurring in complex mosaics with heathland, scrub, marsh and upland or coastal habitats. The degree to which a grassland may support rare plant and/or animal species, and the degree of vulnerability of the grassland, is dependent on all these location-related factors and more. This, combined with the fact that grassland is often a successional stage which naturally tends to progress swiftly towards more woody vegetation communities, makes species-rich grassland one of the most challenging habitats to conserve (see NCC 1989 Ch3 and Crofts & Jefferson 1999).

This chapter covers typical lowland grasslands; **upland pastures** are subject to special ecological and management issues, and are treated separately - see **Chapter 4.7: Upland Habitats**. Because grasses are such a ubiquitous element of Manx habitats, this chapter may also relate closely to the chapters on lowland **heathland**, bogs, **fens** and flushes, and **coastal habitats**.

Assessing lowland grasslands

All of the points on the Checklist of Priority Sites Criteria listed in **Part 3** are relevant to grasslands. Because of the wide range of grassland types and associated management issues on the Island, the description, classification, status and distribution of the four main grassland types are each treated separately here, for clarity and ease of reference. However, when assessing a grassland it should be borne in mind that this is an artificial separation. Sites which do not "fit" the broad categories are not necessarily poor examples of species-rich communities, and may be as important for wildlife as more "typical" grassland types. Many sites of nature conservation importance support a range of different grassland types, sometimes grading into each other, sometimes distinct, and quite possibly from more than one of the four main categories. Indeed, some grassland types on the Island are *only* present as small patches within larger habitat mosaics. For this reason, **the assessment process should aim to recognise good examples both of large areas of a single grassland type, and patches of grassland within larger mosaics of different habitats.**

The majority of the grasslands which have a higher-than-average nature conservation value share factors which favour less vigorous and competitive plants - for example, low nutrient status, high exposure levels, very wet or very dry conditions, extreme pH levels, or heavy exposure to salt spray. Situation and management history (e.g. continuity of traditional farming methods) are therefore of special concern. The position of the grassland within the surrounding ecological unit can also make a big difference to its importance for wildlife, especially where invertebrate species are concerned. Other factors, such as size, level of disturbance, and adjacent land use should all be taken into account when assessing the potential merits of designating a grassland site.

In the case of very rare grassland communities, **all** good examples will qualify for selection as ASSIs provided that they are of sufficient size and fulfil the minimum criteria as listed in the relevant section of this Chapter. In most cases, however, candidate sites will be selected from the best good examples of each grassland type and variant. Grasslands will be considered to be "good examples" where they not only fulfil all the minimum criteria described in this chapter, but also one or more of the Priority Site Criteria listed in Part 3.

Lowland grassland types

For the purpose of these Criteria, grasslands on the Isle of Man have been grouped as follows:

Acid grassland: Dry grassland and pastures on base- and nutrient-poor soils, usually grazed and/or subject to exposed conditions. Also known as **calcifugous** grassland. Often in association with heath and/or bog. The commonest grassland type on the Island.

Note: Upland pastures are treated separately in **Chapter 4.7: Upland habitats**.

Neutral grassland: Dry or wet species-rich grassland, wet rush pastures and a range of semi-improved and improved grasslands, on neutral soils. Also known as **mesotrophic** grasslands. These may be grazed, cut for hay, or unmanaged. These sites generally have good potential for agricultural improvement, hence neutral, species-rich grasslands are now a rarity on the Island.

Calcareous grassland: Dry grassland, often very species-rich, on base-rich soils, usually grazed (at least by rabbits if not by livestock) and/or subject to exposed conditions. Also known as **calcicolous** grassland. Calcareous soils are limited to the limestone areas in the south east of the Island, and provide some of the better agricultural land on the Island. Species-rich calcicolous grasslands are therefore both rare and local, and usually restricted to exposed, coastal or inaccessible sites. For this reason Manx calcicolous grasslands often have affinities with coastal grassland and sea cliff and slope habitats.

Coastal grassland: Cliff-top and slope communities, often extending some way inland ("sub-maritime"). Sometimes grazed, always subject to heavy exposure to wind and salt deposition. *Note:* in the *UK Guidelines for selection of biological SSSIs*, sub-maritime grassland is covered under Coastal Habitats rather than Lowland Grassland. The island nature of Manx ecology means that sub-maritime grassland tends to occur further inland and more in conjunction with other lowland grassland types - and hence with similar management issues - than in the UK. It has therefore been treated here alongside other lowland grassland types. More strictly maritime habitats with grassland affinities include cliff crevice, saltmarsh and dune vegetation, but as these have a different ecology to most grasslands they are treated separately under **Chapter 4.7: Coastal habitats**.

These Manx grassland types, and the National Vegetation Classification (NVC) categories they most closely resemble, are listed in detail on **Tables 4.2a-4.2d**, below. It should be noted when assessing a site that some characteristic UK grassland species are naturally absent from the Isle of Man; to help with assessment, the full range of vascular plants which typically occur in species-rich lowland grassland communities on the Island are listed in **Table 4.2e**.

"Similar communities"

Very few grassland samples were taken from the Isle of Man to contribute to the original research that formed the basis of the National Vegetation Classification (Rodwell, 1992). Perhaps because of this, Manx vegetation communities do not always fit well within NVC categories. This does not mean that the NVC is worthless here, but it has led to a situation where the NVC alone is not usually sufficient for assessing the nature conservation importance of a Manx site. See **Section 1.3.3**, above, regarding the use of the NVC in the Manx context.

In order to take into account the shortcomings of the NVC with respect to classifying Manx vegetation, most of the detailed selection criteria for grasslands refer to an NVC type and "similar communities". Similar communities are here defined as vegetation communities which have the majority of species in common with the NVC type, but which may lack some of the determining species from the relevant Floristic Table in the NVC for any of the following reasons:

- due to local conditions, some species are typically missing from this habitat on the Island;
- some NVC determinant species don't occur on the Island at all due to the general paucity of flora compared with the UK;
- due to local conditions, one or more determinant species of the NVC type in the UK are typically "replaced" by different species on the Island (e.g. because they fit a similar niche in the community and are more tolerant of maritime conditions; or e.g. because they represent a "western British" variant).

As a general guide, any community which supports a good range of the species listed in **Table 4.2e** will be considered for selection as a species-rich grassland. If sites selected in this way support vegetation communities that aren't a good "fit" for the NVC categories listed in the main selection criteria, the reasons for the choice will be made clear in the ASSI citation details.

4.2.1 Acid grasslands

Description and Distribution

Acid grassland, or calcifugous, communities consist of species which are capable of tolerating base-poor, acidic conditions. Such vegetation is common on the Isle of Man because the predominant bedrock is Manx slate, which naturally tends to form fairly acidic, podzolic soils when weathered, whilst the cool, wet climate keeps the soil pH low, with an average of pH 5-6. Additional organic deposits which are also acid in nature often form over these soils, mostly derived from heather and gorse litter, but also from other accumulated vegetation. In some areas this humus layer is thin, but it may reach 30cm or more in depth, and is usually present to a greater or lesser degree in all upland situations (Fullen, Harris & Kear 1999). Where the topography is such that dead vegetation becomes too waterlogged to decompose away, true peat layers can build up which may reach over a metre in depth (Robinson & McCarroll 1990 Ch5). These unusual areas commonly form a basin, with curragh, bog and swamp communities on the wetter peat at the centre, and grasslands around the edges. Due to this range and prevalence of acidic habitats, the majority of grassland on the Island is more or less acidic in character (see Sayle *et al*, 1995). Such grassland is naturally less species-rich than grassland on base-rich soils because of the more challenging conditions.

Heavy leaching, particularly in the uplands where rainfall levels are very high, can result in peaty "mor humus" topsoil with very low levels of nutrients available to plants, and a pH as low as 3.2, i.e. very acidic (Fullen, Harris & Kear 1999). Upland pasture on this kind of soil is frequently interspersed with bog and/or heath; for this reason, the assessment of very acid, upland grasslands cannot usually be divorced from consideration of associated upland habitats, with which grassland will often form one part of a complex mosaic. **Acid upland pastures are therefore treated separately - see Chapter 4.7: Upland habitats.**

The two commonest types of lowland acid grassland on the Island are:

- mainly dry, extensive sheep pasture on the margins of the uplands; and
- mainly wetter pastures, often rather scattered and isolated in the modern farming landscape, and often associated with heath and bog.

Both the main types of lowland acid grassland are the result of habitat modification by human activities, and both tend to "revert" to other vegetation types if left unfarmed. In less exposed areas, dry acid grassland is frequently subject to agricultural improvement such as drainage, liming and re-seeding. Wetter and more acid areas are not usually considered worth a lot of agricultural effort, but may still receive drainage and control of unwanted species. In both cases the dominance of grasses is maintained by grazing, as almost all the calcifugous grassland on the Island tends towards heathland if left unmanaged, and is very prone to encroachment from European gorse and bracken. These issues mean that farming activities are a very important factor in all acid grassland conservation. It should be noted, however, that whilst calcifugous grasslands will disappear if taken over completely by heathland, bracken or gorse, the presence of any of these is not necessarily a sign that the conservation status of the grassland is poor. For example, stands of bracken may support a "ground flora" of species which flower in spring and complement the grassland habitat early in the year.

Classification of lowland acid grasslands

Lowland acid grassland on the Island may be classified broadly as outlined in **Table 4.2.a**.

Lowland acid grassland status

Acid grassland is common throughout most of the Isle of Man, making up around 7% of the Island's land use total. This equates to over a quarter of all the semi-natural habitat present, and nearly half of all the grassland (Sayle *et al*, 1995). The mean average size of mapped, unimproved acid grassland area is 12ha (BDIS). However, this figure includes upland grassland (which is covered in Chapter 4.7).

Unimproved lowland acid grasslands were not mapped separately, but correspond more closely with the next category, semi-improved acid grassland, which has a mean average size of 1.8ha and is much more scattered in distribution (BDIS). Good examples of lowland acid grassland generally occur singly or in small groups of fields which have been spared agricultural intensification for reasons of inaccessibility, high water-table, exposed conditions or other obstacles to farming. The situation with the uplands is rather different, as several specific areas of key ecological importance have long been recognised, all of which support a complex mosaic of interrelated habitats. Uplands are therefore dealt with separately in **Chapter 4.7**.

The Manx Wildlife Sites Handbook states that all lowland acidic grassland of at least 2 hectares, or 1 hectare if in association with other semi-natural habitats, are eligible for Wildlife Site selection if the site supports a good range of characteristic vascular plant species (Keehan 1999). The importance of lowland acid grassland is recognised in the UK by the inclusion of Lowland dry acid grassland as a Biodiversity Action Plan Priority Habitat – as with other dry lowland grassland types. The importance of this habitat for birds is also noted (UK Biodiversity Group 1998 Vol II – see UKBAP.org.uk).

Minimum selection criteria for lowland acid grassland

U2 – Wavy hair-grass *Deschampsia flexuosa* grassland

Examples of U2 grassland or similar communities will qualify for selection if they are over 10ha in total extent and fit at least one of the Priority Sites Criteria as listed in **Part 3**. Further priority will be given to examples which form part of a habitat mosaic.

U4 - Sheep's fescue/common bent grass/ heath bedstraw *Festuca ovina*- *Agrostis capillaris*-*Galium saxatile* grassland - outside of upland areas

Examples of U4 grassland or similar communities will qualify for selection if they are over 10ha in total extent and fit at least one of the Priority Sites Criteria as listed in **Part 3**. Further priority will be given to examples which form part of a habitat mosaic.



Sheep's-bit scabious *Jasione montana*

Table 4.2.a: Broad categories of acid grassland on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (see Rodwell, 1992 ³)	Rarities present?	Protected species present?
<p>Lowland acid grassland</p> <p>The commonest lowland grassland type on the Island, often occurring as a heath/grass mosaic or in conjunction with bog and upland pasture.</p> <p>May occur in flat lowland farmland or lower and middle slopes of hills. Usually grazed; sometimes cut for hay.</p>	<p>Generally fairly species-poor due to the poor nature of the substrate, but may have quite a good mix of herb species, particularly where it forms part of a grass/ heath mosaic.</p>	<p>U2 - <i>Deschampsia flexuosa</i> grassland Frequent but not particularly extensive on the Island; patches may occur throughout "heathy" U4 grassland and in between patches of heather on upland margins. Often quite diverse, but vulnerable to burning, liming, fertilizing, scrub invasion and overgrazing, all of which can adversely affect the species diversity. No Island sample in NVC maps.</p> <p>U4 - <i>Festuca ovina</i>- <i>Agrostis capillaris</i>-<i>Galium saxatile</i> grassland One of the commonest grassland types, often covering extensive areas especially on upland fringes. See also Chapter 4.7. One Island NVC sample.</p>	<p>A possibility of invertebrate rarities. May form part of a feeding territory for some upland waders and birds of prey, especially in conjunction with heath mosaic.</p>	<p>Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing.</p> <p>Species-rich acid grassland may also support orchids, notably heath-spotted orchid, <i>Dactylorhiza maculata</i> ssp <i>ericetorum</i>, and northern marsh orchid <i>Dactylorhiza purpurea</i>.</p>
Upland pasture	see Chapter 4.7 - Upland Habitats			

4.2.2 Neutral grasslands

Description and Distribution

Neutral, or mesotrophic, grassland communities do not show a distinctive dominance by either acid- or lime-loving species. They are generally associated with enclosed agricultural land on neutral or slightly acid soils. Throughout the Isle of Man the majority of neutral grasslands are found on the more productive agricultural land in lowland areas, away from the more acid soil conditions in the hills.

The main types of neutral grassland on the Island are:

- Improved rotation and permanent pasture dominated by perennial ryegrass and white clover;
- Semi- or un-improved species-rich permanent pasture and hay-meadows;
- Rough grassland with little or no grazing, dominated by false oat-grass.

Semi-improved and unimproved pastures, which result from extensive farming practices, are becoming increasingly rare (BDIS). Most of the potential sites for species-rich neutral grassland have long since been improved to support arable, rotation grassland or silage crops, replacing diverse vegetation communities with more productive ryegrass leys, which are now overwhelmingly prevalent. Such improved pastures are not generally considered to have any value for wild plant conservation, but may be important in the conservation of farmland birds, particularly choughs (see www.rspb.org.uk/birds/guide 2005).

As with other grassland types, neutral grassland is closely associated with scrub which, if allowed to develop, will overtake the low-growing grasses and herbs. This can reduce the species richness of grassland quite rapidly, especially where a stand of one particular shrub comes to dominate the whole site. However, the presence of occasional scrub such as hawthorn, willows, blackthorn or gorse does not automatically indicate that grasslands have a low conservation status. Scattered scrub can add considerably to the conservation value of grasslands, providing a diversity of structure that offers a richer habitat for birds and invertebrates, especially where there are young shrubs and a good variety of species. Butterflies in particular are known to benefit from a mix of scrub patches in otherwise open grassland (Asher et al, 2001 Ch2). This needs to be borne in mind when assessing the quality of a potential ASSI.

Although neutral grassland is primarily dependent on farming activities, mesotrophic species also occur on alternative habitats such as road verges and "brownfield" sites, where farming activities are either limited or non-existent. Road verges in particular form an important habitat refuge on the Island, and may support a rich diversity of herb species more commonly associated with traditional hay-meadows (Scott 1997 Ch4). Hence the conservation of neutral grassland species is not entirely dependent on farming methods; however, whether on farmland or not, the biological diversity of any neutral grassland will ultimately depend on carefully-timed management practices and the maintenance of low nutrient levels.

Classification of neutral grasslands

Neutral grassland on the Island may be classified broadly as outlined in **Table 4.2.b**.

Neutral grassland status

Unimproved neutral grassland is one of the rarest semi-natural habitats on the Island, occupying just 0.13% of the total grassland habitat area. Indeed, only 11ha of unimproved neutral grassland was identified during the Phase 1 Habitat Survey. Even semi-improved neutral grassland is rare compared to calcifugous (acid grassland) communities, amounting to only 634ha or 7.37% of the total grassland area (Sayle et al, 1995).

Although semi-improved sites have been modified to a certain extent by agricultural improvements, many are quite diverse and constitute some of the more species-rich grasslands on the Island (BDIS). The Criteria have therefore been designed to reflect this.

In the wider context of habitats throughout the British Isles, neutral grasslands have undergone a severe decline, particularly traditional hay-meadows. In the UK *Guidelines for Selection of Biological SSSIs* (NCC 1989) it was stated that less than 3% of such meadows remained; this figure has been steadily decreasing since then, and the majority of UK examples of the habitat are now confined to SSSIs and other protected areas. Consequently this habitat features in the UK Biodiversity Action Plan as a Priority Habitat – Lowland meadows. Some of the best and largest examples of the type are protected by international as well as national designations.

Minimum selection criteria for neutral grassland

MG1 False oat-grass *Arrhenatherum elatius*-dominated rough grassland

Examples of the most species-rich variants of MG1 grassland or similar communities will qualify for selection if they are over 1ha in extent and fit at least one of the priority sites criteria as listed in Part 3. Further priority will be given to examples which form part of a habitat mosaic and/or which are known to have important invertebrate populations.

MG5 Crested dog's-tail and knapweed *Cynosurus cristatus* - *Centaurea nigra* grassland

- **All** examples of MG5 grassland and similar communities greater than 1ha will qualify for selection, with greatest priority given to sites in regular hay-meadow management;
- Examples of MG5 grassland and similar communities on roadside verges or other permanent linear sites (e.g. disused railway lines) will qualify for selection if the total extent of the grassland is greater than 0.1ha and the site fits at least two of the Priority Sites Criteria as listed in **Part 3**. Priority will be given to Conservation and Sensitive Verges as identified in the DoT/MNCT Road Verge Survey (Scott, 1997).

Note: 0.1ha is equivalent to a roadside verge 4m wide and 250m long; or a green lane 10m wide and 100m long; or 250m of 2m verge along both sides of a road.

MG6 Perennial ryegrass and crested dog's-tail *Lolium perenne*-*Cynosurus cristatus* grassland

Examples of the most species-rich MG6 grasslands, i.e. those which share some determinant MG5 species, or similar communities, will qualify for selection if they are over 1ha in extent and fit at least one of the priority sites criteria as listed in Part 3. Further priority will be given to examples which are known to have important invertebrate populations and/or significant populations of wild orchids.

MG7 Perennial ryegrass *Lolium perenne* leys

Grasslands dominated by perennial ryegrass and cultivated clover obviously do not warrant protection on the grounds of semi-natural habitat importance, but may occasionally provide crucial feeding areas for significant populations of choughs. Sites which are exceptional in this respect may therefore qualify for inclusion within an ASSI under bird protection criteria - see **Chapter 4.9**.

MG 9, 10, 11, 13 – poor semi-improved wet and inundation grasslands

As with MG7 leys, these species-poor grassland types will not be considered for selection on the grounds of semi-natural habitat importance, but they may form part of important bird territories which are covered under bird protection criteria – see **Chapter 4.9**.

Table 4.2.b: Broad categories of neutral grassland on the Isle of Man

Type and description	Species-rich?	Closest NVC category (see Rodwell, 1992 ³)	Rarities present?	Protected species present?
<i>Arrhenatherum elatius</i> -dominated rough grassland, often unmanaged	not usually botanically rich; one sub-type MG1e may be fairly diverse in plants; all types may have good invertebrate diversity but only if the sward structure is suitable.	MG1 <i>Arrhenatherum elatius</i> grassland One Island NVC sample	not usually	Rough MG1 grassland provides feeding territory for owls; damper sites may rarely support common orchid species; sites reverting to scrub may form valuable songbird or bush-cricket habitats; undisturbed sites may form important lizard feeding areas.
Semi-natural neutral grassland, characterised by grasses such as <i>Festuca rubra</i> , <i>Cynosurus cristatus</i> , <i>Koeleria cristata</i> , <i>Anthoxanthum odoratum</i> , and <i>Poa pratensis</i> . Has frequent meadow flower species such as <i>Centaurea nigra</i> , <i>Prunella vulgaris</i> , <i>Succisa pratensis</i> and <i>Lotus corniculatus</i> . May also support a rich assortment of sedges such as <i>Carex caryophyllea</i> , <i>C. flacca</i> , and, in damper areas, <i>C. panicea</i> . May be traditional hay meadows or pastures.	potentially very species-rich in both plants and invertebrates.	MG5 <i>Cynosurus cristatus</i> - <i>Centaurea nigra</i> grassland No Island NVC samples	Several rare plant and invertebrate species rely on this habitat.	Damper sites may support common orchid species; hay meadows may support corncrakes.
Semi-improved, often re-sown grasslands where a long-term perennial ryegrass and white clover crop is accompanied by grasses more typical of an MG5 grassland, such as <i>Cynosurus cristatus</i> , <i>Agrostis capillaris</i> and <i>Festuca rubra</i> . May support a range of herb species, such as <i>Lotus corniculatus</i> , <i>Achillea millefolium</i> and <i>Leontodon autumnalis</i> . If left unimproved, may approach MG5 in character.	not usually; however, low-input and/or organic examples with short swards may support a good range of fungi	MG6 <i>Lolium perenne</i> - <i>Cynosurus cristatus</i> grassland Three Island NVC samples	not usually; however, low-input and/or organic examples with short swards may support unusual fungi	Short MG6 grassland may form valuable chough feeding areas provided animal husbandry is appropriate.
Perennial ryegrass leys, often with white clover (usually in rotation with arable crops).	no.	MG7 <i>Lolium perenne</i> leys and related grasslands	no.	Short MG7 grassland may form valuable chough feeding areas provided animal husbandry is

				suitable.
<p>Poor semi-improved wet pasture, dominated by <i>Holcus lanatus</i> and with frequent <i>Juncus effusus</i> and/or <i>Deschampsia cespitosa</i>, usually forming rough clumps and tussocks. A few herb species such as <i>Ranunculus repens</i> and <i>Cardamine pratensis</i> may be present, usually as a small overall proportion. A tough and persistent grassland resulting from grazing on poorly-drained ground. <i>Deschampsia cespitosa</i> may occur with or instead of <i>Juncus effusus</i> in less consistently wet soils; <i>J. effusus</i> is generally the more common constituent. Frequently a target for drainage and/or reseeded to improve productivity. May be found in association with more species-rich mire and swamp areas.</p>	no.	<p>MG9 <i>Holcus lanatus</i>- <i>Deschampsia cespitosa</i> grassland No Island NVC samples MG10 <i>Holcus lanatus</i>- <i>Juncus effusus</i> rush pasture (more frequent) No Island NVC samples</p>	<p>May occasionally attract winter waders such as redshank, especially if near the coast.</p>	<p>May support breeding and feeding lapwing, especially if kept short. May support curlew and snipe if allowed to become tussocky. Tussocky and varied wet pasture may form a good habitat for a wide range of invertebrates; more research is needed regarding its importance for invertebrates on the Island.</p>
<p>Poor semi-improved periodically flooded pasture, characteristic of low-lying, free-draining floodplain and coastal areas. Dominated by a mat of <i>Festuca rubra</i>, <i>Agrostis stolonifera</i> and <i>Potentilla anserina</i>. Waterlogged, freshwater patches may also be dominated by <i>Alopecurus geniculatus</i>. Often rather short and with a flattened appearance, and subject to periods of inundation interspersed with intense drying. May grade into strandlines and saltmarsh, or into more permanently wet or dry grassland.</p>	no.	<p>MG11 <i>Festuca rubra</i>- <i>Agrostis stolonifera</i>- <i>Potentilla anserina</i> grassland frequent, especially near the coast; No Island NVC samples MG13 <i>Agrostis</i> <i>stolonifera</i>-<i>Alopecurus</i> <i>geniculatus</i> grassland frequent, especially in occasionally-flooded patches in inland pastures. No Island NVC samples.</p>	not usually.	<p>May support lapwing, especially if kept short. May offer valuable feeding areas for winter waders when flooded, especially near the coast.</p>

4.2.3 Calcareous grasslands

Description and Distribution

The term "calcareous grassland" covers a range of plant communities in which low-growing, calcicole (lime-loving) plants are prominent. It is also referred to as **calcicolous** grassland. Grassland calcicoles are typically short herbs and fine-leaved grasses that only thrive where lime or other basic (alkaline) soil chemicals are present, and competition from coarser and more vigorous species is low. The habitat is commonest on shallow, well-drained soils which are base-rich (principally in calcium carbonate) due to the weathering of chalk and other types of limestone or base-rich rock. For this reason the habitat is commonly referred to as "chalk grassland" or "limestone grassland" depending on the underlying rock.

The "Manx Group" geology that underlies the majority of the Isle of Man does not provide a base-rich substrate, and lime-loving plant communities are typically restricted to rock and soils on the small areas that fall outside this zone. Typical calcareous grassland therefore occurs only in the south of the Island, centred on Castletown and Billown, in places where the underlying Carboniferous limestone is at or close to the surface. Chalk does not occur on the Isle of Man; all the examples of typical calcareous grassland occur on limestone areas. In general, these sites have a soil pH of 6.5 to 7.6 (Harris, Fullen & Hallett, 2001).

Limestone grassland is a famously attractive habitat, rich in colourful flowers and supporting a wide range of butterflies and other invertebrates. Unfortunately it is known to be declining throughout the British Isles, with a high proportion of all calcareous grassland now lost - either through farm improvement on the best and flattest land, or due to lack of grazing on the poorest and least accessible sites (UK Biodiversity Group, 1998; also see

<http://www.jncc.gov.uk/ProtectedSites/SACselection/habitat.asp?FeatureIntCode=H6210>, 2005). The Isle of Man is no exception to this, hence most examples of calcareous grassland are restricted to well-drained slopes, quarries or rocky outcrops, as is the case in the UK.

The main types of calcareous grassland on the Island are:

- Dry grassland on rock outcrops and old limestone quarries;
- Coastal turf on limestone cliffs and rock outcrops;
- Calcareous dune vegetation

Localised variations in rock type and soil depth can also produce a mosaic of grassland types, with scattered small calcicole-dominated patches within predominantly neutral grassland. This mirrors the situation in the UK, where calcicolous species are commonly confined to very small and isolated patches, often surrounded by more nutrient-rich, improved grassland: "*Zonations through various grades of improved sward have become a common feature of almost every tract of calcicolous grassland in the southern lowlands of Britain*" (Rodwell, 1992³ p114). Manx calcareous grasslands, like those in the UK and Eire, may therefore have a mixture of strict calcicoles and more neutral grassland species.

Calcicoles are not confined to typical calcareous grasslands. Plants characteristic of limestone grassland may additionally be found on a range of shallow, well-drained soils with neutral grassland and coastal turf, wherever low nutrient levels and/or exposure have led to a severe reduction in competition from more vigorous plants. The Manx coast has narrow but extensive sections of very exposed, red fescue-dominated coastal turf, which may also contain calcicoles. Calcicolous vegetation also occurs on the most base-rich of the dune grasslands at the Ayres, where the sand is shell-derived and rich in calcium. Both these communities are somewhat different in ecological requirements to typical grasslands, and are consequently covered in other parts of the Criteria - see **4.2.4**, below, and Chapter **4.6: Coastal Habitats**.

Scrub is frequently associated with calcicolous grassland, and can contribute to local biodiversity by providing shelter, food and scrub/grassland margin conditions suitable for invertebrates such as ground beetles. Scrub also provides breeding habitat for birds such as linnet (*Carduelis cannabina*) and nectar for a range of flying insects, hence a small amount of scrub is usually beneficial to wildlife (see Cullen &

Jennings, 1986; www.rspb.org.uk, 2005; Asher et al, 2001). However, many scrub species will encroach onto grassland, including blackthorn *Prunus spinosa*, hawthorn *Crataegus monogyna*, and old man's beard *Clematis vitalba*, and the scrub cover can quickly overtake short calcareous turf to the exclusion of low-growing calcicoles. Scrub control is therefore a management concern on the majority of calcareous grassland sites.

Classification

Calcareous grassland on the Island may be classified broadly as outlined in **Table 4.2.c:**

Table 4.2.c: Broad categories of calcareous grassland on the Isle of Man

Type and Description	Species rich?	Closest NVC category (see Rodwell, 1992 ³)	Rarities present?	Protected species present?
Limestone grassland Restricted to the south of the Island, and often closely associated with coastal habitats. Other examples are restricted to road verges and quarries.	Yes. Species richness is a distinctive characteristic.	CG2 <i>Festuca ovina-Avenula pratensis</i> grassland Note: several spp. typical of this category in the UK do not occur on the Isle of Man. One NVC sample for CG2d) (<i>Dicranum scoparium</i> sub-community)	Plant and invertebrate rarities are both likely.	Several orchid species including Bee orchid <i>Ophrys apifera</i> , and Pyramidal orchid <i>Anacamptis pyramidalis</i> . Choughs in coastal areas, but only on closely-grazed sites.
Coastal grassland on limestone cliffs	Restricted to the Castletown area. This habitat is treated separately in Section 4.2.4 - Coastal grassland			
Calcareous dune grassland	Restricted to shell-sand derived dunes on the Ayres. This habitat is treated separately in Chapter 4.6: Coastal Habitats			

Calcareous grassland status

Calcareous grassland is the rarest grassland type on the Island, and is restricted to such small areas that it was not assessed as part of the DAFF Phase 1 Habitat Survey report (Sayle *et al*, 1995). However, there are some notable examples of calcicole-rich grassland, usually in conjunction with other grassland types. At a non-statutory level, the Manx Wildlife Sites Handbook contains criteria for Wildlife Sites selection in which, due to the rarity of the habitat, **all** examples of lowland calcareous grassland will be eligible; this mirrors many Wildlife Sites criteria publications for counties in the UK.

Elsewhere in the British Isles, the potential for calcareous grasslands to support a very rich plant and invertebrate diversity has resulted in widespread protection of many of the best examples. At a local level, calcareous grasslands feature in the UK network of Local Nature Reserves, reflecting the importance of this kind of habitat within areas which are enjoyed by the public. Nationally, the sharp decline in the extent of lowland calcareous grasslands has led to them being made a UK Biodiversity Action Plan Priority Habitat, whilst the majority of the best sites are covered by statutory protection (UK Biodiversity Group, 1998). Calcareous grasslands therefore feature strongly in the UK SSSI system, with an estimated 60-70% of sites in both England and Wales falling within SSSIs. The value of the habitat is also recognised in the National Nature Reserve series, with a total of 30 NNRs in England and Wales in which calcareous grassland makes up a major habitat constituent (www.jncc.gov.uk , 2005).

In the wider context of European law, lowland calcareous grassland is included within the *Festuco-Brometalia* grassland identified in Annex 1 of the EC Habitats Directive as of "Community interest". The habitat is a further priority if important orchid populations are present. The international importance of lowland calcareous grasslands is recognised in the UK , where the best calcareous grassland examples also form part of the Natura 2000 network (www.jncc.gov.uk, 2005).

Minimum selection criteria for calcareous grassland

CG2 Sheep's fescue and meadow oat-grass *Festuca ovina* - *Avenula pratensis* grassland

- **All** examples of CG2 grassland and similar communities greater than 0.5ha will qualify for selection;
- Sites which have a mosaic of grassland types in which CG2 grassland or similar communities feature as a proportion of the overall vegetation will qualify for selection if their total extent is greater than 1ha and they fit at least one of the Priority Sites Criteria as listed in Part 3;
- Examples of CG2 grassland and similar communities on roadside verges or other permanent linear sites (e.g. disused railway lines) will qualify for selection if the total extent of the grassland is greater than 0.1ha and the site fits at least one of the Priority Sites Criteria as listed in Part 3. Priority will be given to Conservation and Sensitive Verges as identified in the DoT/MNCT Road Verge Survey (Scott, 1997).

Note: 0.1ha is equivalent to a roadside verge 4m wide and 250m long; or a green lane 10m wide and 100m long; or 250m of 2m verge along both sides of a road.

4.2.4 Coastal grassland

Description and Distribution

Coastal grassland differs from inland grassland communities because of the influence of salt spray and strong maritime winds. The harsh conditions limit the range of species which would normally be able to colonise similar rock and soils elsewhere, resulting in a distinctive flora that grades into more typically inland vegetation communities as soon as the terrain becomes less exposed. A cliff and cliff-top may therefore support a range of grassland types, with more maritime communities on the exposed parts and typical inland grasslands in more sheltered gullies and ledges, or on parts of the cliffs which face away from the prevailing winds (Ratcliffe, 1997 Ch4). Both inland and maritime type communities can occur very close to the strandline if conditions are suitable; or they may occur at a field's distance, or more, from the sea.

Around one tenth of all the strictly coastal habitat around the Isle of Man is grassland, almost always dominated by red fescue *Festuca rubra* (BDIS). This distinctive fine-leaved grass forms a cushion-like mat where ungrazed, or a short, fine turf where grazed by sheep and/or rabbits. Ungrazed examples of coastal grassland are often rather species-poor, but shorter turf may support a colourful range of wild flowers, especially where it grades into other species-rich grassland types on calcareous rocks and soils. These include typical seaside species such as thrift *Armeria maritima*, sea plantain *Plantago maritima*, kidney vetch *Anthyllis vulneraria*, and sea campion *Silene uniflora*. Other species which thrive in the absence of competition from larger, coarser plants (which cannot stand the exposed conditions) may also occur. These include calcicoles such as purple milk-vetch *Astragalus danicus*, and dry grassland species such as mouse-ear hawkweed *Pilosella officinarum*, English stonecrop *Sedum anglicum*, and wild thyme *Thymus polytrichus*.

The vertical nature of much of the habitat means that, when mapped, coastal grassland appears to cover a very small area in a thin strip around the coast of the Island; however, in practice it can represent a much larger available habitat due to the surface area of the sloping land not being fully represented on the map.

The diversity of the grassland is dependent to a certain extent on the nature of the coastline concerned. Soft, crumbling cliffs do not offer a stable habitat for perennial turf, and consequently favour more open vegetation communities. Cliffs of Manx slate are generally surrounded by acid-loving communities such as coastal heath, bracken stands (often with a woodland-like ground flora) and acid grassland, all of which may influence the species composition of the coastal turf. Likewise, the most species-rich coastal

grassland on the Island occurs in conjunction with limestone rocks in the area around Castletown, where it grades into more typical calcareous grassland. The areas which catch the highest rainfall tend to have a lower salinity, so stretches of coastline, whether of slate or limestone, can vary in the proportion of characteristic salt-tolerant plants present, depending on the level of rainfall they receive. These varied factors all contribute to the great diversity of Manx coastal vegetation types, even within relatively short stretches.

Coastal grassland may occur on sites where there is no grazing pressure at all, maintained only by the extremely exposed conditions within range of salt spray and maritime gales. In general, such communities are not very rich in plant species, although they may provide a sheltered mat of vegetation in which invertebrates can thrive. Areas which are closely grazed by sheep or rabbits or both usually support a much wider range of plants, and offer a warm, open, flower-rich turf which favours sun-loving invertebrates. Small amounts of scrub - particularly in association with coastal heath - can enhance the value of coastal grassland for wildlife. However, on the most diverse sites invasion by taller species can become a management concern, as gorse, bracken and bramble can rapidly take over the more delicate species if grazing pressure is removed, resulting in the loss of rare maritime herbs that cannot withstand competition.

When assessing the species richness of Manx coastal grassland it should always be borne in mind that some characteristic UK coastal and lowland grassland species are naturally absent from the Island. In general, a species-rich example of a Manx grassland might be expected to have slightly fewer species than its equivalent in the UK, even where other factors appear to be equal. Coupled with the close proximity of small patches of different vegetation types on the varied coastal terrain, this means that Manx coastal grassland tends not to express the typical makeup of NVC grassland categories. As a guide to assessing the species richness of a coastal grassland, the same approach is taken here as with Woodlands in Chapter 4.1. As an aid to site assessment Table 4.6.e) lists plant species typical of semi-natural grassland around the Island, with a guide to those species particularly associated with calcareous coastline (the rarest grassland category on the Island).

Classification

Coastal grassland on the Island may be classified broadly as outlined in **Table 4.2.d**.



Coastal grassland at Dalby

Table 4.2.d: Broad categories of coastal grassland on the Isle of Man

Type and Description	Species rich?	Closest NVC category (see Rodwell, 1992 ³⁺⁵)	Rarities present?	Protected species present?
<p>Maritime cliff and cliff-top grassland, generally dominated by Red fescue <i>Festuca rubra</i> agg.</p> <p>Note that cliff crevice communities are covered in Chapter 4.6: Coastal habitats.</p>	<p>Very species-rich when in conjunction with calcareous grassland habitats; otherwise variable. See Table 4.2.e for a list of species typical of species-rich coastal grassland</p>	<p>MC8 Red fescue and thrift <i>Festuca rubra</i>-<i>Armeria maritima</i> maritime grassland One Island NVC sample.</p> <p>MC9 Red fescue and Yorkshire fog grass <i>Festuca rubra</i>-<i>Holcus lanatus</i> maritime grassland No Island NVC sample.</p> <p>MC10 Red fescue and plantain <i>Festuca rubra</i>-<i>Plantago</i> spp. maritime grassland No Island NVC sample.</p>	<p>Plant and invertebrate rarities may occur.</p>	<p>Orchids may be present. Short coastal turf may also form a feeding territory for protected bird species, notably choughs, and forms the habitat of the lesser mottled grasshopper <i>Stenobothrus stigmaticus</i>.</p>
<p>Dune and saltmarsh communities</p>	<p>In some cases dunes and saltmarsh may be dominated by grasses, but they exhibit a different ecology to most grasslands, with different management issues. They have therefore been treated separately – See Chapter 4.6: Coastal habitats.</p>			

Coastal grassland status

Although the Isle of Man naturally has a high ratio of coastal to inland habitats, the overall proportion of unimproved coastal grassland is low: just 0.1% of the total Island area and 0.34% of semi-natural habitats. Mapped coastal grassland areas range in extent from 0.12 – 4.2ha, with a mean size of 0.8ha (Sayle et al, 1995). Even amongst coastal habitats, unimproved coastal grassland amounts to only 0.67% (Lamb/MWT 1999). Unimproved neutral grassland was mapped at 2.17% of all coastal habitats, i.e. also scarce compared with improved or poor semi-improved grassland. One reason for the relatively small area of semi-natural coastal grassland is the presence of improved farmland right up to the edge of cliff-tops in some areas. Another reason is that suitable sites for the development of a coastal turf are scarce, with many semi-natural areas supporting more ephemeral vegetation, or more overgrown habitats, or coastal heath. It should also be noted that the areas of habitat in question have been derived from orthographic mapping, which will not have measured the full surface area of the land involved due to the steepness of the terrain around much of the coast.

The importance of maritime cliff and slope grassland is recognised in the UK Biodiversity Action Plan, which listed Maritime Cliff and Slopes as a Priority Habitat. The Action Plan also notes that another Priority Habitat, Lowland Calcareous Grassland, may occur in coastal locations. In the European context, coastal grassland is recognised as a component of *Vegetated sea cliffs of the Atlantic and Baltic coasts*, which is listed in Annex 1 of the EC Habitats Directive.

Minimum selection criteria for coastal grassland

- Coastal sites which have a mosaic of grassland types in which calcareous grassland communities of type CG2 or similar (see chapter 4.2.3, above) feature as a proportion of the overall vegetation will all qualify for selection where they fit at least one of the Priority Sites Criteria as listed in Part 3;
- **All** examples of species-rich coastal grassland (i.e. grassland which support more than half of the typical Manx coastal species listed in Table 4.2e below) greater than 0.5ha will qualify for selection;
- Smaller, fragmented and/or linear examples of unimproved or species-rich semi-improved coastal grassland will qualify for selection if the total extent of the grassland is greater than 1ha and the site fits at least one of the Priority Sites Criteria as listed in Part 3;
- Priority will be given to sites which have a naturally-occurring range of coastal habitat features, e.g. ephemeral soft cliff communities, hard cliffs, coastal heath and adjacent shingle banks.



In some parts of the British Isles, Harebells *Campanula rotundifolia* have distinctive large “bells”. This is due to a minor genetic variation. Manx harebells often have this characteristic, especially near the coast. Protected habitats help to maintain this kind of locally distinctive diversity.

Table 4.2e: Vascular plants characteristic of semi-natural grassland on the Isle of Man

Scientific name	English name	Manx name	Typical grassland type/Notes
<i>Achillea millefolium</i>	Yarrow	Ayr lossey	Neutral, calcareous, coastal
<i>Achillea ptarmica</i>	sneezewort		Wet grasslands
<i>Agrimonia eupatoria</i>	Agrimony		Neutral, calcareous, coastal Rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Agrimonia procera</i>	Fragrant agrimony		Neutral, calcareous, coastal. Hard to tell apart from agrimony, and likely to be mis-recorded for fragrant varieties of agrimony. May escape from gardens
<i>Agrostis capillaris</i>	Common bent	Shaslagh lheannagh	Neutral, acid
<i>Agrostis stolonifera</i>	Creeping bent		Neutral, calcareous, coastal
<i>Agrostis vinealis</i>	Brown bent	Shaslagh	Neutral, acid
<i>Aira caryophylla</i>	Silver hair-grass		Acid, coastal, thin sandy soils
<i>Aira praecox</i>	Early hair-grass		Acid, coastal, thin sandy soils
<i>Alopecurus geniculata</i>	Marsh foxtail		Wet
<i>Anacamptis pyramidalis</i>	Pyramidal orchid		Neutral, calcareous, coastal protected under Schedule 7 of the Wildlife Act 1990
<i>Anagallis tenella</i>	Bog pimpernel		Acid, wet
<i>Anthoxanthum odoratum</i>	Sweet vernal grass	Faiyr sonnys	Neutral
<i>Anthyllis vulneraria</i>	Kidney vetch		Neutral, calcareous, coastal
<i>Arenaria serpyllifolia</i>	Thyme-leaved sandwort		Neutral, calcareous, coastal
<i>Armeria maritima</i>	Thrift		Coastal
<i>Astragalus danicus</i>	Purple milk-vetch		Calcareous, coastal
<i>Campanula rotundifolia</i>	Harebell	Clag ferrish	Neutral, acid. NB. conspicuously large-flowered harebells may be hexaploid, rather than the more usual British tetraploid plants.
<i>Cardamine pratensis</i>	Cuckooflower		Wet
<i>Carduus tenuiflorus</i>	Slender thistle		Coastal
<i>Carex arenaria</i>	Sand sedge		Coastal, sandy ground
<i>Carex binervis</i>	Green-ribbed sedge		Acid
<i>Carex caryophyllaea</i>	Spring sedge		Neutral, calcareous, coastal
<i>Carex echinata</i>	Star sedge		Acid, wet
<i>Carex flacca</i>	Glaucous sedge	Shast (sedge)	Neutral
<i>Carex hirta</i>	Hairy sedge		Neutral, wet
<i>Carex nigra</i>	Common sedge	Shast dhoo	Neutral
<i>Carex ovalis</i>	Oval sedge		Neutral
<i>Carex pallescens</i>	Pale sedge		Neutral Rare; protected under Schedule 7 of the Wildlife Act 1990
<i>Carex panicea</i>	Carnation sedge		Neutral
<i>Carex pulicaris</i>	Flea sedge		Acid, wet
<i>Carex viridula subsp viridula</i>	Yellow-sedge		Acid, wet

<i>Carlina vulgaris</i>	Carlina thistle		Calcareous, coastal, dry ground Rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Catapodium marinum</i>	Sea fern-grass		Bare to thin soils, coastal only
<i>Catapodium rigidum</i>	Fern-grass		Bare to thin soils, coastal
<i>Centaurea nigra</i>	Common knapweed	Lus ny gramman	Neutral
<i>Centaurium erythraea</i>	Common centaury		Coastal, well-drained soils
<i>Centaurium littorale</i>	Seaside centaury		Coastal only; only one doubtful record for the Isle of Man, but possibly present on coastal turf.
<i>Cerastium arvense</i>	Field mouse-ear		Thin, well-drained soils, coastal
<i>Cerastium diffusum</i>	Sea mouse-ear		Coastal only
<i>Cerastium semidecandrum</i>	Little mouse-ear		Thin, well-drained soils, coastal
<i>Cirsium palustre</i>	Marsh thistle		Wet
<i>Cochlearia danica</i>	Danish scurvy-grass		Coastal
<i>Cochlearia officinalis</i>	Common scurvy-grass		Coastal
<i>Conopodium majus</i>	Pignut	Corlan	Neutral
<i>Crepis capillaris</i>	Smooth hawksbeard		Neutral
<i>Cuscuta epithymum</i>	Dodder		Acid, coastal – a parasite of heather, thyme and legumes, mostly found on the coast Rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Cynosurus cristatus</i>	Crested dog's-tail	Connane	Neutral, coastal
<i>Dactylorhiza fuchsii</i>	Common spotted-orchid	Bwoid saggart	Neutral protected under Schedule 7 of the Wildlife Act 1990
<i>Dactylorhiza incarnata</i>	Early marsh orchid		Neutral protected under Schedule 7 of the Wildlife Act 1990
<i>Dactylorhiza maculata ssp. ericetorum</i>	Heath spotted-orchid		Neutral protected under Schedule 7 of the Wildlife Act 1990
<i>Dactylorhiza purpurella</i>	Northern marsh orchid		Neutral protected under Schedule 7 of the Wildlife Act 1990
<i>Danthonia decumbens</i>	Heath grass		Acid, coastal
<i>Daucus carota</i>	Wild carrot		Neutral, calcareous, coastal
<i>Deschampsia cespitosa</i>	Tufted hair-grass		Neutral, wet
<i>Deschampsia flexuosa</i>	Wavy hair-grass		Acid
<i>Erodium cicutarium</i>	Common stork's-bill		Sandy, very dry soils, coastal
<i>Erodium maritimum</i>	Sea stork's-bill		Sandy, very dry soils, coastal
<i>Erophila verna</i>	Whitlow grass		Calcareous, coastal, thin dry soils
<i>Euphrasia spp.</i>	Eyebright		Short, permanent grassland, coastal. A complex aggregate of related species, often hard to tell apart. Semi-parasitic on roots of a range of other

			plants. Indicative of relatively nutrient-poor, well-established grassland.
<i>Festuca ovina</i>	Sheep's fescue		Acid, calcareous, coastal
<i>Festuca pratensis</i>	Meadow fescue		Neutral, damp meadows; native but may sometimes be sown in grassland mixes
<i>Festuca rubra</i>	Red fescue		Neutral, calcareous, coastal
<i>Filago minima</i>	Small cudweed		Neutral, acid, rather bare ground
<i>Filipendula ulmaria</i>	meadowsweet		Wet
<i>Galium palustre</i>	Marsh bedstraw		Wet
<i>Galium saxatile</i>	Heath bedstraw		Acid
<i>Galium verum</i>	Lady's bedstraw	Lus y volley	Neutral, coastal
<i>Gentianella campestris</i>	Field gentian		Neutral, acid, coastal. Intolerant of over-grazing.
<i>Hydrocotyle vulgaris</i>	Marsh pennywort		Wet
<i>Hypochaeris radicata</i>	Common cat's-ear	Cleaysh chiyt	Neutral, coastal
<i>Hypochoeris glabra</i>	Smooth cat's-ear		Acid, sandy, dry ground. Rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Jasione montana</i>	Sheep's-bit	Bossan gorrym	
<i>Juncus acutiflorus</i>	Sharp-flowered rush		Wet
<i>Juncus articulatus</i>	Jointed rush		Wet
<i>Juncus bufonius</i>	Toad rush		wet
<i>Juncus conglomeratus</i>	Compact rush		wet
<i>Juncus effusus</i>	Soft rush		Wet
<i>Juncus foliosus</i>	Leafy toad rush		Wet
<i>Juncus squarrosus</i>	Heath rush		Acid
<i>Knautia arvensis</i>	Field scabious	Lus ny gramman gormey	Neutral
<i>Koeleria macrantha</i>	Crested hair-grass		Calcareous, coastal
<i>Lathyrus linifolius</i>	Bitter-vetch		Acid, damp ground
<i>Lathyrus pratensis</i>	Meadow vetchling	Pishyr bwee	Neutral
<i>Leontodon autumnalis</i>	Autumn hawkbit	Croag phartan	Neutral
<i>Leontodon taraxacoides</i>	Lesser hawkbit		Dry turf, coastal
<i>Listera ovata</i>	Common twayblade		Shady, damp ground. Protected under Schedule 7 of the Wildlife Act 1990
<i>Lotus corniculatus</i>	Common bird's-foot trefoil		Neutral, calcareous, coastal
<i>Lotus pedunculatus</i>	Greater bird's-foot trefoil		Damp grassland
<i>Luzula campestris</i>	Field woodrush	Leaghyr cheylley	Neutral
<i>Luzula multiflora</i>	Heath woodrush		Neutral A species with two types: <i>Luzula multiflora</i> ssp. <i>multiflora</i> and a subspecies with clustered fruiting heads, <i>Luzula multiflora</i> ssp. <i>congesta</i> . The two are not always easy to tell apart, and may grow together on similar sites.

<i>Lychnis flos-cuculi</i>	Ragged robin		Wet
<i>Mentha aquatica</i>	Water mint		Wet
<i>Mentha arvensis</i>	Corn mint		Wet (but prefers drier grassland to <i>M. aquatica</i>)
<i>Mentha pulegium</i>	Pennyroyal		Wet, seasonally inundated grassland. Very rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Molinia caerulea</i>	Purple moor-grass		Acid, wet
<i>Myosotis discolor</i>	Changing forget-me-not		Dry short turf, coastal
<i>Myosotis ramosissima</i>	Early forget-me-not		Dry, thin, infertile soils, coastal
<i>Myosotis secunda</i>	Creeping forget-me-not		Wet
<i>Myosotis laxa</i>	Tufted forget-me-not		Wet, often on trampled ground
<i>Nardus stricta</i>	Mat grass		Acid
<i>Odontites vernus</i>	Red bartsia		Short, often trampled grasslands, coastal
<i>Ononis repens</i>	Rest harrow		Neutral, calcareous, coastal
<i>Ornithopus perpusillus</i>	Bird's-foot		Thin, dry soils, coastal
<i>Pedicularis palustris</i>	Marsh lousewort		Wet
<i>Pedicularis sylvatica</i>	Lousewort		Acid, wet
<i>Phleum arenarium</i>	Sand cat's-tail		Sandy ground, coastal
<i>Phleum pratense</i>	Timothy grass		Neutral
<i>Pilosella officinarum</i>	Mouse-ear hawkweed		Calcareous, thin dry soils, coastal
<i>Pimpinella saxifraga</i>	Burnet saxifrage		Neutral, calcareous, verges Rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Plantago coronopus</i>	Buck's-horn plantain		Thin dry soils, coastal, bare ground
<i>Plantago lanceolata</i>	Ribwort plantain		Any semi-improved grassland, coastal
<i>Plantago maritima</i>	Sea plantain		coastal
<i>Platanthera chlorantha</i>	Greater butterfly orchid		Neutral Rare, Protected under Schedule 7 of the Wildlife Act 1990
<i>Poa pratensis</i>	Smooth meadow-grass		Neutral
<i>Polygala serpyllifolia</i>	Heath milkwort		Acid, coastal
<i>Polygala vulgaris</i>	Common milkwort		
<i>Potentilla anglica</i>	Trailing tormentil		Neutral, acid, damp ground
<i>Potentilla anserina</i>	Silverweed		Disturbed and/or seasonally damp ground, sometimes coastal
<i>Potentilla erecta</i>	tormentil		Acid
<i>Potentilla palustris</i>	Marsh cinquefoil		Wet
<i>Potentilla reptans</i>	Creeping cinquefoil		Neutral, calcareous, coastal
<i>Potentilla x mixta</i> and <i>P. x suberecta</i>	Hybrid cinquefoils		Neutral, acid, verges and banks. Various crosses between tormentil, trailing cinquefoil and creeping cinquefoils (the hybrids are hard to tell apart). Abundant in some unimproved grasslands, even where one or both

			parent species is absent.
<i>Prunella vulgaris</i>	Selfheal	Lus y choine cast	Neutral
<i>Pulicaria dysenterica</i>	Fleabane		Damp grassland
<i>Ranunculus acris</i>	Field buttercup	Cass feeagh	Neutral
<i>Ranunculus flammula</i>	Lesser spearwort		Wet
<i>Ranunculus repens</i>	Creeping buttercup		Heavy soils, disturbed pastures, often damp
<i>Rhinanthus minor</i>	Yellow rattle	"Clabberey traagh"	Neutral Characteristic of traditional hay crop management.
<i>Rosa spinosissima</i>	Burnet rose		Calcareous, coastal
<i>Rumex acetosa</i>	Sorrel		Neutral, acid, damp ground
<i>Rumex acetosella</i>	Sheep's sorrel		Acid, coastal, well-drained thin soils
<i>Sagina subulata</i>	Heath pearlwort		Dry, open ground, coastal
<i>Scilla verna</i>	Spring squill		Coastal
<i>Sedum anglicum</i>	English stonecrop		Dry ground, coastal
<i>Senecio aquaticus</i>	Marsh ragwort		Damp soils. Treated as ragwort (above).
<i>Senecio jacobaea</i>	Ragwort	cushag	Dry, often disturbed sandy soils. Of ecological importance, but poisonous to stock and listed under the Weeds Act
<i>Silene uniflora</i>	Sea campion		Coastal
<i>Spergularia rupicola</i>	Rock sea- spurrey		Coastal
<i>Spiranthes spiralis</i>	Autumn ladies'- tresses		Calcareous, coastal. Manx record for this plant is the northernmost in the British Isles. Very rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Stachys palustris</i>	Marsh woundwort		wet
<i>Stellaria graminea</i>	Lesser stitchwort		Neutral, acid, damp ground
<i>Stellaria uliginosa</i>	Bog stitchwort		Acid, damp ground
<i>Succisa pratensis</i>	Devil's-bit scabious	Bit y jouyll	Neutral, wet
<i>Thymus polytrichus</i>	Wild thyme		Thin soils, coastal
<i>Trichophorum cespitosum</i>	Deer-grass		Acid pasture and heath
<i>Trifolium fragiferum</i>	Strawberry clover		Damp ground, coastal
<i>Trifolium ornithopodioides</i>	Bird's foot clover		Thin soils, coastal
<i>Trifolium pratense</i>	Red clover	Shamrag ny gabbil	Neutral
<i>Trifolium scabrum</i>	Rough clover		Thin soils, coastal
<i>Trifolium striatum</i>	Soft clover		Dry ground, coastal
<i>Trifolium suffocatum</i>	Suffocated clover		Thin soils, coastal. Manx records are the most northerly in the British Isles.
<i>Trifolium arvense</i>	Hare's-foot clover		Dry sandy ground, coastal Rare, protected under Schedule 7 of the Wildlife Act 1990
<i>Triglochin maritimum</i>	Sea arrowgrass		Saline wet ground, coastal only
<i>Triglochin palustris</i>	Marsh		Usually calcareous or neutral, wet

	arrowgrass		
<i>Valeriana officinalis</i>	Valerian		Neutral, usually on wet sites
<i>Vicia cracca</i>	Tufted vetch	Pishyr lughag	Neutral
<i>Vicia lathyroides</i>	Spring vetch		Thin, sandy soils; coastal
<i>Viola canina</i>	Heath dog-violet		Acid, coastal
<i>Viola lutea</i>	Mountain pansy		Acid, usually upland. Very rare, possibly extinct. Protected under Schedule 7 of the Wildlife Act 1990
<i>Viola tricolor ssp curtisii</i>	Seaside pansy		Sandy ground, coastal. Rare.
<i>Viola tricolor ssp tricolor</i>	Heartsease		Acid, sandy ground, coastal
<i>Wahlenbergia hederacea</i>	Ivy-leaved bellflower		Acid, heathy, damp pastures



Moonwort *Botrychium lunaria*, a tiny fern of short grassland, with a pound coin for comparison

4.3 Lowland heathland

Description

Lowland heathland is restricted to the more acid, nutrient-poor soils of the British Isles and parts of North-west Europe, where woodland has been cleared in the past and the resultant low-growing shrub layer is maintained by human activities – usually extensive grazing and rotational burning. This management favours species such as heathers, whose ecological niche would otherwise be limited to exposed coastal sites, or woodland areas which are recovering from wildfire. Like other habitats which have formed as a result of very long-established human activity, lowland heathland supports a range of distinctive ecological communities with plants and animals which rely on the managed habitat for their survival. The majority of lowland heaths therefore occupy areas which would have supported woodland in prehistoric or historic times, and which still have a strong tendency to revert to woodland in the absence of management.

For most of the UK, “lowland” is defined as altitudes less than 300m; in the case of northerly and coastal regions, the vegetation tends to progress to upland types at a lower altitude. Thus, in the case of the Isle of Man, 250m is a more useful cut-off point for defining lowland as opposed to upland habitats. Most land above this altitude has not recently been subject to enclosure, and consists of upland heathland types, semi-improved grassland, and plantations. Some areas of upland type moorland do occur at lower altitudes, but such instances are generally limited to the lower margins of larger, upland blocks of heather. For the purposes of these criteria, the upper defining altitude for lowland heathland will therefore be taken as 250m.

Most lowland heathland consists of a low to medium growth of heather *Calluna vulgaris*, with varying amounts of bell heather *Erica cinerea*, cross-leaved heath *Erica tetralix*, gorse *Ulex spp* and bilberry *Vaccinium myrtillus*. Specialised local conditions may also give rise to abundant lichen and/or moss communities. In the Isle of Man, the native gorse is *Ulex gallii* (“Western” or “Manx gorse”), but the introduced European gorse *Ulex europaea* also occurs in profusion on some lowland heathland areas, especially in places where there has been some enrichment of the soil for agricultural purposes. Heathland management is usually aimed at maintaining a high proportion of heather species rather than either type of gorse, heather having a generally higher value for extensive grazing as well as for conservation and the landscape. However, lowland heathland needn’t just consist of a carpet of heather: interspersed areas of developing trees and scrub, and microhabitats such as seepages and bogs, are important elements of a diverse site.

Heathland in general is not very floristically diverse (Ratcliffe, 1977), but this is partly because management efforts tend to encourage single-stage heather growth. Where heathlands support a range of habitat types, particularly wet habitats, along with a diverse structure of different stages of heather growth, they may have a striking range of birds, plants and invertebrates, some of them rare and restricted in range. Details of some of the species which occur on the best examples of lowland heathland on the Island are listed below in Sections 4.3.1-4.3.3.

In European terms, lowland heathland is largely oceanic in distribution, occurring where the temperature range is steady and the climate relatively mild in terms of temperature and dryness. The maximum altitude of lowland heathland, as opposed to upland and upland margins, is variously defined as between 250m and 300m (RSPB, 2005; English Nature 2002). In the British Isles, lowland heathland tends to occur further inland as well as at the coast, probably due to the generally temperate conditions (Ratcliffe, 1977). In this respect the Manx situation mirrors that of the British Isles as a whole (BDIS).

Lowland heathland is a priority for nature conservation because it is a rare and threatened habitat. The UK has an important proportion (about 20%) of the international total of this habitat - 58,000ha, of which the largest proportion (55%) is found in England. However, even in England only one sixth of the heathland present in 1800 now remains (UK BAP).

The range of European heathland types has been categorised by different organisations in different ways, leading to a multitude of overlapping descriptions, criteria and statistics (see **Types of lowland heathland**, below). For the sake of clarity, this chapter covers all lowland heathland types, including coastal and dune heathland, and both wet and dry habitats. Other coastal habitats – which may occur in conjunction with coastal and dune heathland – are covered in **Chapter 4.6: Coastal habitats**. **Upland heathland** is treated alongside related upland habitats in **Chapter 4.7**.

Assessing lowland heathland

In general, the same principles of site assessment apply to lowland heathland as to grasslands: all of the Priority Site Criteria listed in Part 3 are relevant. Of these, the following are particularly important when assessing the quality of lowland heathland:

Size

The suitability of much of the Island's lowland heathland for agricultural improvement has led to loss and fragmentation of the habitat just as it has in the UK (see JNCC Annex 1 habitat descriptions). Thus it may not always be possible to protect the kind of extent of habitat that would be possible with upland heathland; indeed, it may be necessary to protect relatively small lowland heathland areas just in order to represent all the types and species which are associated with this habitat on the Island. However, larger sites are more likely to contain the diversity of microhabitats and conditions which favour biodiversity, and for this reason it is always desirable to select the largest available examples.

Location

The varied geology, topography and climate on the Island has led to a perhaps surprising range of heathland types given the limited extent of the land. In order to represent this range in full, the ASSI series should include lowland heathland examples from both coastal and inland sites, and from the full range of geological and hydrological situations. Natural climax heathlands are regarded as having especially high scientific value (Ratcliffe, 1977), so sites such as established dunes at the Ayres, where the heathland at least on the seaward side is likely to be a natural climax vegetation, are particularly important.

Another location consideration is the presence of species or habitat sub-types which are known to be at the edge of their natural range. Such species are of value in monitoring the effects of climate change, as well as being of interest in their own right. The geographic position of the Isle of Man and the ameliorating influence of the sea on its climate have resulted in the development of several habitat types which otherwise tend to occur further south. This is particularly marked in the case of lowland heathland, most types of which are largely confined to England and South-west Wales (see <http://www.forestry.gov.uk/forestry/Lowlandheath>). Conversely, some species may be at the southernmost end of their range in the Isle of Man. Such species are at risk of losing their Island station if climate change forces them further north. Details of some of the habitat sub-types and species which are thought to be at the edge of their native range on the Isle of Man are listed in sections 4.3.1 – 4.3.3, below.

Diversity

Diversity of habitat and micro-habitat is one of the most important features of good lowland heathland examples. The natural cycle of heather is one of structural, and hence ecological, development over a period of twenty years or more, with young, vigorous shoots giving way to uniform bushes which then gradually grow more "leggy", to collapse and be overgrown by new shoots. Each stage is of importance for different bird, invertebrate and lower plant species. Regenerating woodland is also a natural feature of lowland heath, although most management efforts will aim to prevent the heather layer from disappearing altogether.

Other features which lead to greater biodiversity include pools and mires, streams, mature hedgerows, and patches of grassland. With this in mind, **the assessment process should aim to protect the full range of Manx lowland heathland biodiversity by recognising good examples of varied mire/heath/grassland mosaic habitats as well as the best large areas of a single heathland type.**

Rarity, fragility and vulnerability

The rarity of lowland heathland, and the limited distribution of types such as dune heathland, make this a prime habitat for rare and local species on the Island. Even if the overall site is small, it may therefore be important to protect it if it contains the sole or dominant population of a rare species. Other factors to take into account are the presence of rare micro-habitats – e.g. pools which support breeding dragonflies – and the vulnerability and fragility of the site – e.g. lichen heath, which may take a very long time to regenerate if damaged. Heathlands are particularly vulnerable to changes in nutrient status and can be lost through gradual changes to grassland through inappropriate grazing management.

Naturalness and representativeness

Selecting the most natural and representative lowland heathland depends a lot on assessments of habitat diversity on site (see Diversity, above). The most natural examples are likely to have a range of stages of heather growth, a mix of different hydrological features, and some associated habitats such as regenerating woodland and scrub. A clue to the naturalness of a lowland heathland might be the presence of a range of hydrological features such as wet heath patches and pools; this would indicate that the area has not been subject to drainage recently. Sites with single-age “monoculture” stands of heather, whilst important for some bird species, are less likely to fulfil the diversity and naturalness criteria.

Ecological position

The tendency of heathland to form an “intermediate” stage between cleared areas and woodland means that it often occurs amidst other habitats, particularly grassland and scrub. A mosaic of habitats, particularly those which represent a natural gradation from one heathland type to another, can greatly increase the biodiversity of a lowland heathland site, offering opportunities for species to shift and adapt with changes in conditions, and providing valuable “edge” habitats, such as the interface between drier heathland and mire. Sites which exhibit good gradients of microhabitat types should be a priority for selection, as these features are often indicators of the most natural lowland heathland sites.

History

Although there are probably small examples of naturally-occurring climax heathland on the Island, the majority of both lowland and upland heathland appears to have developed in response to human activities. A history of appropriate management is therefore likely to have a bearing on the current quality of a lowland heathland site, and may also indicate where the most natural areas – e.g. those which have not been ploughed, drained or limed in the past – remain. Conversely, sites where the management has plainly not been supportive of high biodiversity are likely to need some remedial management if their full conservation potential is to be achieved; such sites may be an important part of any action plan to increase the resource of lowland heathland on the Island, but they should not normally be prioritised for protection ahead of heathlands which have a history of good management.

Types of lowland heathland

A straightforward classification of lowland heathlands and their status is difficult. The range of heathland types around the British Isles has led to a number of classification approaches, some of which refer to vegetation communities, whilst others relate to distribution and altitude.

This can lead to problems when attempting to compare the conservation status of heathland types. For example, in the case of coastal heath:

- The National Vegetation Classification (NVC) includes dune heath dominated by heather and sand sedge *Carex arenaria*, and describes it as a vegetation type of coastlines and also some inland dune systems (Rodwell, 1992⁵);
- Annex 1 of the EC Habitats Directive lists *Atlantic decalcified dunes* as a category which includes some heathland elements along with other species more typical of mobile and semi-fixed dunes;
- The UK Biodiversity Action Plan lists Lowland heathland as a Priority Habitat, regardless of whether it is coastal or inland, wet or dry, but also lists Coastal dune systems as a separate Priority Habitat, which may have some heathland elements present;
- The DAFF Phase 1 Habitat Survey treats Dry dwarf shrub heath as a category regardless of whether it is upland or lowland, but treats coastal heath as a separate category even though it, too, is strictly a type of dry dwarf shrub heath.

Remaining areas and rates of loss have been calculated for several of the classes of habitat above, but determining how these statistics relate to practical categories such as “all coastal dune heath” or “all dry lowland heath” can be difficult. Perhaps because of the choice of ways in which lowland heathland has been categorised, most of the available statistics about the status of the habitat are general, rather than for individual sub-types such as dune or wet heathland. Given that several broad types of heathland may occur adjacent to each other in any case, this document also takes a broad view of lowland heathland status – see **Overall status of lowland heathland**, below.

For the purpose of determining detailed minimum criteria for site selection, and bearing in mind the above points about other ways of classifying heathland, lowland heathlands on the Isle of Man have been grouped as follows:

Dry lowland heathland – heather-dominated habitats, found in scattered inland areas up to the edge of the hill-land, often on enclosed fields and up to an altitude of around 300m;

Wet lowland heathland – heather and cross-leaved heath-dominated habitats, often in conjunction with mires, restricted to small, poorly-drained areas of drier heaths, valley mires, and amongst marshy grassland.

Coastal heathland – heather- and Manx gorse-dominated habitats on cliffs and adjacent to coastal slopes, and heather- and lichen-dominated fixed, decalcified dunes.

These heathland types, and the National Vegetation Classification (NVC) categories they most closely resemble, are listed in detail in sections 3.4.1 – 3.4.3 below.

Overall status of lowland heathland

"Open heathland is rarer than rainforest. In the UK we have only about 16% left of the area that existed in 1800. That means that from an area similar to the size of Cornwall, only the equivalent of the Isle of Wight remains. The process of loss and disintegration has been particularly fast in recent decades. However, this country still holds 20% (more than 60,000 hectares) of the whole world's lowland heathland. Thus there is a need, not only to preserve and improve our remaining heathlands, but if possible, to re-create them in areas where they have recently been lost.

There is a special case for linking small fragments of heathlands, where the few remaining species are stretched for space and risk disappearing in the event of a fire, to create areas which can maintain a wider range of wildlife and can survive in the future."

(from English Nature, 2002)

Lowland heathland on the Isle of Man tends to be restricted to small, often isolated patches of inland farmland, or narrow zones along the coastline. The DAFF Phase 1 Habitat Survey does not distinguish between upland and lowland dwarf shrub heath, but the area of heathland below 250m may be ascertained from the mapping as less than one tenth of the overall heathland area (BDIS). Dry dwarf shrub heath is one of the commonest semi-natural habitats on the Island, but a glance at the habitat maps reveals that very little of this falls within lowland areas – around 6% is estimated to occur at altitudes under 250m. Wet dwarf shrub heath is rare in any case, with just 310ha, or 0.55% of the Island, in total (much of which also appears to be in the uplands) (Sayle *et al*, 1995). Only coastal heaths were measured separately, accounting for 322ha, most of it dune heathland at the Ayres. It may therefore be inferred that the rarity of both wet lowland heath and coastal heath throughout Europe is mirrored on the Island. The importance of these habitats is reflected in the Wildlife Sites Handbook, where all areas of lowland heathland which have at least 10% cover of heathers, preferably over 1ha in extent, should be considered for selection as Wildlife Sites (Keehan, 1999), along with **all** examples of coastal heathland which have more than 25% dwarf shrub cover.

Lowland heathland is vital for four bird species of high UK conservation concern: stone-curlew, nightjar, woodlark and Dartford warbler. Other birds of importance on heathlands include linnet (a Red List species), and stonechat, short-eared owl and curlew (Amber List species). The latter species all occur on heathland on the Isle of Man; short-eared owl and curlew are both protected under Schedule 1 of the Wildlife Act 1990. Heathland is also very important for insects, some of which are dependent on the

habitat. Heathland pools and streams support most of the UK's 38 species of dragonfly (RSPB, 2005). Manx invertebrate species of interest include lesser bee-fly *Bombylius minor* on coastal heathland and lesser mottled grasshopper *Stenobothrus stigmaticus* on coastal heathland/grassland, both of which are protected under Schedule 5 of the Wildlife Act 1990.

In the past, heathland in the UK was lost primarily to agriculture, forestry, mineral extraction and development. Uncontrolled burning has also been a particular threat to bryophyte and lichen-rich heathland. The UK Biodiversity Action Plan lists Lowland heathland as a Priority Habitat, and gives the main factors affecting the habitat at present as:

- Encroachment of trees and scrub and the simplification of vegetation structure due to a lack of conservation management such as light grazing, controlled burning and cutting.
- Nutrient enrichment, particularly deposition of nitrogen compounds emitted from intensive livestock farming, or from other sources.
- Fragmentation and disturbance from developments such as housing and road constructions.
- Agricultural improvement including reclamation and overgrazing, especially in Northern Ireland.

These factors apply in varying degree to the Isle of Man and indeed other parts of Europe (JNCC).

The scarcity of lowland heathland in the British Isles, and the responsibility of protecting one of the world's rare habitats, has led to a corresponding amount of conservation effort. For example, the Ministry of Defence – who own around 30% of the remaining UK lowland heathland - is represented on the UK Lowland Heathland Habitat Action Plan Steering Group, and has produced a Lowland Heathland Biodiversity Action Plan. A relatively high proportion of the lowland heathland resource in the UK has been notified as a Site of Special Scientific Interest SSSI under the Wildlife and Countryside Act 1981.

In the European context, both upland and lowland heaths are well-represented in the SAC series, and there is a high proportion of the rarer and more species-rich lowland types, which are particularly threatened by human activities (JNCC). Whilst not treated as a single habitat type (unlike the UK BAP), the EU Habitats Directive lists the following types, of which some or all occur on lowland heathlands:

- **European dry heaths** (much of this habitat is in the uplands)
- **Northern Atlantic wet heaths with *Erica tetralix*** (much of this habitat is also upland)
- **Atlantic decalcified fixed dunes** (usually occurs at around sea-level)

Lowland heathland may also be found within Ramsar Sites, especially where they form part of a lowland wet peatland system.

Where information is available for individual lowland heathland types, they are noted in the Status sections of parts 4.3.1 – 4.3.3, below.



Bell heather *Erica cinerea*, growing with "Spaghetti lichen" *Cladonia portentosa* in coastal heath

4.3.1 Dry lowland heathland

Description and Distribution

Dry lowland heaths typically occur on well-drained, acidic to circumneutral soils with low nutrient content. Heathers and other dwarf-shrubs dominate the vegetation. The most common is heather *Calluna vulgaris*, which often occurs in combination with gorse *Ulex* spp., bilberry *Vaccinium myrtillus* or bell heather *Erica cinerea*, though other dwarf-shrubs are important locally. Nearly all dry heath is semi-natural, being derived from woodland through a long history of grazing and burning. Most dry heaths are still managed as extensive grazing for livestock. Criteria for dry heaths in upland areas – which make up a high proportion of all the semi-natural vegetation on the Island – are listed in **Chapter 4.7: Upland habitats**.

At low to moderate altitudes in warm oceanic parts of southern Britain, the typical NVC form of the habitat is H8 heather-gorse *Calluna* – *Ulex* heath, characterised by abundant heather *Calluna*, Western/"Manx" gorse *U. gallii* and bell heather *Erica cinerea*. Western gorse becomes scarce in the cooler oceanic climate further north. It should be noted that this particular species of gorse is an important constituent of oceanic and southern heath in the British Isles (NCC, 1989), hence site selection on the Isle of Man needs to ensure that it is well-represented despite its apparent abundance here, not least because it is near the edge of its northerly native range.

Heather and bell heather are abundant together in H10 *Calluna* – *Erica* heath, especially on more southerly-facing slopes; bell heather *E. cinerea* becomes dominant in the hyper-oceanic fringes of the north-west, but in most cases appears to be the less prevalent species in Manx lowland heaths. In this respect they can resemble those of England, and as such are probably also at the northernmost limit of their range.

Dry lowland heathland is of high importance for birds. Many moorland birds feed and/or breed on dry lowland heathland, including hen harrier, nightjar, kestrel, merlin and short-eared owl – all of which are protected under Schedule 1 of the Wildlife Act 1990.

Classification of dry lowland heathland

Dry lowland heaths on the Island may be classified broadly as outlined in **Table 4.3.a**:

Status of dry lowland heathland

The DAFF Phase 1 Habitat Survey does not distinguish between dry dwarf shrub heath on lowland as opposed to upland areas. Overall, it is a common habitat on the Island; however, lowland examples only comprise around 6% of the total habitat resource (BDIS). Dry lowland heathland on the Isle of Man is not only rare, but also exceedingly fragmented. Whilst the DAFF Phase 1 Habitat GIS records 365 individual parcels of dry lowland heathland (i.e. under 250m altitude), over three-quarters of the total area of the habitat on the Island occurs within just twenty sites – only five of which are greater than 10ha. Added together, the hundreds of examples which are less than 0.5ha in extent (often much less) only make up 10% of the total area of the habitat. The Island is not alone in this respect: whereas dry heaths occur throughout the UK, and are particularly abundant in the uplands, where they may dominate the landscape, they are more localised and fragmented in lowland areas, especially in south and central England, where they have declined in extent due to afforestation, agricultural improvement and other factors (JNCC).

European dry heaths are an Annex 1 habitat under the EU Habitats Directive. They are found in every EU Member State except for Greece, but are only extensive in the western oceanic fringes of Europe. A high proportion of the EU resource of European dry heaths occurs in the UK, although this proportion is not as high as that of the other Annex 1 habitat, Northern Atlantic wet heaths with *Erica tetralix* (see 4.3.2, below). Dry heaths in the UK exhibit exceptional diversity in comparison with examples found elsewhere in the EU. (www.JNCC.gov.uk, 2005).

For more details see **Overall status of lowland heathland**, above.

Table 4.3.a: Broad categories of dry lowland heathland on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (see Rodwell 1992²)	Rarities present?	Protected species present?
Manx gorse heath – often found near the coast, especially in the south of the Island, but not necessarily right next to the sea.	Not usually	H8a & b <i>Calluna vulgaris</i> – <i>Ulex gallii</i> (heather and western or “Manx” gorse) heath Six Island NVC samples, all within the species-poor or <i>Danthonia decumbens</i> (a moss) sub-communities. One other sub-community may be present on the coast – see Table 4.3.c, below.	Not usually	Yes. Protected bird species are likely, including, kestrel, merlin and possibly feeding chough.
Heather heath – occurs throughout, including upland fringes. Often subjected to burning and liming.	Not usually	H10 <i>Calluna vulgaris</i> – <i>Erica cinerea</i> (heather and bell heather) heath One Island NVC sample	Not usually; rarely supports unusual clubmosses such as fir clubmoss <i>Huperzia selago</i> and stag’s-horn clubmoss <i>Lycopodium clavatum</i>	Yes. Protected bird species are likely, including hen harrier, kestrel, merlin and short-eared owl. Potential for nightjar though records rare now.

Minimum selection criteria for dry lowland heathland

- All areas of dry lowland heath greater than 5ha in extent should be considered for selection;
- Any smaller areas of dry lowland heath should be considered for selection where they form part of a mosaic of semi-natural habitats of 2ha or more in extent, and the site as a whole satisfies at least one of the Priority Site Criteria listed in Part 3.

4.3.2 Wet lowland heathland

Description and Distribution

Wet heathland usually occurs on acidic, nutrient-poor substrates, such as shallow peats or sandy soils with impeded drainage. The vegetation is typically dominated by mixtures of cross-leaved heath *Erica tetralix*, heather *Calluna vulgaris*, grasses, sedges and *Sphagnum* bog-mosses. It is highly likely to occur in conjunction with other habitats such as dry lowland heath, marshy grassland, mire and bog.

M15 *Scirpus* – *Erica* wet heath is found in areas with a moderate to high rainfall, and is the typical form of wet heath in the north and west of the UK. *E. tetralix* and *Calluna* are typically accompanied by abundant deer grass *Trichophorum cespitosum* and purple moor-grass *Molinia caerulea*. Where there is movement of mildly base-rich water through the peat, sedges *Carex* spp. and a wide range of species favoured by flushing occur. The latter include distinctive variants that are often characterised by abundant bog-myrtle *Myrica gale*, or black bog-rush *Schoenus nigricans*. These habitat variants are rare on the Island, as elsewhere.

M16 bell heather/*Sphagnum* moss *Erica* – *Sphagnum* wet heath is characteristic of drier climates in the south and east of the UK, and is usually dominated by mixtures of cross-leaved heath *Erica tetralix*, heather *Calluna* and purple moor-grass *Molinia caerulea*. The bog-moss *Sphagnum compactum* is typically abundant. Both the above habitats are relatively diverse and colourful compared with many dry heaths, and may support good populations of protected species such as heath-spotted orchid *Dactylorhiza maculata ssp ericetorum*.

Wet heaths occur in several types of ecological gradient. In most lowland areas they are very local, and often restricted to the transition zone between dry lowland heaths and constantly wet valley mires. This effectively means that the majority of wet lowland heathland on the Island is restricted to many very small patches within larger semi-natural areas. Similar habitat in the uplands occurs most frequently in gradients between dry heath or other dry, acid habitats and blanket bogs – **see Chapter 4.7: Upland habitats**.

Classification of wet lowland heathland

Please note that in upland margin areas, some wet heathland types may occur which are more typical of uplands; these may be important habitats to include as part of the gradient between lowland and upland heathlands on some sites. They are described in more detail in Chapter 4.7: Upland habitats. Strictly lowland wet heathland on the Island may be classified broadly as outlined in **Table 4.3.b:**

Table 4.3.b: Broad categories of wet lowland heathland on the Isle of Man

Type and Description	Species rich?	Closest NVC categories	Rarities present?	Protected species present?
Cross-leaved heath wet heath Actually both NVC types listed here are a type of mire; other mire types with less heathy characteristics may occur nearby (see Chapter 4.4 Bogs, fens and flushes).	Yes, often	M15 <i>Scirpus cespitosus</i> – <i>Erica tetralix</i> (deer grass and cross-leaved heath) wet heath No Island NVC sample M16 <i>Erica tetralix</i> - <i>Sphagnum compactum</i> (cross-leaved heath and <i>Sphagnum</i> moss) wet heath No Island NVC sample	May support rare invertebrates and/or plants	Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing. A likely habitat for cranberry <i>Vaccinium oxycoccus</i> , heath-spotted orchid <i>Dactylorhiza maculata ssp ericetorum</i> and pale butterwort <i>Pinguicula lusitanica</i> .

Wet lowland heathland status

The DAFF Phase 1 Habitat Survey does not distinguish between wet dwarf shrub heath on lowland as opposed to upland sites; hence it is difficult to discern the total extent of the habitat on the Island. Unlike dry dwarf shrub heath, which is concentrated in the uplands, wet heath occurs in roughly equal amounts in both upland and lowland habitats, if “upland” is taken as any habitat over 250m. Wet lowland heathland is a very fragmented habitat on the Island, with just 11 out of 81 sites containing 75% of the total habitat area. The remaining 25% is scattered throughout 70 separate blocks of wet lowland heathland, all under 2ha in extent, and most under 1ha (BDIS).

Wet heath is extensively developed in western Britain and there are large areas of poor-quality habitat which do not meet the criteria for SAC selection; however, the vast majority of this is in upland areas. One exception is **Northern Atlantic wet heaths with *Erica tetralix***, which are an Annex 1 habitat under the EU Habitats Directive. They are restricted to the Atlantic fringe of Europe between Norway and Normandy. A high proportion of the EU resource occurs in the UK, where it is restricted to highly localised sites in parts of southern and central England (JNCC). Small fragments of this habitat type occur on lowland sites around the Isle of Man, usually in conjunction with valley mires and fens.

Wet heath is an important habitat for a range of vascular plant and bryophyte species of an oceanic or Atlantic distribution in Europe, several of which have an important part of their EU and world distribution in the UK. The UK is one of the most important parts of the world for Atlantic bryophytes, and this is the most important habitat for many of these oceanic species of restricted world distribution (www.JNCC.gov.uk, 2005). The location of the Island would suggest that it may contribute to the UK's high proportion of bryophyte-rich wet heath; however, the full extent of bryophyte diversity in wet heaths on the Isle of Man is not well known at present. It is therefore an area which requires further study.

Minimum selection criteria for wet lowland heathland

- All areas of wet lowland heathland over 0.5ha in extent which fulfil at least one of the Priority Site Criteria listed in Part 3 should be considered for selection;
- Smaller areas of wet lowland heathland which occur in conjunction with other semi-natural habitats totalling 1ha or more should be considered for selection where the site as a whole satisfies at least one of the Priority Site Criteria listed in Part 3. Sites which consist of a range of related wetland habitats should here be considered a top priority.

4.3.3 Coastal heathland

Description and Distribution

Coastal heathland around the Isle of Man usually takes one of two forms: dune heathland on fixed, decalcified dunes (i.e. those where the calcium content of the shell sand has been leached out by rain, leaving an acidic sandy soil), or gorse-dominated heath on rocky cliffs and slopes adjacent to the sea.

Dune heathland occurs on mature, stable dunes where the initial calcium carbonate content of the dune sand is low. The surface soil layers rapidly lose their remaining calcium carbonate through leaching, and become acidified. The most characteristic community is dune heath in which heather *Calluna vulgaris* is found in combination with sand sedge *Carex arenaria*. The main species present vary considerably throughout the UK. Bell heather *Erica cinerea* is abundant in dune heath on west coast sites, but much scarcer in the east, where heather *Calluna vulgaris* predominates. Manx dune heathland appears to be more or less intermediate in this respect. Associated with the heathland in drier conditions, and forming transitions with it, is acidic dune grassland. This is often a precursor to heath development and in these drier conditions is usually characterised by a combination of the fine-leaved grasses such as sheep's-fescue *Festuca ovina* and common bent *Agrostis capillaris*, with sand sedge *Carex arenaria*.

On very nutrient-poor sand that is also subject to severe drought, dune heath tends to be dominated by lichens, which form a continuous grey carpet over the ground. Such low-growing habitats may depend on rabbit-grazing for their survival, and can become overgrown quite rapidly if grazing pressure is reduced. The Isle of Man has an unusual regional example of this kind of lichen heath: for a short section of the Ayres coastal dune system, the shorter areas of heath are dominated by lichens, principally *Cladonia* spp., and sausage lichen *Usnea articulata* – a delicate species, listed as Near threatened in the UK (JNCC), and more usually found growing on trees in sheltered woodland and orchard margins. The relatively mild conditions on the Manx coastline appear to suit it. As such it represents both a unique sub-type of habitat, and a valuable indicator of low background pollution levels, as *U. articulata* is highly sensitive to sulphur dioxide, and is now extinct in most of its former range in England (DEFRA air pollution factsheet, 2005).

Dune heathland almost always occurs in a gradient with other coastal dune habitats, flanked on the inland side by other dry lowland heathland types, or by cultivated farmland. This chapter may therefore need to be read in conjunction with **Chapter 4.6: Coastal habitats**.

Rocky coastal heathland consists of maritime heather *Calluna vulgaris* and spring squill *Scilla verna* heaths. These are usually characterised by abundant Western or “Manx” gorse, *Ulex gallii*. Whilst a familiar and common species on the Island, Manx gorse is actually a key element of one of the more restricted types of coastal heath in the UK, represented here at the northernmost end of its natural range. This kind of coastal heath is at its most diverse when in conjunction with other, related habitats, with which it often forms a gradient – such as coastal grassland and scrub.

Classification of coastal heathland

Bearing in mind the comments about classification made in the introduction above, coastal heathland on the Island may be classified broadly as outlined in Table 4.3c.

Table 4.3.c: Broad categories of coastal heath on the Isle of Man

Type and Description	Species rich?	Closest NVC categories	Rarities present?	Protected species present?
<i>Rocky coastal heath</i> A type of heath which is notable for being a natural climax community in most of the places where it occurs; often found on cliff tops and slopes within range of sea spray.	May be species rich, especially where found in conjunction with coastal grassland on base-rich sites.	H7 <i>Calluna vulgaris</i> – <i>Scilla verna</i> (heather and spring squill) heath No Island NVC sample H8d <i>Calluna vulgaris</i> – <i>Ulex gallii</i> (heather and Manx gorse) heath, <i>Scilla verna</i> (spring squill) sub-community No Island NVC sample; likely to be at the northern limit of its range	Yes, especially when in conjunction with coastal grassland; species may include species such as purple milk-vetch <i>Astragalus danicus</i> , and see right.	May form part of chough feeding territory. Possible protected plants include Autumn ladies tresses <i>Spiranthes spiralis</i> , Portland spurge <i>Euphorbia portlandica</i> , and spring sandwort <i>Minuartia verna</i> . Lesser mottled grasshopper <i>Stenobothrus stigmaticus</i> in coastal heathland/grassland.
<i>Dune heath</i> Heathland generally forms at some distance from the sea; lowland heathland types, including coastal heath, are described in more detail in Chapter 4.3 .	Yes – particularly in lichens and invertebrates, but also in vascular plants.	H11 <i>Calluna vulgaris</i> – <i>Carex arenaria</i> (heather and sand sedge) heath One Island NVC sample – at the southern end of the range of this habitat. May grade into H10 <i>Calluna vulgaris</i> – <i>Erica cinerea</i> (ling and bell heather) heath on the landward side.	A very specific lichen habitat, supporting unusual species such as sausage lichen <i>Usnea articulata</i> . Unusual plants include suffocated clover <i>Trifolium suffocatum</i> , and see right.	Offers good habitat for lizards <i>Lacerta vivipara</i> . Hare’s-foot clover <i>Trifolium arvense</i> , and orchids may be present. Protected invertebrates known to occur on this habitat include heath bee-fly <i>Bombylius minor</i> .

Coastal heathland status

Coastal heath is listed separately from other lowland heathland types in the DAFF Phase 1 Habitat Survey. The actual overall area of coastal heath measured is very low; dune heathland, restricted to the Ayres, amounts to just 0.55% of the Island (311ha), and lichen/bryophyte heath (defined there as having less than 30% vascular plant cover) is only 3.12ha in total. Other coastal heath is restricted to rocky cliff tops and slopes, and comes to 11.44ha. Whilst this last figure may partly be low due to false representation of steep habitats on the orthographic mapping system, it is nevertheless clear that coastal heath is not common on the Island, and is restricted to narrow strips. Nor is it a continuous feature as might be expected, many areas of coastline having now been cultivated right up to the edge of the cliff-top. In recognition of the rarity and vulnerability of coastal heathland of all kinds, the Manx Wildlife Sites Handbook recommends that all areas of coastal heath should be eligible for Wildlife Site selection.

In the UK, both sand dunes and other coastal heath have been recognised as important, with a high proportion of sand dune sites designated as SSSI, or ASSI in Northern Ireland. All coastal sand dunes are a UK BAP Priority Habitat.

In the European context, Atlantic decalcified fixed dunes (*Calluno-Ulicetia*) are not just an Annex 1 habitat under the EU Habitats Directive, but are also listed as a Priority Feature. Atlantic decalcified fixed dunes are widespread in Europe, and the UK lies at the centre of the north – south range of this habitat type (JNCC). The dune heathland at the Ayres fits this category.

Minimum selection criteria for coastal heathland

- All examples of dune heath should be considered for selection; top priority should be given to lichen-rich examples, with the aim of protecting all examples of lichen heath with *Usnea articulata* as ground cover;
- All examples of coastal heath on rocky cliffs and slopes should be considered for selection where the site in question satisfies at least one of the Priority Site Criteria listed in Part 3; further priority should be given to sites where the heathland occurs in natural proximity to other coastal habitats such as coastal grassland and scrub.



Coastal heathland at Langness ASSI

4.4 Bogs, fens and flushes

Description

Bogs, fens and flushes are all types of **mire** - a broad term used to describe waterlogged habitats where water is permanently at or just below ground level. Mires may result from a constant flow of surface water, or, more commonly, impeded drainage. In either case, the waterlogged conditions can lead to a range of problems for plants, such as low soil oxygen levels, high acidity and limited nutrient availability. Because of this, mires tend to contain specialised vegetation communities which are well adapted to the difficult conditions, but very sensitive to change - and hence vulnerable to loss through agricultural improvement and drainage. In spite of all this, mires can support a surprisingly high degree of biodiversity. They are of importance for many rare and delicate plants, and they also provide a home for specialised invertebrates and a range of upland and lowland birds, most notably waders.

Mire habitats contribute a very distinctive element to both upland and lowland landscapes, with purple moor-grass pastures full of bog asphodel and heath spotted-orchids, cotton-grass bogs, and their resident calling curlews all ranking amongst the most popular and evocative elements of the Manx countryside. Unfortunately, mires throughout the British Isles are vulnerable to loss and damage, particularly from commercial peat extraction and agricultural improvement. The high rate of loss of these very special communities has led to their inclusion in both the EC Habitats Directive and the UK Biodiversity Action Plan as priority habitats for conservation.

Because the term "mire" is such a general one, it is usually split into bogs, fens or flushes, depending on the type of water source involved:

- True **bogs** receive all their nutrient input from rainwater, and tend to have a very nutrient-poor, acid environment, often characterised by abundant *Sphagnum* mosses;
- **Fens** are different, in that they additionally receive groundwater input, either from the surrounding water table, or from springs, seepages and surface water runoff. The nature of a fen depends a lot on the source of the groundwater: base-rich groundwater (e.g. from limestone bedrock) offers more plant nutrients than a true bog, whereas acidic groundwater (e.g. from Manx slate) tends to create fen conditions that are similar to bogs fed by rain alone;
- If a patch of fen results from a localised source of surface water flow (a spring, for example), it is usually classed as a **flush**. Flushes result in the familiar "bright green patches" on hillsides.

The different categories of bog, fen and flush vegetation known to occur on the Isle of Man are detailed in **Types of Mire**, below. For the purpose of these Criteria, bogs, fens and flushes are taken to include all waterlogged habitats, whether seasonal or permanent, *except*:

curragh - which is covered under the Woodland and Scrub criteria;

inundation grassland - which is covered along with other Grassland types; and

swamp - which is associated with aquatic habitats, as opposed to mire which has waterlogged soil but little actual standing water. Swamp is covered under the Freshwater and Swamp criteria.

Note: Bogs and fens are treated separately in the UK *Guidelines for Selection of Biological SSSIs*. However, many of the most distinctive types of fen found in the UK do not occur on the Island, whilst the bogs and fens that do occur are often closely interrelated and can be hard to tell apart. For these reasons bogs, fens and flushes have here been included within the same set of Criteria.

Assessing bog, fen and flush habitats

In general, the same principles of site assessment apply to mires as to grasslands: all of the Priority Site Criteria listed in Part 3 are relevant. As with grasslands, the position of a mire within the surrounding ecological unit can make a big difference to its importance for wildlife, especially where bird and invertebrate species are concerned. Other factors, such as size, level of disturbance, presence of rare species and extent of the associated peat should all be taken into account when assessing the potential merits of designation.

In particular, the question of water source in relation to nutrient input into the site may be crucial to the current and ongoing nature conservation value of a mire. Hydrology and adjacent land use (e.g. traditional low-input farming methods) are therefore of special concern.

Because of the considerable range of mire habitats known to occur on the Island, their detailed descriptions, classification, status and distributions are treated separately here for reasons of clarity and ease of reference. However, as when assessing a grassland, it should be borne in mind that this is an artificial separation. In fact, the majority of mires contain more than one type of vegetation community, with many bog areas grading into fen, blanket mire into valley mire, and so on. Likewise it is not uncommon for some mire communities to occur *only* within a larger mosaic of wet heath or grassland habitats. Therefore, sites which do not “fit” the broad categories are not necessarily poor examples of species-rich communities, and may be as important for wildlife as more “typical” habitats.

National Vegetation Classification (NVC) categories have been listed below in order to help with the identification of individual bog, fen and flush communities, but it should be noted that when the NVC was being compiled even fewer mire habitat samples were taken from the Island than grassland ones. The NVC should therefore be used as a guide in this context rather than a direct key to the value of a given habitat. This is especially true of upland areas, where a closely-related mixture of habitats may occur even within a small area. Because successful protection of upland biodiversity relies on inclusion of a wide range of closely-related habitat features, upland ASSI selection is treated specifically in Chapter 4.7: Upland habitats.

In the case of very rare bog, fen and flush communities, particularly those which support rare and/or protected species, **all** examples will qualify for selection as ASSIs provided that they are of sufficient size and fulfil the minimum criteria. In most cases, however, candidate sites will be selected to represent the best good examples of each mire type and variant. They will be considered to be “good examples” if they not only fulfil all the minimum requirements but also one or more of the Priority Sites Criteria listed in Part 3.

It should always be borne in mind that the distinction between bog, fen and flush vegetation communities can be very subtle, based on small differences in water source, peat depth, aspect, grazing levels, nutrient enrichment, seasonal drying and a host of other factors. To add to the ecological complexity, both bogs and fens often form part of a mosaic of wet, acid grassland and heath habitats, the whole of which supports an ecology which requires all the separate habitat elements in order to thrive. The delicate balance required in order to maintain bog habitats in particular has led to a widely-recognised category of “modified bog”, which encompasses habitats which result from drying out, scrub-encroachment, over-grazing or nutrient enrichment of established bogs. Despite losing some of their characteristic bog species, modified bogs may also be of high value for wildlife. With this in mind, **the assessment process should aim to protect the full range of Manx bog and fen biodiversity by recognising good examples of varied mire/heath/grassland mosaic habitats as well as the best large areas of a single mire type.**

Types of mire

Two main factors govern the classification of mire types: the source of water and nutrients – which governs the distinction between bogs, fens and flushes as described above - and the depth and type of soil medium. Although it is possible for mires to form in waterlogged mineral soils with no peat present, the majority of bogs, fens and flushes on the Isle of Man are associated with peat deposits. These result from a gradual build-up of dead vegetation – *Sphagnum* mosses in particular - wherever conditions are too waterlogged to allow it to decompose away. This build-up of organic matter can result in a body of peat several metres in depth.

Places where peat is known to be accumulating are known as “active mires” or “active bogs”. Any of the types of mire listed below may be either inactive or active regarding the build-up of peat; in practice however this can be difficult to tell. In some cases one site will have some areas of active peat deposition, and others where the peat layer is no longer being added to, perhaps because of heavy selective grazing, or because the mire in that area is starting to dry out. In general, bogs are most likely to be active, whilst fens and flushes may or may not create a build-up of peat.

Examples of mire types on the Isle of Man include basin mires, where peat has developed within a waterlogged depression in the surrounding land; valley mires, where sloping ground is kept waterlogged by springs, seepages and overflows, often at the base of hills or along either side of mountain streams and rivers; and blanket mires, which can cover substantial areas of mountain top and hillside, and receive most of their water from rainfall alone. Blanket mire includes the familiar "blanket bog" that covers wet hill-land throughout the uplands of the British Isles. Whilst convenient for the purposes of ecological survey, the three main types of mire are not always very distinct from each other in the field. For example, blanket mire often grades into slightly more nutrient-rich valley mire on the lower slopes of hills and alongside streams and rivers. This is particularly the case on the Isle of Man, where some habitat categories are actually very closely associated with each other and probably best lumped together for convenience.

Site selection criteria for mire habitats on the Isle of Man have been grouped as follows:

Bog

Mires which receive all or nearly all of their nutrient input from rainwater alone.

Lowland "raised bogs" are very rare; "blanket bog" is the commonest type of mire on the Island and throughout the British Isles. Blanket bog usually occurs on deep peat in the uplands, and is here taken to include the category of "modified bog" .

Valley mires - fens associated with groundwater or surface water flow from springs, river heads, and
flushes - seepages and overflows on hillsides, sloping river and stream valleys and valley floors.

Basin mires

Areas of fen which form where water is "ponded" or trapped in depressions in the surrounding land. Basin mires can cover a large area, but on the Isle of Man they are mostly limited to shallow scrapes and ponds which have filled up with vegetation and now have waterlogged soil but no standing water.

Fen meadows

Waterlogged meadows which are subject to grazing and/or hay cutting. These are often treated as marshy grassland, but they are strictly mires and their vegetation and ecology reflect this. Although rare, they represent one of the most diverse mire habitats on the Isle of Man. Fen meadows are a remnant of a type of farmland which would once have been more common; unlike most mire communities, they are directly dependent on farming methods. This kind of pasture is now thought to be one of the fastest-disappearing wildlife habitats in the British Isles.

These Manx mire types, and the National Vegetation Classification (NVC) categories they most closely resemble, are listed in detail below.

4.4.1 Bog

Description and Distribution

Bogs are mires that support vegetation that is usually peat-forming and which receive mineral nutrients principally from rainwater, with little or no surface flow or groundwater input. This is referred to as ombrotrophic (rain-fed) mire. The vegetation of bogs which have not been modified (e.g. by surface drying and aeration or heavy grazing) is dominated by acid-loving plant species such as bog-mosses *Sphagnum* spp., cotton-grass *Eriophorum* spp. and cross-leaved heath *Erica tetralix*. The water-table on these types of bogs is usually at or just below the surface. The term "bog", like "mire", is a broad category, usually split into two main types, namely raised bog and blanket bog, both of which are capable of a lot of variability. Raised and blanket bogs have broadly similar ecosystems, although some plant and invertebrate species are clearly more specialist than others.

Raised bogs are a rare habitat type both on the Isle of Man and throughout the British Isles. They generally form on flat lowland sites, where the accumulated dead mosses and other organic matter have built up to above the original ground level, forming a characteristic "dome" of waterlogged peat. In the 1994 Ecological Habitat Survey, only one small area of raised bog was identified on the Island. However,

it is possible that small localised patches of raised bog may also occur on more or less flat ground within larger areas of Manx bog or fen, resulting in small pockets of unusual invertebrate or vegetation communities.

Blanket bog is not confined to areas of poor drainage but rather can cloak whole landscapes, even developing on slopes of up to 30°. The period over which blanket peat has been accumulating and the depth it can attain are very variable and not necessarily related. UK studies indicate that most blanket peat development began 5000-6000 years ago, but the range extends from 9000 - 1500 years ago. There is evidence to suggest that some areas of blanket bog began to form following clearance of the original forest cover by early man, but the relative significance of this activity and changing climate on the historical and contemporary extent of the resource has yet to be determined either for the Isle of Man or elsewhere in the British Isles. There is no agreed minimum depth of peat which can support blanket bog vegetation. Throughout the British Isles as a whole, peat depth is very variable, with an average of 0.5-3 m being fairly typical but depths in excess of 5m not unusual. The majority of bogs on the Island are associated with peat deposits of between 0.5 and 3.0 metres.

Throughout their range, blanket bogs support a very wide range of terrestrial and aquatic plants, vertebrates and invertebrates. Some of these are widespread and common, such as heather *Calluna vulgaris*, cross-leaved heath *Erica tetralix*, deer grass *Trichophorum cespitosum*, and cotton grass *Eriophorum* species. Some are much more local, such as the UK BAP species pillwort *Pilularia globulifera*, a protected plant on the Isle of Man. Some are of international interest, for example Eurasian golden plover *Pluvialis apricaria*, which is an infrequent visitor to upland bog and modified bog on the Island, and a rare breeder. One striking feature of very nutrient-poor bogs is the presence of carnivorous plants, which supplement the low nutrients available in the bog with ingenious ways of catching and digesting insects. These species are too specialised to thrive on richer habitats, and depend on mire habitats for their survival.

The presence, extent and type of surface patterning is an important feature of blanket bogs, often exerting a strong influence on the abundance and diversity of plants and animals present. Patterning can range from a relatively smooth surface, with the only irregularities being those created by vegetation features (eg hare's-tail cotton-grass *Eriophorum vaginatum* tussocks and *Sphagnum* moss hummocks) to the extreme patterning associated with suites of bog pools and the intervening ridges. Whilst not as dramatically varied as bogs elsewhere in the British Isles, the full range of patterns of Manx bogs should be represented by the ASSI selection process, in order to reflect the full range of biodiversity involved.

In addition to supporting a diverse ecosystem, blanket bogs fulfil an important role as repositories of archaeological and palaeoecological material and have functional values as agricultural rough grazing, sporting estate and water catchments. In the context of climate change the role of blanket bogs as a carbon store is also now considered significant.

Blanket bog is one of the most extensive semi-natural habitats in the UK and ranges from Devon in the south to Shetland in the north. As with the rest of the British Isles, blanket bog is by far the commonest type of mire on the Isle of Man, covering substantial areas of the uplands; however, despite its regional abundance, blanket bog is very restricted globally, and is one of the habitats listed in the EC Habitats Directive as internationally threatened. For this reason, countries within the British Isles are considered to have a special responsibility for the conservation of these distinctive and ecologically fragile areas.

The bog habitat category also includes **modified bog** vegetation that essentially resembles wet or dry dwarf shrub heath but occurs on deep acid peat which would have once supported peat-forming vegetation. Modified bog includes impoverished vegetation dominated by purple moor-grass *Molinia caerulea* or hare's-tail cotton-grass *Eriophorum vaginatum*. Although there is no agreed minimum depth of peat that can support ombrotrophic vegetation, unmodified bog can be identified floristically by the presence of characteristic species such as cotton-grass *Eriophorum* spp. and peat-forming *Sphagnum* mosses – a feature that tends to be absent or much-reduced once a bog has been modified. Wet modified bog is rare on the Isle of Man, being mostly confined to the Ballaugh Curragh, but dry modified bog is fairly frequent; on the Island it usually takes the form of wet heather or purple moor-grass dominated moorland, modified by farming activities or changes in hydrology, but with some bog species remaining.

The extensive nature of both unmodified and modified blanket bog is such that other mire habitats frequently occur in close association with it. It is therefore very variable and may include localised areas of fen habitat wherever the "blanket" is interrupted by surface water flow, seepage and springlines. Such localised variations in the strict definition of a bog are usually included within the same category as blanket bog to avoid confusion with more clearly differentiated types of fen. In the Manx uplands blanket bog can thus encompass some areas classed as 'intermediate bog' (i.e. sharing features of both raised and blanket bog) together with examples of spring, flush and poor fen, a range of oligotrophic (nutrient poor) water bodies whose catchment is largely or entirely blanket bog, and those relatively small areas of heath and grassland which occur on better drained slopes and by the many streams and rivers which drain areas dominated by blanket bog. Not only are all such areas in hydrological connection with the surrounding peat mass, they frequently contribute to the overall habitat requirements of the peatland fauna. This inextricable mosaic of upland habitats is thought to require a more comprehensive approach to site selection than more clearly-defined lowland habitats, hence the selection of blanket bog ASSIs is covered specifically in Chapter 4.7: Upland habitats.

Status of bogs

The total extent of blanket peat in the UK amounts to just under 1.5 million ha. There is no agreed figure for the extent of blanket bog vegetation. In terms of national cover of blanket peat soil (in the main >0.5 m deep) England supports some 215,000 ha, Scotland approximately 1,060,000 ha, and Wales has around 70,000 ha. Northern Ireland has approximately 140,000 ha of blanket bog vegetation. Significant proportions of peat soil, probably in excess of 10%, no longer support blanket bog vegetation (for example because it has been modified by agricultural activities, or subjected to commercial peat extraction).

Comprehensive data for changes to the total UK resource are lacking, but studies in Scotland (where most of the resource lies and where it accounts for some 13% of the land area) suggest a 21% reduction in the extent of blanket mire between the 1940s and the 1980s. The greatest single cause of this reduction (51%) is afforestation, and substantial losses to forestry are reported from Wales. Further losses of extent and condition can be attributed to drainage and heavy grazing, peat cutting and atmospheric pollution, resulting in significant habitat change in, for example, mid and south Wales and the Pennines.

Within the Isle of Man, bogs are rarely dug out for peat, but may lose some of their diversity due to farming activities or, occasionally, development such as pipeline construction. The 1994 Ecological Habitat Survey report identified a total of just 0.36ha of true raised bog, 105.72ha of blanket bog, 26.08ha of wet modified bog, and 67.36ha of dry modified bog. It is not known how much previously-unmodified blanket bog has since been modified, but the process is plainly still under way on the Island due to the need to utilise farmland to the best ability. The introduction in 2002 of the Manx Agri-environment Scheme may contribute to reducing the rate of modification of otherwise undisturbed upland bogs.

Classification of bogs

Bearing in mind the comments above regarding the use of the National Vegetation Classification (NVC), bogs on the Island may be classified broadly as outlined in **Table 4.4a**:

Table 4.4a: Broad categories of bog on the Isle of Man

Type and Description	Species rich?	Closest NVC categories	Rarities present?	Protected species present?
Bog pools	yes	M1 <i>Sphagnum auriculatum</i> bog pool community. No Island NVC sample.	Highly likely to support rare invertebrates	Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing.
Cotton-grass bog	yes	M17 <i>Scirpus cespitosus</i> - <i>Eriophorum vaginatum</i> blanket mire One Island NVC sample M19 <i>Calluna vulgaris</i> - <i>Eriophorum vaginatum</i> blanket mire (restricted to highest upland areas) No Island NVC sample	May support rare plants and/or invertebrates	Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing.
Cross-leaved heath bog (sometimes modified bog)	yes	M18 <i>Erica tetralix</i> - <i>Sphagnum papillosum</i> raised and blanket mire No Island NVC sample	May support rare invertebrates and/or plants	Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing. A likely habitat for cranberry <i>Vaccinium oxycoccus</i> and rare sedges such as white sedge <i>Carex curta</i> .

Minimum selection criteria for bogs

Raised bog

Any unmodified examples of raised bog will qualify for selection if they are over 0.25ha in total extent; Examples of smaller extent will be considered if they form a significant element of a larger area of related habitats that fits at least one of the priority sites criteria as listed in **Part 3**.

Blanket bog

Blanket bog is confined to the uplands; selection of blanket bog will therefore be covered entirely under the specific upland site selection process in Chapter 4.7.

Wet modified bog

Examples of wet modified bog over 1ha in extent will qualify for selection; Examples of a smaller extent will be considered as a part of a larger mosaic of related habitats, and/or where they fulfil at least one of the Priority Sites Criteria listed in Part 3.

Dry modified bog

Dry modified bog will usually be selected only as part of a larger mosaic of related habitats; this may include heathland and upland sites (See Chapters 4.3 and 4.7 respectively).

4.4.2 Valley mires and flushes

Description and Distribution

A distinctive, complex habitat on the Isle of Man, often associated with blanket bog areas, but also frequently grading into swamp and open water, grassland or heath. Valley mire is naturally marginal to other habitats, but it can also represent the only remaining semi-natural vegetation within improved farmland, especially in or adjacent to upland areas where all the surrounding habitats have been modified or improved. Flushes may occupy quite large areas, particularly at the head of rivers, but usually they are small, appearing as characteristic "bright green patches" on hillsides and valleys. Even small flushes can house unusual or very local invertebrate and plant populations. Valley mires and flushes on the Isle of Man are usually on peat soils, with a neutral to acid chemistry, and hence often have similar species to bogs.

Classification

Bearing in mind the comments above regarding the use of the National Vegetation Classification (NVC), valley mires and flushes on the Island may be classified broadly as outlined in **Table 4.4b**:

Table 4.4b: Broad categories of valley mires and flushes on the Isle of Man

Type and Description	Species rich?	Closest NVC categories	Rarities present?	Protected species present?
Sedge-dominated mires	Not always botanically rich	M4 <i>Carex rostrata</i> - <i>Sphagnum recurvum</i> mire No Island NVC sample M6 <i>Carex echinata</i> - <i>Sphagnum recurvum/auriculatum</i> mire One Island NVC sample	May support rare plants and/or invertebrates	Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing. A likely habitat for rare sedges such as white sedge <i>Carex curta</i> .
Cross-leaved heath wet heathland (also, see Chapter 4.3 Lowland heathland)	yes	M16 <i>Erica tetralix</i> - <i>Sphagnum compactum</i> wet heath No Island NVC sample	May support rare invertebrates and/or plants	Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing. A likely habitat for cranberry <i>Vaccinium oxycoccus</i> and rare sedges such as white sedge <i>Carex curta</i> .
Purple moor-grass pastures	sometimes	M25 <i>Molinia caerulea</i> - <i>Potentilla erecta</i> mire 1 Island NVC sample	May support rare invertebrates and/or plants	Protected plants such as orchids are highly likely to be present.
Bog pondweed flushes	Not always	M29 <i>Hypericum elodes</i> - <i>Potamogeton polygonifolius</i> soakaway	May support rare invertebrates and/or plants	A possible habitat for Pillwort <i>Pilularia globulifera</i> and other unusual plants.

Valley mire and flush status

In the UK the majority of fens as a whole are notified as SSSI/ASSIs and many are notified as Wetlands of International Importance under the Ramsar Convention and as SPAs under the EC Birds Directive. Several of the larger fens are managed as NNRs by EN and CCW, and as reserves of the RSPB and County Wildlife Trusts. Several types of fen are listed in the Habitats Directive including transition mire, poor and rich fen, alkaline fens (rich-fen not found on the Isle of Man). A number of fens have been proposed as SACs under the EC Habitats Directive for these types.

Fens as a whole are now a Priority Habitat in the UK Biodiversity Action Plan; small flushes and patches of valley mire which fall within active upland blanket bog areas would also be covered under the Priority Habitat: Blanket Bog. Within the Isle of Man true valley mire is very scattered and often limited in extent, totalling only 14.56ha in the 1994 Ecological Habitat Survey Report. Flushes, which include some extensive areas eg. at the head of the Blaber River, amount to 455.24ha, although it should be noted that much of this occurs on one or two large sites; the rest is very scattered.

Minimum selection criteria for valley mires and flushes

- **All** examples of valley mires and flushes greater than 2ha will qualify for selection;
- Sites which have a mosaic of habitats in which valley mires and/or flushes feature as a proportion of the overall vegetation will qualify for selection if their total extent is greater than 2ha and they fit at least one of the Priority Sites Criteria as listed in Part 3.
- Flushes which occur in association with saltmarsh, cliff or other coastal habitats will be covered by the relevant coastal habitat criteria.
- The importance of valley mires that occur within upland areas has been taken into account in the upland site selection process. Valley mires and flushes that lie within upland areas will therefore be selected by the specific upland site selection process detailed in Chapter 4.7: Upland habitats.

4.4.3 Basin mires

Description and Distribution

Basin mires are a form of fen which develops when water is “ponded up” within a depression in the ground. This may be a natural topographical feature, such as a kettle hole or pingo, or a man-made depression such as an ancient quarry working or abandoned peat-digging. In either case, the vegetation depends on the water chemistry involved. Manx groundwater is generally acidic in nature, hence basin mire vegetation is likely to be similar to bog and acid flush vegetation in nature.

Basin mires are not usually large in extent; the 1994 Ecological Habitat Survey Report noted that several basin mires had been seen during survey work around the Island, but all of them had been too small to map. In spite of this, basin mires may support diverse vegetation and unusual or local invertebrate species, often extending the biodiversity of a larger bog, fen, heath or marshy grassland site.

Classification

Basin mires generally fall within similar NVC categories to valley mires, flushes and bog pools; see Tables 4.4a and 4.4b.

Basin mire status

Basin mires of more than a few hectares in extent are not particularly common anywhere in the British Isles, and calcareous basin mires (not found on the Island) are especially rare. Within the Isle of Man they occur in both the uplands and lowlands, but mostly the uplands, within larger areas of wet semi-natural habitat. Basin mires with good examples of fen habitat are covered under the UK Biodiversity Action Plan Priority Habitat: Fens. Smaller basin mires within upland active blanket bog areas are covered under the UK BAP Priority Habitat: Blanket Bogs.

Minimum selection criteria for basin mires

- Basin mires occurring in upland areas are covered under the Upland site selection process detailed in Chapter 4.7;
- Other basin mires will generally qualify for ASSI selection only where they form part of a larger mosaic of habitats which fulfils at least one of the Priority Sites Criteria listed in Section 3.2. However, exceptional examples of basin mire, e.g. those which fulfil one or more of the Priority Sites Criteria in their own right, will be considered on their own merit, with priority given to those which support very rare or local plant or invertebrate communities.

4.4.4 Fen meadows

Description and Distribution

Unlike most mire habitats, fen meadows are usually part of enclosed farmland, and have been produced by management (grazing, mowing and/or burning). Cessation of management usually results in taller herb and woody species invading and taking over, usually following a gradual succession to curragh. Fen meadows are usually dominated by purple moor-grass *Molinia caerulea*, rushes *Juncus spp.* and sedges *Carex spp.*, and include the category now generally known as “Rhos pasture”, or Purple moor-grass pasture, which is often very diverse and which has suffered a severe rate of loss throughout the British Isles in recent years.

Manx fen meadows have a neutral to acid peat soil, and often contain a wider range of plants than the bogs and valley mires which they otherwise resemble. True flood-plain fens such as those typical of lowland England do not occur on the Island, but some river valleys are wide enough to encompass narrow fen meadows – enclosed valley mire - which may be at least partly dependent on flooding for their water source.

The main types of fen meadow that occur on the Island are **purple moor-grass and rush pastures**. Purple moor grass and rush pastures occur on poorly drained, usually acidic soils in lowland areas of high rainfall in western Europe. In the UK, they are found in south-west England, particularly in Devon, southern Wales, south-west Scotland, perhaps extending as far north as northern Argyll, and in Northern Ireland, especially Fermanagh. Elsewhere in Europe they are particularly characteristic of the oceanic and sub-oceanic regions of the western seaboard, from Portugal to the Low Countries, extending eastward into central Europe. Their vegetation, which has a distinct character, consists of various species-rich types of fen meadow and rush pasture. Purple moor-grass *Molinia caerulea*, and rushes, especially sharp-flowered rush *Juncus acutiflorus*, are usually abundant. Just as the best examples of lowland heath contain a wide range of plant communities, so the same is true for this habitat: the characteristic plant communities often occur in a mosaic with one another, together with patches of wet heath, dry grassland, swamp and scrub.

Key species associated with purple moor grass and rush pastures include: whorled caraway *Carum verticillatum*, marsh hawk's beard *Crepis paludosa*, greater butterfly orchid *Platanthera chlorantha*, curlew *Numenius arquata*, snipe *Gallinago gallinago*, and barn owl *Tyto alba*.

Classification

Fen meadows on the Island may be classified broadly as outlined in **Table 4.4c**:

Table 4.4c: Broad categories of fen meadow on the Isle of Man

Type and Description	Species rich?	Closest NVC categories	Rarities present?	Protected species present?
Rush pastures	sometimes	M23 <i>Juncus effusus</i> / <i>acutiflorus</i> – <i>Galium palustre</i> rush pastures 1 Island NVC sample. Possibly also M27c <i>Filipendula ulmaria</i> - <i>Angelica sylvestris</i> mire, <i>Juncus effusus</i> - <i>Holcus lanatus</i> sub- community No Island NVC sample	May support rare invertebrates and/or plants	Protected plants such as orchids may be present.
Purple moor-grass pastures	sometimes	M25 <i>Molinia caerulea</i> - <i>Potentilla erecta</i> mire 1 Island NVC sample	May support rare invertebrates and/or plants	Protected plants such as orchids are highly likely to be present.

Fen meadow status

Fen meadows are one of the few mire habitats which can profitably be improved for agricultural purposes, usually through drainage, liming and fertilizing. In times of increasing pressure for farmers, this has led to a sharp reduction in the overall area of fen meadow in the British Isles, as previously low-input fields are brought into more intensive use. This is true of the Isle of Man as elsewhere. Purple moor grass and rush pastures – the main types of fen meadows on the Island - are a special priority for nature conservation because they are highly susceptible to agricultural modification and reclamation throughout their range. In Devon and Cornwall, where the habitat is known as Culm Grassland, only 8% of that present in 1900 remains, with a staggering 62% of sites and 48% of the total area being lost between 1984 and 1991. In Northern Ireland, between 1990 and 1993, the rate of loss of fen meadow was reckoned to be 3.3% per annum. Fragmentation and isolation of stands have been common.

The total area of this habitat in the Isle of Man is hard to ascertain, as much of it is likely to have been included in the Marshy grassland Phase 1 habitat category. However, as this whole category amounts to only 1096ha or 1.94% of the Island, some of which may already have been modified or destroyed, the total area of fen meadow is not likely to be great.

Because of the high rate of loss, purple moor-grass and rush pastures are now a Priority Habitat in the UK Biodiversity Action Plan, and several key sites have been designated SSSIs and/or SACs in recent years. In Wales it is estimated that there is now about 24,000 ha of lowland purple moor-grass and rush pasture. In south west England 530 purple moor grass and rush pastures sites are known to survive on the Culm Measures, covering 3,981 ha, 400 sites on Dartmoor covering 1,000 ha with a further 90 sites covering about 300 ha on the Blackdowns. In Northern Ireland it was estimated that there was about 24,600 ha in 1993. No area estimates are available for Scotland, but the total extent is thought likely to be in the region of 2,000 ha. Thus it is probable that the total extent of the habitat in the UK is now about 56,000 ha. This is thought to be considerably more than survives in the rest of Europe, with the possible exception of the Republic of Ireland. Countries in the British Isles therefore have a responsibility for the conservation of this habitat, which is in danger of ecological extinction if the current rate of loss continues.

Minimum selection criteria for fen meadows

- **All** good examples of fen meadows over 1ha in extent will qualify for selection;
- Smaller areas of fen meadow may qualify for selection if they fulfil at least one of the Priority Sites Criteria listed in Part 3, with priority given to sites which form part of a larger mosaic of interrelated habitats.
- Fen meadows may also fall within upland areas; these will be covered under the upland site selection process detailed in Chapter 4.7.



Wet pasture at Glen Rushen ASSI; the furry appearance is due to the presence of whorled caraway *Carum verticillatum* leaves.

4.5 Open water and swamp

Description

The size of the Isle of Man and the absence of extensive chalk or limestone areas naturally limits the range of open water bodies present compared with adjacent countries. For example, there are no large lowland river systems, lakes are few and mostly limited to man-made reservoirs, and marl water bodies and chalk streams are absent altogether. Instead, the most typical watercourses are small, soft-water streams in the uplands, which often reach the sea via dramatic waterfalls and deep coastal glens.

Flooded quarry workings and other industry-related man-made water bodies are rare on the Island. Standing water is most often found in reservoirs, or in dubs (shallow farm ponds) on farmland, most of which are concentrated in the north of the Island (Allen, 1984). Another farmland feature which forms a significant habitat in parts of the Island is the presence of drainage ditch systems, some of which have a well-established aquatic flora and fauna resulting from decades of continuous management. Whilst not natural watercourses, some ditch systems do connect with rivers, and most are designed to have a slight flow in order to carry excess water away. The ecology of well-established drainage ditches is therefore intermediate between that of ponds and streams (although species of fast-flowing upland streams are likely to be absent).

Despite the high proportion of coastline, there are few examples of salt or brackish standing water bodies on the Island, due to the lack of suitable topography. Those that there are – such as the seasonally-wet dune slacks on the Ayres National Nature Reserve – tend to support rare and local plants and invertebrates, and to have a high level of importance for coastal wading birds. Such brackish pools will be covered in Chapter 4.6: Coastal habitats, along with strictly tidal areas such as river estuary saltmarsh.

A distinctive ecological feature of all open water on the Island is the tendency for water levels to fluctuate, often sharply, following changes in the weather. For example, small ponds and dubs may dry out in summer, but remain sufficiently flooded for the rest of the year to enable aquatic plants and animals to survive from year to year (see Samson, 2002). Their ecology is adapted to this situation. Likewise, some upland streams with small, steep catchment areas may dry to a trickle in warm summer months, but fill up to a powerful torrent after one night of heavy rain. This has necessitated flood protection measures in many parts of the Island. For these reasons, both permanent and seasonal open water features are covered by the Criteria, and those watercourses which retain the most natural structure will be considered a priority for protection.

Aquatic habitats are one type of site for which buffer zones may be required in order to carry out effective species conservation, even where the biodiversity of the buffer zone itself is low. This is because of the disproportionate vulnerability of wetland habitats to runoff, spray drift and other “edge effects”, which can quickly damage the entire ecology of a site once they affect the water. Buffer zones can also protect standing water from bank erosion by grazing livestock, and may be important habitats in their own right, e.g. rank grassland and scrub that offer cover for frogs and a feeding area for birds and semi-aquatic invertebrates. No cultivation, spraying or fertilizer applications should be carried out in a buffer strip. It may also be necessary to exclude livestock, at least for certain periods, in order to protect the banks and water quality.

This chapter covers all freshwater features, including submerged, floating and emergent (swamp) vegetation communities. Rivers and streams are both covered here, but it should be noted that in the uplands, most watercourses will be selected as part of an overall upland area, using the Upland habitats criteria in Chapter 4.7.

Assessing open water and swamp

All of the Priority Sites Criteria listed in Part 3 of this document apply to the selection of open water and swamp sites. However, the special nature of freshwater ecology – its motability and its susceptibility to very quick changes in chemistry, clarity and flow rates – means that the following criteria should be given special consideration when assessing an open water site:

Naturalness, representativeness and water quality

Water bodies are one category of habitat where **man-made sites** may offer good substitutes for natural pools and lakes. This is particularly true of sites where the water quality is good (see below) and has been allowed to develop over a long period of time – e.g. old, flooded quarries, which may in exceptional cases support submerged, floating or swamp vegetation communities very similar to those found on natural lakes (NCC 1989, *Ch6*). In view of this, although naturalness is still a very desirable feature, both semi-natural and man-made sites will be considered for selection.

Whether subject to human influence or not, the **flow rate** of watercourses is an important factor when assessing the naturalness and representativeness of a site, because the ecology of the watercourse will be adapted to the prevailing speed of water flow, in particular the available oxygen levels and the need to secure a stronghold. The level of human modification – especially flood relief measures and other control structures – is therefore an important consideration when assessing the naturalness and representativeness of rivers and streams. Because most water bodies on the Island are not large, other questions of water movement, such as wind and wave effects on lake shoreline vegetation, will not normally apply, although reservoirs with very hard, wave-swept edges are clearly less well-suited to the formation of emergent vegetation fringes than those with softer margins.

Position within a larger mosaic of habitats can indicate the naturalness of an open water site – see *Position in ecological unit*, below.

Although different shapes and sizes of water body may look very distinctive, from an ecological perspective the **chemical quality** of the water itself is often the more important factor, because this tends to determine the range of animals and plants which can live there; it may also indicate the degree of human modification currently affecting the site. The need to assess the water quality of a site using chemical analysis varies according to circumstances; in most cases the classification of the habitat will be obvious from the diversity of species present. Water analysis is more helpful in assessing the condition of an existing open water habitat, particularly where there appears to be a problem, for example with escalating algal growth or sudden fish deaths. The results of water quality testing can help with comparing similar watercourses for selection.

Where chemical analysis is to be carried out, the first consideration is usually trophic status – i.e. how nutrient-enriched the water is. Water with plenty of available plant nutrients (particularly nitrogen and phosphorus) is classed as eutrophic, whilst a mesotrophic water body has only moderate nutrient levels, and oligotrophic water is relatively nutrient-poor. Water which contains a high level of dissolved substances such as tannins, but little available plant nutrients, is referred to as dystrophic, and is usually associated with peatland areas where the water is acidic and dark-stained. All these categories are naturally associated with a particular ecology, and all may contain rarities in certain cases; a change in trophic status can therefore result in considerable damage to an open water ecosystem, whilst a stable level of nutrient input is usually associated with a more natural and representative site.

The **water quality** of a river, pond or lake can be affected by a variety of factors, both man-induced and naturally occurring. Some of the most important considerations are:

Biological oxygen demand (BOD) – a high BOD is a sign that water is highly nutrient-enriched, leading to excessive algal and/or bacterial activity and a corresponding increase in the demand for oxygen. The effects on organisms which need high levels of dissolved oxygen can be severe, with extreme cases – such as those caused by sudden milk or silage spills – leading to mass fish kills (DEFRA, 2000). An unhealthily high BOD may be due to an ongoing situation, or may relate to a single event. In either case the biodiversity of the water body is likely to be adversely affected.

pH (acidity/alkalinity)

The pH of a water body has a direct effect on important aspects of water chemistry such as the balance of calcium in the water. A higher pH is usually taken to indicate a higher trophic status, i.e. eutrophic, rather than mesotrophic or oligotrophic water. With the exception of sites in the vicinity of Castletown, where limestone is the underlying rock, most Manx water is relatively acidic, soft water (with low levels of dissolved calcium and magnesium salts). The ecology of Manx rivers and streams reflects this and is adapted to soft water conditions. In the case of bog pools, a steady, very acidic pH of 4.5 or less is vital

in order to support the few, specially-adapted species that can survive in the extreme conditions. An increase in alkalinity, for example due to run-off from nearby building works, can threaten the special interest of such sites, not least because increased pH is often associated with increased nutrient levels. However, it should be noted that in the case of lakes and ponds which naturally have a lot of plant and algal growth in summer, the water can vary considerably in pH – becoming more alkaline when vegetation growth is at its strongest.

Plant nutrients

The plant nutrients most often tested for are ammonia, nitrates and phosphates, although a range of other chemical tests – eg. for potassium levels and sulphates – may also be useful in assessing the condition of a water body. Healthy eutrophic ponds, lakes and rivers have relatively high levels of plant nutrients and support a high diversity of plants and animals as a result; however, artificial sources of nutrients such as phosphates and ammonia can far exceed healthy levels. Ammonia, nitrates and nitrites are all potentially dangerous to fish at high levels, and are associated with a high BOD. Phosphates are of particular concern because high phosphate levels result in greatly accelerated algal growth, creating a soaring oxygen demand and decreased water clarity and light penetration. This situation can be difficult to remedy.

Conductivity

The electrical conductivity of soft water such as that naturally found in Manx rivers and streams is naturally very low. Inputs such as farm or domestic sewage can hugely increase the conductivity by introducing a high concentration of sodium and other ions. Whilst a good indication that something is amiss with otherwise clean, soft-water streams, conductivity is a less useful test in salt or brackish water, as these water-bodies already have a high conductivity due to the natural salt load of the water.

Undesirable readings on any of the above tests will be considered **negative** points when considering a site for selection, except where the problem is plainly short-term and/or a satisfactory solution is possible. The same applies to more obvious sources of water quality problem, such as oil pollution, mine water contamination or discarded agrochemical containers.

Diversity

In general, well-lit, nutrient-rich, base-rich eutrophic waters are capable of sustaining the greatest diversity of aquatic life, whereas only a very limited range of species can survive in very acidic or dystrophic pools. However, bog pools and dystrophic standing water may support vulnerable rarities such as unusual dragonflies, so the desirability of species richness at any given site needs to be tempered with consideration of how rare both the habitat and species are.

Many animals which inhabit open water require a mixture of feeding areas and cover, and semi-aquatic species are adapted to using more than one habitat in order to complete their life-cycle. Hence a range of adjacent vegetation types is a very desirable feature, e.g. proximity to related habitats such as fen meadows and mires.

Position in ecological unit

The majority of natural watercourses exist within a wider ecological context of upland or lowland wetland situations, hence position within an ecological unit should be one of the prime considerations when assessing two otherwise similar watercourses. Likewise, the biodiversity of ponds and dubs and particularly bog pools may depend to a great extent on the nature and quality of the surrounding habitat. This affects not just the species present (see Diversity, above) but also potential water quality issues such as whether or not the water is shaded, whether grazing stock have access, and whether the nutrient status is being affected by fertilizer runoff.

Open water and swamp types

For the purpose of these Criteria, areas of open water and swamp on the Isle of Man have been grouped broadly as follows:

Still water and swamp: Natural pools, ponds, lakes and seasonally-flooded areas, man-made ponds and dubs, reservoirs and other man-made water-bodies. These are the commonest sites for **swamp** vegetation, which usually occurs around the shallower margins of still water. Swamp may also occur as a habitat on its own, with no open water visible. This is especially true of shallow water bodies, which can fill up with extensive stands of pure swamp such as bottle sedge *Carex rostrata* or common reed *Phragmites australis*. **Brackish pools** and **dune slacks** also occur on the Island; these are covered in Chapter **4.6: coastal habitats**.

Ditch systems: These consist of linked, man-made ditches, occasionally incorporating minor natural watercourses. On the Isle of Man, ditch systems usually result from a desire to drain farmland (Allen, 1984), either on an ongoing or seasonal basis. They may be strategically placed or, more commonly, randomly associated within a particular area. Well-established ditch systems may support a good range of aquatic plant and invertebrates, and may be particularly important for semi-aquatic species.

Flowing water: Rivers and streams, both lowland and upland. Some swamp may form alongside the margins of slower-flowing rivers, but it is usually restricted to areas where there is a "bay" or widening out of the main watercourse. Note: river estuary saltmarsh is covered in Chapter **4.6: Coastal habitats**.

These Manx open water and swamp types, and the National Vegetation Classification (NVC) categories they most closely resemble, are each treated separately in detail below.

4.5.1 Still water and swamp

Description and Distribution

The main types of still water bodies on the Island are:

Bog pools, which occur in both upland and lowland habitats. Those which occur in the uplands are covered by the Upland Habitats criteria described in **Chapter 4.7**. Those which occur in lowlands are usually confined to the wetter parts of curragh and fen sites. They are amongst the smallest open water bodies on the Island, usually occurring as a collection of small pools and troughs within a larger site.

Bog pools may be too dark and dystrophic to support any vegetation other than *Sphagnum* mosses; however, lowland examples frequently support a mat of floating vegetation (usually dominated by bogbean *Menyanthes trifoliata*) and a fringe of emergent sedges and rushes. Bog pools – including very dystrophic examples – are known to support a small but significant range of rare and local invertebrate species.

Ponds and dubs (shallow field ponds, some of which may dry out in summer) are scattered throughout the Island's farmland, with the majority concentrated in the north. Where ponds and dubs occur within a large semi-natural area – and particularly if several occur in close proximity – their overall biodiversity and value for wildlife is likely to be greatest (q.v. Samson, 2002). Collections of ponds are also more likely to contain rare plants and invertebrates than isolated sites. The beetle fauna has received some study and the resulting species lists can be used to compare site quality. Unlike bog pools, ponds and dubs often occur in the absence of associated semi-natural habitat, surrounded by improved farmland. This can limit their value for wildlife, but even completely isolated examples may still support a good range of aquatic and emergent species. Shallow dubs within improved farmland may also have a special role to play in supporting farmland birds such as lapwing, curlew and snipe.

Lakes and reservoirs are rare on the Isle of Man, and most are the result of human activities. Natural lakes, such as Lough Cranstal (a relic of an ancient, much larger area of open water in the north of the

Island), are also affected by human activities due to the proximity of improved farmland and managed drainage systems. Despite the scarcity of large semi-natural water bodies, the Island's few larger open water sites still support a range of associated vegetation, along with significant populations of both breeding and visiting waterfowl.

Most ponds, dubs and larger water bodies are likely to fall within the broad category of eutrophic standing water, i.e. water bodies with high levels of plant nutrients such as nitrates, calcium salts and phosphates. However, some are more mesotrophic in nature, with species characteristic of less nutrient-rich waters.

Classification of still water and swamp habitats

Still water and swamp habitats on the Island may be classified broadly as outlined in Table 4.5.a



A flowing stream and pools at Glen Rushen ASSI, with whorled caraway *Carum verticillatum* and bog pondweed *Potamogeton polygonifolius*.

Table 4.5.a: Broad categories of still water and swamp habitats on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (after Rodwell, 1992 ⁴)	Rarities present?	Protected species present?
Bog pools These are rare, scattered and tend to be very limited in extent; typical sizes range from 0.01ha to 0.1ha. Most are confined to upland bog areas and will be covered by the criteria for Upland habitats; some occur in the lowlands, where they represent a rare and vulnerable habitat.	Not usually; it is not in the nature of very acid, dystrophic water bodies to support a wide diversity of species.	A14 – <i>Myriophyllum alterniflorum</i> (alternate water-milfoil) community A24 – <i>Juncus bulbosus</i> (bulbous rush) aquatic community (no NVC samples mapped as ubiquitous throughout north west Britain)	This community may occasionally contain unusual sedges, mosses or liverworts; may also support rare dragonflies and other specialised invertebrates.	Sometimes. White sedge <i>Carex curta</i> is known sometimes to occur on the edges of bog pools; Pillwort <i>Pilularia globulifera</i> also prefers shallow acidic waters.
Ponds, dubs and lakes Most ponds, dubs and larger water bodies are likely to fall within the broad category of eutrophic standing water, i.e. water bodies with high levels of plant nutrients such as nitrates, calcium salts and phosphates. However, some are more mesotrophic in nature, with low levels of calcium salts and species characteristic of more acid, less nutrient-rich waters.	Variable, according to degree of nutrient enrichment and modification. May be very species rich in good examples.	A2 – <i>Lemna minor</i> (common duckweed) community A7 <i>Nymphaea alba</i> (native white water-lily) community A8 – <i>Nuphar lutea</i> (yellow water-lily) community A9 – <i>Potamogeton natans</i> (floating pondweed) community A10 – <i>Polygonum amphibium</i> (amphibious bistort, now named <i>Persicaria amphibia</i>) community A11 – <i>Potamogeton pectinatus</i> - <i>Myriophyllum spicatum</i> (fennel pondweed and spiked water-milfoil) community A12 – <i>Potamogeton pectinatus</i> (fennel pondweed) aquatic community A15 <i>Elodea canadensis</i> (Canadian waterweed) community A16 <i>Callitriche stagnalis</i> (common water-starwort) community A19 <i>Ranunculus aquatilis</i> (common water crowfoot) community (all these communities are fairly widespread throughout the British Isles, hence NVC samples were not mapped)	Often. Although not always species-rich, this is a relatively scarce habitat on the Island, and may contain unusual stoneworts, pondweeds and invertebrates.	Fennel pondweed <i>Potamogeton pectinatus</i> is itself a protected plant on the Island.

<p>Bottle sedge swamp</p> <p>Usually found in association with bogbean (e.g. in the Curragh) and water horsetail (e.g. around the edges of larger water bodies). This kind of swamp is typical of nutrient-poor, acid waters on peat soils, and may grade into bog pools.</p>	<p>Not very; although this is one of the more diverse acidic swamp vegetation communities, the habitat is not usually capable of supporting a high degree of species richness.</p>	<p>S9 – <i>Carex rostrata</i> (bottle sedge) swamp, especially the <i>Menyanthes trifoliata</i>-<i>Equisetum fluviatile</i> sub-community. No Island sample in NVC maps.</p> <p>S10 – <i>Equisetum fluviatile</i> (water horsetail) swamp, <i>Carex rostrata</i> sub-community also see below. No Island NVC samples.</p>	<p>Often forms a good habitat for unusual aquatic invertebrates; some rare plants may also occur.</p>	
<p>Water horsetail swamp</p> <p>May occur as dense stands of horsetail (e.g. at the edge of lakes) or smaller mats of mixed vegetation with similar species to bottle sedge swamp.</p>	<p>Not usually.</p>	<p>S10 <i>Equisetum fluviatile</i> (water horsetail) swamp No Island NVC sample.</p>		
<p>Tussock-sedge swamp</p> <p>A distinctive mix of large sedge tussocks and emergent vegetation in between; can provide a unique habitat for invertebrates. Very rare on the Island.</p>	<p>Moderately species-rich</p>	<p>S3 <i>Carex paniculata</i> (greater tussock-sedge) sedge swamp No Island NVC sample</p>	<p>May support rare invertebrates and unusual emergent plant species</p>	
<p>Spike-rush swamp</p> <p>Typically forms a fringing zone around larger lakes and reservoirs.</p>	<p>No.</p>	<p>S19 <i>Eleocharis palustris</i> (common spike-rush) swamp</p>	<p>May form part of a larger habitat for unusual invertebrates.</p>	
<p>Reedmace/bulrush swamp</p> <p>This species is characteristic of broad pond and river margins, where it forms dense stands (although rarely on the scale of bulrush swamp in the English lowlands). In the absence of reedbeds this forms a valuable bird habitat.</p>	<p>Variable; may be a single-species only stand of bulrush</p>	<p>S12 <i>Typha latifolia</i> (greater reedmace) swamp</p>	<p>May support unusual invertebrates and birds.</p>	<p>May attract protected bird species including rare visiting waterfowl.</p>

<p>Common club-rush swamp</p> <p>A tall and distinctive vegetation forming a sheltered water margin habitat; may have similar associated plants to bottle sedge swamp (see above). May occur in bays and sheltered stretches of river margins. A rare habitat on the Island.</p>	<p>Often species-poor, but may contain a range of submerged and floating species amongst the rushes.</p>	<p>S8 - <i>Scirpus lacustris</i> ssp. <i>lacustris</i> (common club-rush, now known as <i>Schoenoplectus lacustris</i>) swamp One Island NVC sample.</p>	<p>A number of unusual plants may be present, as this is a rare type of vegetation on the Isle of Man. Unusual invertebrates may also be present in this habitat for at least part of their life-cycle.</p>	<p>Yellow water-lily, white sedge and common bladderwort may all occur within this vegetation type, but very rarely do so on the Island; Common club-rush is itself a protected species. May form part of the favoured habitat for shy protected water birds such as the water rail and bittern.</p>
<p>Greater pond-sedge swamp</p> <p>Typical of lowland English lake and canal margins; at the north end of its usual range here, with only one occurrence (Loch Cranstal)</p>		<p>S6 - <i>Carex riparia</i> (greater pond-sedge) swamp No Island NVC sample</p>		
<p>Bur-reed swamp</p> <p>This distinctive plant is at the northern edge of its main range here. It rarely forms large stands, tending to occur in zones along the edge of slower-moving rivers and ditches, and around the edges of sheltered ponds.</p>	<p>Some stands may be moderately diverse due to the presence of emergent species amongst the bur-reed</p>	<p>S14 <i>Sparganium erectum</i> (branched bur-reed) swamp No Island NVC samples.</p>		
<p>Reed swamp</p> <p>Rare on the Island, and never forming the huge areas of "monoculture" habitat found in England. May occur in brackish habitats, e.g. alongside muddy tidal reaches of rivers.</p>		<p>S4 - <i>Phragmites australis</i> (common reed) swamp One Island NVC sample.</p>		<p>May attract protected bird species including rare visiting waterfowl.</p>
<p>Sea-clubrush swamp</p> <p>Typical of more or less still, brackish or salt water alongside estuaries, lagoons and the deeper areas of saltmarshes.</p>	<p>This is very much a halophyte (salt-loving) community, and is therefore covered in Chapter 4.6 - Coastal Habitats, in the section on Saltmarsh.</p>			

Status of still water

Naturally-occurring water bodies with unmodified fauna and vegetation are rare throughout the British Isles, chiefly because of their strong associations with human activities – notably farming, fishing and water supply. The more natural and unspoilt water bodies have therefore received the highest levels of recognition, with regionally restricted features (e.g. turloughs and Mediterranean temporary ponds) included as Priority habitats in Annex 1 of the EC Habitats Directive. A range of oligotrophic lakes, natural dystrophic lakes and ponds, and natural eutrophic lakes – all of which are vulnerable to loss of biodiversity due to pollution and nutrient enrichment - are also listed as Annex 1 habitats. The UK Biodiversity Action Plan recognises both eutrophic and mesotrophic lakes as Priority Habitats, along with reedbeds.

The DAFF Phase 1 Habitat Survey recorded an average size of water body on the Island of just 0.273ha, with a maximum of 25.4ha and a total area of 105ha (which indicates the rarity of large bodies of open water) and just 19ha of true swamp (Sayle *et al*, 1995). Whilst there are over three hundred standing water bodies recorded, many are much smaller than the mean average size, often as small as 0.003ha; many more were too small to be included in the survey. Such ponds and dubs are rarely found in association with other wetland habitats and tend to be vulnerable to drying, pollution, stock poaching and other ecological “edge effects”. Many therefore have a limited biodiversity; for those that are species-rich, a **buffer zone** is usually vital for effective conservation of both the aquatic and emergent species.

Whilst the majority of distinctive British open water types do not occur on the Isle of Man, and many of the water bodies present are low in diversity, the small number of diverse lakes, ponds and dubs which are present contain some of the rarest Manx plant and invertebrate species, several of which are protected under the Wildlife Act 1990 (see Samson, 2002). The Manx Wildlife Sites Handbook criteria for selection of Wildlife Sites therefore includes all open water or wetland that has rich marginal, inundation or aquatic vegetation communities (dominated by native species) and/or is of particular value for invertebrates, birds, fish or bats (excepting sites that are severely and irreversibly degraded). It also includes all areas of swamp greater than 0.5ha in extent, or 0.25ha when in mosaic or juxtaposition with other semi-natural habitats, and dominated by one or more of a list of typical native swamp species (Keehan/MWT, 1999).

The Isle of Man is signatory to the Ramsar Convention, which requires Member States to promote “wise use of wetlands” and to protect sites which fulfil the criteria for Ramsar Site status. These criteria refer to wetlands as a whole rather than individual habitat types. They contain several points of relevance to open water, including:

Criterion 1: A wetland should be considered internationally important if it contains a representative, rare, or unique example of a natural or near-natural wetland type found within the appropriate biogeographic region.

Criterion 2: A wetland should be considered internationally important if it supports vulnerable, endangered, or critically endangered species or threatened ecological communities.

Criterion 3: A wetland should be considered internationally important if it supports populations of plant and/or animal species important for maintaining the biological diversity of a particular biogeographic region.

Criterion 4: A wetland should be considered internationally important if it supports plant and/or animal species at a critical stage in their life cycles, or provides refuge during adverse conditions.

It is likely that the majority of sites which are considered for Ramsar Site selection on the Isle of Man will contain at least an element of still water habitat.

Minimum selection criteria for still water habitats

Note: The majority of still water habitats will require a buffer zone of at least 5m from the water's edge to enable effective conservation measures.

Bog pools

- Upland bog pools will be selected as part of larger upland sites – see **Chapter 4.7 Upland Habitats**.
- All lowland sites with a single or several pools totalling 0.5ha or more (including any buffer zone) of bog pools will be considered for selection where the pools in question form part of a larger area of related semi-natural habitat;
- Semi-natural areas with a collection of bog pools totalling less than 0.5ha (including any buffer zone) will be considered for selection where the pools in question fulfil at least one of the Priority Site Criteria listed in Part 3.

Ponds and dubs

- An individual lowland pond/dub will be considered for selection where it fulfils one or more of the Priority Site Criteria listed in Part 3, *and* forms part of a larger area of related semi-natural habitat;
- Series and/or groups of lowland ponds will be considered for selection where they collectively fulfil at least one of the Priority Sites Criteria listed in Part 3. This includes groups of ponds which are situated within improved farmland.
- In all cases, priority will be given to sites which have a good structural diversity and/or a good range of both submerged and emergent plant species, including the swamp categories listed in Table 4.5.a.

Individual, isolated ponds and dubs will not usually be considered for ASSI selection except on species criteria, and upland ponds and dubs of interest will be selected as part of larger upland sites – see **Chapter 4.7 Upland Habitats**. Criteria for selection of brackish pools will be covered in **Chapter 4.6: Coastal habitats**.

Lakes and reservoirs

- Lakes and reservoirs will be considered for ASSI selection where they fulfil at least one of the Priority Site Criteria listed in Part 3 *and* have a high structural diversity including more than one of the swamp categories described in Table 4.5.a (above).
- Priority will be given to lakes and reservoirs which occur in association with other wetland habitats such as streams, curragh, and mire.

4.5.2 Ditch systems

Description and Distribution

When unspoilt by pollution or excessive nutrient enrichment, man-made drainage ditches can provide a collection of interrelated habitats capable of supporting a wide range of plants and animals typical of both stagnant and slow-flowing open water. Those which are managed sparingly, but on a regular basis, generally provide the best opportunities for emergent vegetation; more undisturbed ditches may offer better opportunities for semi-aquatic invertebrates (NCC, 1989 Ch6:5). In some cases they may provide the only source of continuous wetland habitat for native species which would otherwise have no place in a modern farmed landscape; for this reason they may contain both plant and invertebrate rarities.

There is rarely enough space in managed ditches to allow the formation of swamp except where the ditches have become choked with vegetation through lack of maintenance, or where small “bays” occur along the main course. However, ditches often provide good habitat for submerged and emergent plants, and they are particularly well-suited to those species which thrive in a mixed, “semi-aquatic” environment, e.g. insects such as dragonflies which rely on the adjacent terrestrial habitats for feeding territory.

The Isle of Man does not have a tradition of flood-meadow and levels management on a par with lowland England and Wales, but there are some areas of farmland, notably the Ballaugh Curragh, in which the wetland environment has a long history of water level control via ditches and sluices (MNH, 1998). These habitats complement the adjacent mire and scrub, extending the possibilities for native wildlife. The lack of an overall plan of management until relatively recently, and the varying nature of the land use and ownership over the years, do not detract from the value of the Ballaugh Curragh drainage ditches for wildlife; indeed, these factors have probably contributed to the ecological diversity of the system over a long period of time.

Classification of ditch systems

Drainage systems may or may not reflect an overall drainage scheme for an area; they may simply result from the efforts of several landowners working individually to control water levels on their land. They fall broadly into two types: those on peat soils, and those on mineral soils, which tend to have less acidic waters and a higher overall nutrient status. There are no established classification methods for species-rich ditch systems, which tend to fall into similar categories to ponds and streams, based on the quality of water involved. Ditches on the Isle of Man may contain any of the pond and swamp vegetation communities listed in Table 4.5.a above, with the exception of those which prefer large areas of open water.

Ditch system status

Lowland ditch systems have long been recognised as a potential haven for aquatic species, but have only been fully taken into account in national designations relatively recently. The series of Welsh and English SSSIs which includes the Gwent Levels (Rhumney and Peterstone) and the Somerset Levels aims to encompass the best examples of ditch system habitats, along with the adjacent wet grassland and mire habitats. On the Isle of Man, the central parts of the Ballaugh Curragh, which includes some ancient ditch and sluice systems of archaeological as well as ecological interest, has been protected by the Curragh Acquisition Act (1963) and subsequent transfers of ownership to the Manx Museum. The Manx Wildlife Trust also manages parts of the Curragh land, some of which contain associated ditches and drains.

Certain of the Ramsar Site criteria (see *Status of still water*, above) may apply to ditch systems within wetland areas, as does the overall requirement for Ramsar Convention members to practice wise use of wetlands.

Species-rich ditch systems are vulnerable to agricultural intensification, wholesale heavy mechanical clearance, and pollution from adjacent developments - hence good examples of thriving ditch system ecology are rare. Unfortunately, the interconnected nature of ditch systems means that the introduction

of foreign aquatic plants or invertebrates can soon affect the species composition of quite a large area; for this reason it is important to safeguard species-rich ditch systems from thoughtless introductions, which can be difficult or impossible to reverse. Pollution incidents can likewise affect ditch systems very drastically. The *Code of Good Agricultural Practice for the Protection of Water* is aimed at preventing damaging pollution events that could otherwise find their way into ditch systems with catastrophic effects on the ecology.

Minimum selection criteria for ditch systems

Note: The majority of ditch habitats will require a buffer zone of at least 5m from the water's edge to enable effective conservation measures.

Only well-established systems of drainage ditches with a diverse ecology will be considered for selection.

- The best example of a lowland ditch system on peat soils will be considered for selection, as will the best example of drainage ditches on mineral soil; both examples should fulfil at least one of the Priority Site Criteria listed in Part 3.
- When choosing the best example of each type of ditch system, special priority should be given to those systems which are known to support rare species that wholly or partly rely on drainage ditches for their survival on the Island.

4.5.3 Flowing water

Description and Distribution

The short catchments and steep terrain of the Isle of Man have resulted in many small rivers and streams, the majority of which fall into upland or upland margin categories by British standards. Few get a chance to develop to more than two or three metres in width, and there are no extensive lowland river systems with accompanying high levels of sediment and nutrient build-up. However, the low-lying parts of the Island do support small, relatively eutrophic rivers with a range of associated marginal habitats, and there are some examples of rivers that follow a full course of upland, marginal, lowland and estuarine habitats before entering the sea. The more ecologically diverse rivers and streams on the Island all occur in close association with related habitats such as moorland, glens, wooded valleys and unimproved farmland.

The dynamic nature of river systems has led to a long history of efforts to control their natural fluctuations and changes in course, in particular to avoid flooding of farmland and settlements, but also to alter meandering watercourses to a more convenient and controllable structure. Few rivers are unmodified by man, either by direct means or as a result of exacerbated erosion due to human factors upstream. This is the case on the Isle of Man as elsewhere in the British Isles – although effects of sediment loading and other features of large lowland river systems are obviously less marked here. If possible at least one representative watercourse should be selected which shows a natural progression from fast upland stream to a slower, more level, lowland river as it enters the sea, in order to ensure that all types of ecological niche are covered by the designation.

Classification of flowing water

Flowing water habitats on the Island fall broadly into upland and lowland rivers and streams, with the majority likely to be wholly or partly covered by the Upland habitats criteria described in Chapter 4.7. Lowland stretches of rivers such as the Sulby and the Silverburn may be affected by human intervention and enrichment from farmland runoff, but not to the extent of heavily nutrient-enriched English lowland river systems. Bays and slower-flowing stretches of Manx rivers may therefore support a variety of the swamp and emergent vegetation communities described in Table 4.5a above, such as common reed *Phragmites australis* and greater reed-mace *Typha latifolia*, as well as submerged aquatics such as common water crowfoot *Ranunculus aquatilis*.

Flowing water status

The importance of unspoilt high-quality watercourses on the Island for sport fishing and water supply is

reflected in a range of protective measures. Game fisheries are protected by a variety of laws and bylaws, both from pollution incidents and from the activities of poachers. *The Code of Good Agricultural Practice for the protection of water* is aimed at protecting watercourses from polluting agricultural practices. Whilst upland and smaller streams are frequent on the Island, relatively unmodified stretches of semi-natural lowland rivers are rare, and this is reflected in the Manx Wildlife Sites Handbook, which recommends that all natural watercourses which exhibit no obvious artificial modification to bed and water level and have a high proportion of semi-natural habitat on both banks should be considered for selection as a Wildlife Site (Keehan/MWT, 1999).

In the wider context, *Water courses of plain to montane levels with the Ranunculion fluitantis and Callitriche-Batrachion vegetation* are now an EC Habitats Directive Annex 1 habitat for which the UK has a special responsibility. Manx rivers have some affinities with Subtype 3 of this river category, which relates to less nutrient-rich rivers in the north and west of Britain, often on upland margins.

Certain of the Ramsar Site criteria (see *Status of still water*, above) may apply to rivers and streams including estuarine reaches; as does the overall requirement for Ramsar Convention members to practice wise use of wetlands.

Minimum selection criteria for flowing water

Note: The majority of flowing water habitats will require a buffer zone of at least 5m from the water's edge to enable effective conservation measures.

Please note that river estuary saltmarsh will be covered in **Chapter 4.6: Coastal habitats**, and upland watercourses will generally be selected according to the Upland habitats criteria in **Chapter 4.7**.

- The best example of a natural and unmodified river from source to sea should be selected along with its catchment of tributaries;
- Shorter sections of lowland river which fulfil at least one of the Priority Site Criteria listed in Part 3 will be considered for selection where they possess a good range of structural diversity and good examples of at least two of the vegetation communities described in Table 4.5a above and/or other key habitats such as semi-natural broadleaved woodland cover and valley mire;
- In all cases, priority will be given to watercourses which occur in association with other semi-natural habitats such as wet grassland, wet woodland and scrub, and valley mire.



Fast-flowing water at Dhoon Glen ASSI creates high humidity, encouraging lots of fern growth.

4.6 Coastal habitats

Description

Defining exactly what is coastal and what is inland is not always straightforward. In general, coastal habitats may be referred to as either “maritime”, “sub-maritime” or “para-maritime”:

- Strictly **maritime** habitats are strongly affected by salt spray and exposure to strong winds and high humidity. It is rare for such habitats to occur at more than a narrow field’s width from the shore;
- **Sub-maritime** habitats may continue for one or two fields’ distance from the shore, and have less extreme conditions. Both maritime and sub-maritime habitats attract a distinctive range of flora and fauna, including some specialised plants and animals which only occur on the coast;
- **Para-maritime** areas are affected by the climate of the sea (i.e. high humidity, relatively small temperature range and exposure to strong winds) but not by the high soil salinity and extreme conditions that occur near the shore. In this sense, much of the Isle of Man could be said to have para-maritime conditions.

For the sake of clarity, this chapter covers strictly **maritime** habitats only, i.e. those which occur on rocks, sands, mud and cliffs between astronomical low tide level and points inland which are regularly within range of salt spray (Ratcliffe, 1977). Sub-maritime habitats all have strong associations with inland habitat types, and will therefore be covered in the relevant chapter – e.g. the Coastal grassland section of Chapter 4.2: Lowland grasslands, and the Dry scrub section of Chapter 4.1: Woodland.

This chapter may need to be considered in tandem with the separate document entitled *Guidelines for the selection of Marine Nature Reserves in the Isle of Man*, which covers sub-tidal marine communities, some of which relate closely to the intertidal habitats described below.

Assessing coastal habitats

All of the points on the Checklist of Priority Sites Criteria listed in **Part 3** are relevant to coastal sites - with the possible exception of points relating to Management History, as there will be some coastal habitats which have never received any management intervention. In all cases, a key consideration will be whether the coastline is generally soft – and hence liable to shift and change both in structure and actual location – or hard, with a relatively constant substrate. In addition to this, the following criteria are of particular importance when assessing coastal sites:

Size

One reason that coastal habitats have all been covered within the same chapter is the transitional nature of the terrain involved: habitats on the coast usually fall into zones, which grade into each other and represent an interrelated series, often in a state of continual development. Site assessments should therefore take into account the need to include all of the habitats that make up a particular coastal ecosystem, rather than just one or two features. This ensures not only that the full biodiversity of the coastline in question will be protected, but also that some of the processes are safeguarded – such as dune system development – for which adjacent habitats are needed in order to accommodate species which are naturally advancing or retreating as the coastal profile develops. This requirement tends to favour larger sites, and the largest examples of particular habitats.

The positive effects of size apply to species populations as well as habitat extent. The presence of breeding seabird colonies and other localised species of interest means that coastal sites are more likely to be chosen if they support the largest and/or most thriving communities of a particular species. The need for this is increased by the restricted nature of some coastal locations (see Location, below). However, it should be borne in mind that steep and/or vertical cliff habitats are unlikely to register a high area on orthographic maps, hence area is not the primary attribute for selection of cliff coasts (NCC, 1989).

Location

Although the Isle of Man is well-endowed with coastline, some coastal habitats are naturally restricted, and for certain features there is very little choice of good sites suitable for ASSI designation. The location of a coastal site is therefore a crucial consideration. There are several reasons for this:

- **Geology.** Most of the Island has a Manx slate bedrock with neutral to acid soils; the small, localised limestone features have a different ecology to the rest of the Island's coastline and as such they do not offer much choice in terms of site comparison for selection;
- **Aspect.** Most sloping or steep coastal sites are one-sided in nature, i.e. the zoning and ecology of the habitats present depends on their position in relation to the sea, which is usually only along one side of the site. The species present will therefore be those which are suited to the aspect, whether it is a cool north-facing cliff or a low south-facing limestone shelf. South-facing cliffs are rare on the Isle of Man; their sunny aspect appears to have a positive effect on the species diversity of both plant and invertebrate communities of coastal species (Lamb, 1999; Boyce & Fowles, 1989).
- **Exposure.** Very exposed coastal habitats are associated with more specialised plants and animals that can survive the harsh conditions and benefit from the lack of competition. More sheltered sites, particularly those on which a good level of drift material and soil has built up, may support robust weedy species typical of ruderal vegetation inland. Both these types need to be recognised as important;
- **Animal behaviour patterns.** Location affects fauna just as much as flora, either because of animal behaviour patterns – eg. preferred seal pupping beaches – or for the same aspect- and exposure-related reasons that influence plant communities;
- **Coastal development.** Location does not just relate to existing features; it is also important for the future development of habitats where these are drifting or transitory in nature. For example, sand dunes and saltmarsh are both vulnerable to loss from coastal "squeeze", man-made damage and modification on sites where there is no adjacent area onto which the habitat can shift if the coastline in question is retreating or accreting. The availability of adjacent habitats is therefore a positive factor in selecting a good example of a developing coastal ecosystem (NCC 1989). This includes adjacent inland habitat types such as unimproved grassland and scrub.

Diversity

Site selection should aim to include not just the full diversity of habitats and species on a site, but also the full range of transitional, zonal and successional habitat types. This is particularly important on soft coastlines, where dunes, slopes and soft cliffs exist as a continually developing system, with an ecology that is adapted to the shifting substrate.

Rarity, fragility and vulnerability

The restricted and specialised nature of some coastal communities makes them vulnerable to modification and damage. This is partly due to the difficult conditions on exposed coastal sites, where recolonisation may be slow. It is also due to the relatively restricted availability of suitable adjacent habitats for beleaguered species to move onto if conditions become inhospitable.

Whilst capable of withstanding very tough conditions, plant communities which are adapted to harsh environments are often by their nature very fragile when it comes to physical damage from vehicle access or heavy trampling. Rarity, fragility and vulnerability of coastal habitats should therefore be given special attention when comparing sites for selection, and unspoilt sites with especially rare or fragile species and habitats should be given priority where possible.

Coastal habitat types

For the purpose of these Criteria, coastal habitats on the Isle of Man have been grouped as follows:

Intertidal
Strandlines and shingle
Dunes
Saltmarsh
Brackish pools
Sea cliffs, caves and slopes

Elements of these habitats, and, where applicable, the National Vegetation Classification (NVC) categories they most closely resemble, are listed in detail on in sections 4.6.1-4.6.5 below.

4.6.1 Intertidal habitats

Intertidal habitats on the Isle of Man are currently assessed using the UK document *Guidelines for selection of biological ASSIs: Intertidal habitats and saline lagoons* (JNCC), in conjunction with local knowledge.

4.6.2 Strandlines and shingle

Description and Distribution

Strandlines, where the tide deposits loose pebbles, litter and organic material, may occur on any shore where there is space for deposits to build up on a regular basis. Whilst obviously a very mobile and changeable habitat, strandlines nevertheless have a distinctive ecology that includes characteristic annual vegetation, invertebrate communities and consequent opportunities for feeding birds. **Shingle** is broadly and simply defined as sediment particles that are larger than sand or gravel and smaller than boulders. This definition encompasses a wide range of particle sizes and mixtures, and may be differentiated still further where wind and wave action sculpts the loose sediments into permanent or semi-permanent structures such as beaches, spits and banks.

The likelihood that strandline and shingle structures will form a suitable home for flora and fauna depends greatly on the exact size and mix of particles involved, the degree of tidal exposure, and the stability of the shingle. For example, a very exposed area of shingle with large pebbles and few finer particles, which is regularly shifted about by fierce tidal action, is unlikely to support any vegetation at all, and very few invertebrates. However, an area of relatively stable shingle with some smaller particles amongst the pebbles will offer a growing medium to the small number of plant species that can stand the extreme conditions – which include drought, heavy salt load, and strong winds.

The Isle of Man has a great variety of different types of beach throughout its coastline, encompassing several distinctive strandline and shingle habitats within a small area (Lamb, 1999). Shingle and strandlines therefore represent a significant element of Manx coastal ecology, with vegetated shingle being a particularly valuable habitat due to its relative rarity. At their most diverse, vegetated shingle structures can support a very distinctive range of plants and an abundance of birds and invertebrates. Shingle beaches also provide good haul-out and pupping areas for seals.

The designation of the Ayres National Nature Reserve, which contains the largest area of shingle on the Island, was partly in recognition of the importance of this habitat for wildlife. Other areas of shingle occur on beaches along the east coast of the Island, such as Laxey beach, and on the south coast, notably Sandwich shore. Strandlines occur all round the Island, but tend to build up more in places where flat or gently sloping beaches face the prevailing currents, particularly the south and south west coasts.

Worldwide, shingle is home to rare plants and invertebrates, significant populations of breeding birds, and many seal territories, notably grey seals, which utilise both vegetated and non-vegetated shingle beaches as haul-out and pupping areas. Grey seals *Halichoerus grypus* are among the rarest seals in the world: around 40% of the global population of grey seals live and breed around the coasts of the British Isles and Isle of Man, which therefore have a collective global conservation responsibility for this species (see www.jncc.gov.uk, 2005). Several other protected species on the Isle of Man rely on beaches, notably little tern *Sterna albifrons*, which nest on shingle at the Ayres, and plants such as oysterplant *Mertensia maritima*; both of which are protected under the Wildlife Act 1990. Strandlines around the Island on both sandy and shingle shores also support a range of feeding birds, notably chough *Pyrrhocorax pyrrhocorax*, particularly in winter, and turnstone *Arenaria interpres* and a roosting site for other beach feeders at high tide. Plants of note include Isle of Man cabbage *Coincya monensis ssp monensis* and grass-leaved orache *Atriplex littoralis*, both protected under Schedule 7 of the Wildlife Act 1990.

Classification of strandlines and shingle

Vegetation of strandlines and shingle around the Island may be classified broadly as outlined in **Table 4.6.b**. Both **annual** vegetation of strandlines, which may occur on both shingle and sandy beaches, and **perennial** vegetation of shingle banks, are important for wildlife conservation. However, it should be noted that annual strandline vegetation is commoner worldwide than perennial vegetation of shingle banks, which is very restricted, with few extensive examples even within the British Isles (www.jncc.gov.uk, 2005).

Strandline and shingle status

The 1995 Isle of Man Ecological Habitat Survey mentions just four sites around the Island's coast where strandline vegetation was found during the Phase 1 Habitat Survey. 876ha of coastal habitats were mapped in total during this Survey, of which only 43ha is mapped as shingle/cobbles (Sayle *et al*, 1995). It is not recorded which of these shingle patches had any vegetation present, and it is likely that some areas described as strandline vegetation were on shingle structures with perennial as well as annual plants present. The Terrestrial Survey of the Coast of the Isle of Man (Lamb, 1999) goes into more detail, listing "shingle above the high tide mark" as a major habitat element on 25 of the 58 coastal sections surveyed. However, the shingle found rarely supported any permanent plant or invertebrate communities due to the very exposed nature of the shore. A rough estimate based on surveys of key sites suggests that less than a quarter of the shingle mapped supports any kind of vegetation, making vegetated shingle one of the rarest habitats on the Island (BDIS). Good examples of strandline communities are slightly more frequent, but not abundant due to the very exposed, rocky nature of many Manx beaches and the practice of clearing strandline depositions from popular beaches in the holiday season. The importance of both strandlines and shingle is reflected in the Manx Wildlife Sites Handbook, which recommends Wildlife Site status for well-vegetated strandlines and shingle beaches with good invertebrate and bird populations (Keehan, 1999).

Worldwide, vegetated shingle is a category which has received a lot of conservation concern. Although shingle beaches are very familiar to visitors of British coastal resorts, vegetated shingle is in fact an internationally scarce and important habitat, restricted to the coasts of Britain, Ireland, North-west Europe, Japan, and New Zealand, and never in great abundance in any of these areas. There is known to be around 4,000 ha of stable or semi-stable vegetated shingle around the coast of the UK, which is widely distributed (often restricted to small patches isolated from other shingle areas) and which exhibits a wide range of variation. There are only a few extensive examples of perennial vegetation of stony banks in Europe, and the UK hosts a significant part of the European resource of this habitat. In recognition of its restricted range and conservation importance, Coastal Vegetated Shingle is listed as a UK Biodiversity Action Plan Priority Habitat, whilst both Annual vegetation of drift lines and Perennial vegetation of stony banks are listed as Annex 1 habitats under the EU Habitats Directive. Full details of these and related habitats and species can be found at www.jncc.gov.uk.

Minimum selection criteria for strandlines and shingle

Strandlines will not constitute reason for ASSI selection on their own except where associated with perennial vegetation of shingle banks (see below) and/or on the grounds of species criteria. However:

- the presence of 100m or more of unpolluted strandline with associated vegetation and/or bird or invertebrate populations is likely to fulfil the Priority Site Criteria for Naturalness and Representativeness and favourable Ecological Position, as described in Part 3, for a coastal site with one or more other habitat features. Good examples of strandline communities will therefore be a positive factor in the selection of larger coastal sites.

Bare shingle will not constitute reason for ASSI selection as a habitat on its own, but should be included within the boundary of designated coastal sites wherever it occurs adjacent to other selected habitats e.g. dunes or vegetated shingle.

Vegetated shingle, whether sandy in nature or not, will qualify for selection where:

- The shingle supports a range of species associated with any of the NVC categories described in Table 4.6b, and the total extent is greater than 0.2ha*, or
- The total area of vegetated shingle as described above is greater than 0.05ha* and the area fulfils at least one of the Priority Site Criteria listed in Part 3, with priority given to sites which are known to support populations of rare plants and/or animals.

*Because of the naturally narrow nature of most vegetated shingle habitats, it may be convenient to think of shingle in terms of length of site as well as total area:

0.2ha = 1km of 2m width shingle, 0.5km of 4m width shingle, 250m of 8m width shingle

0.05ha = 250m of 2m width shingle, 125m of 4m width shingle, 60m of 8m width shingle etc.



Blackthorn *Prunus spinosa*, usually a substantial shrub, has an almost prostrate form when growing on the harsh environment of bare shingle

Table 4.6b: Broad categories of strandlines and shingle habitats on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (See Rodwell, 1992 ⁵)	Rarities present?	Protected species present?
<p><i>Sandy strandline</i> Consists of seaweed, driftwood and litter deposits, rapidly-changing according to tides and often greatly increased by storm events. Vegetation is likely to be limited to salt-tolerant, weedy annual species. This type of strandline may be important in the formation of embryo dunes (see 4.6.3 below)</p>	Not usually rich in plant species, but can support rich terrestrial and marine invertebrate communities	None, but sand couch foredune vegetation SD4 (see 4.6.3, below, and Table 4.6.c) may occur in association with sandy strandlines.	May form part of breeding and/or feeding territory for rare birds (see right)	Grass-leaved orache <i>Atriplex littoralis</i> ; oysterplant <i>Mertensia maritima</i> . A range of birds may feed along strandlines, including chough <i>Pyrrhocorax pyrrhocorax</i> , starling <i>Sturnus vulgaris</i> , and turnstone <i>Arenaria interpres</i> .
<p><i>Sandy shingle</i> A mixed substrate, with smaller sand and gravel –sized particles caught between larger pebbles, providing a potential growing medium for perennial as well as annual plants. This type of strandline may also be important in the formation of embryo dunes (see 4.6.3 below)</p>	May be species-rich, especially where perennial vegetation is established	<p>SD3: <i>Matricaria maritima</i> – <i>Galium aparine</i> (sea mayweed/cleavers) strandline community No Island NVC sample</p> <p>SD4: (sand couch foredune – see above and Table 4.6.c) SD19: <i>Phleum arenarium</i> – <i>Tortula ruralis</i> (sand timothy grass and moss) dune annual community – also occurs in disturbed areas of sand dunes (see Table 4.6.c below) No Island NVC sample</p>	Sea holly <i>Eryngium maritimum</i>	Oysterplant (as above); Sea wormwood <i>Seriphidium maritimum</i> ; Grass-leaved orache <i>Atriplex littoralis</i> ; Hare's-foot clover <i>Trifolium arvense</i> ; Isle of Man cabbage <i>Coincya monensis subsp monensis</i> ; viper's bugloss <i>Echium vulgare</i> . A range of birds may feed occasionally on vegetated sandy shingle. Some birds such as little tern <i>Sterna albifrons</i> and ringed plover <i>Charadrius hiaticula</i> may nest on sandy shingle.

<p>Shingle Sharply-drained, large pebbles with limited, mostly annual vegetation. May build into sloping banks, the inland side of which can develop perennial vegetation and associated invertebrate communities. The ability of this vegetation to trap sand high up the shore may play a part in dune formation, especially in an island setting.</p>	Not usually	<p>SD1: <i>Rumex crispus</i>-<i>Glaucium flavum</i> (curled dock/yellow horned poppy) shingle community No Island NVC sample</p> <p>SD2: <i>Honckenya peploides</i> – <i>Cakile maritima</i> (sea sandwort/sea rocket) strandline community 2 Island NVC samples</p>	Yes; several rare plants (see right) and invertebrates may occur on this habitat; also some unusual/locally distributed plants such as Sea kale <i>Crambe maritima</i> .	Oysterplant <i>Mertensia maritima</i> ; NB this species is also nationally scarce in the British Isles. Terns <i>Sterna</i> spp. may nest on shingle areas.
<p>Bare shingle This consists of pebbles and cobbles only, often within the most exposed tidal ranges.</p>	No	Not applicable; no vegetation present	May form part of breeding and/or feeding territory for rare birds (see right)	May be used for haul out or pupping area by Schedule 5 protected Grey seal <i>Halichoerus grypus</i> ; breeding terns <i>Sterna</i> spp.



Sea sandwort *Honckenya peploides*, growing on shingle at the Ayres ASSI and National Nature Reserve

4.6.3 Dunes

Description and Distribution

Coastal sand dunes develop where there are onshore prevailing winds and a sufficient supply of sand, coupled with a large enough area of beach, to allow sand deposits to build up and dry out between high tides (Ratcliffe, 1977). A small number of highly specialised grasses, principally marram grass, lyme grass and sand couch, can colonise the shifting sand and develop a vegetation community which both stabilises the developing dunes and helps further build-up by trapping more sand.

Sand dune vegetation forms a number of zones in relation to the time elapsed since the sand was deposited, the degree of stability which it has attained, and the local hydrological conditions:

- **Embryonic and mobile dunes** occur mainly on the seaward side of a dune system where sand deposition is occurring and occasionally further inland in blow-outs. They support very few plant species, the most characteristic being marram grass *Ammophila arenaria*.
- **Semi-fixed dunes** occur where the rate of sand accretion has slowed but the surface is still predominantly bare sand; marram is still common but there is an increasing number of other species.
- **Fixed dune** grassland forms largely closed swards where accretion is no longer significant, the surface is stabilised and some soil development has taken place. Calcareous fixed dunes support a particularly wide range of plant species. On dunes which have become acidified by leaching or **de-calcified**, acid dune grassland or dune heaths develop.
- **Dune heaths** are usually dominated by ling *Calluna vulgaris* and bell heather *Erica cinerea*.
- **Dune slack** vegetation occurs in wet depressions between dune ridges; it is often characterised by creeping willow *Salix repens* and mosses.

Fixed dunes and well-developed dune grassland and heath are potentially very species-rich, attracting warmth-loving insects as well as birds and a wide range of grassland and heathland plants. Dune habitats are further diversified by the presence of dune slacks. All these habitats are slow to recover from damage, and are hence vulnerable to human disturbance and modification, particularly heavy vehicle access, fires, and disturbance of ground-nesting birds. Once "reclaimed" for agriculture or development a dune is lost for good, and changes in coastal defences or adjacent land use can alter the whole dune system by affecting local sand deposition patterns. For these reasons, much UK duneland has been lost or damaged, and a high proportion of the remaining habitat is protected under UK and/or EU legislation.

Dunes can form in a variety of places, always in exposed conditions. Because of the need for a large area of more or less flat, open land, dunes occur predominantly around the north coast of the Isle of Man, where hindshore dunes form on large, open areas of coast (e.g. the Ayres). Elsewhere, the coastline is either too steep and rocky to allow dune formation, or the areas of sand build-up are on beaches adjacent to built-up areas such as Peel and Port Erin.

Classification of dunes

Vegetation of dunes around the Island may be classified broadly as outlined in **Table 4.6.c**.

Dune status

The Isle of Man has one large area of dunes across the north coast, with more open dunes concentrated in the northwest (eg. Blue Point) and dune heath most prevalent in the central to northeast Ayres and Point of Ayre. Elsewhere, only tiny patches of sand deposits occur, none of which show the range of dune development found on the north and northwest coasts. Dune heath (54% of coastal habitats overall) is by far the most extensive dune type, with dune scrub and dune slack the rarest. Although they make up a high proportion of the total area of coastal habitat on the Island, dune habitats are rare overall, with even dune heath making up just 0.55% of the Island total, or 1.89% of the semi-natural habitat total. The rarity and vulnerability of dune habitats is recognised in the Manx Wildlife Sites Handbook: all areas of dune except severely degraded examples are recommended as eligible for Wildlife Site selection (Keehan, 1999).

Dune habitats are treated as threatened habitats throughout the British Isles, with Coastal Sand Dunes listed as a UK Biodiversity Action Plan Priority Habitat. At the European level, a range of dune types are listed in Annex 1 of the EC Habitats Directive, including Embryonic Shifting Dunes, Dunes with *Salix repens ssp. argentea* (Salicion arenariae), Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes'), Fixed dunes with herbaceous vegetation ('grey dunes') (a priority habitat) and Atlantic decalcified fixed dunes (Calluno-Ulicetea) (also a priority habitat). Humid dune slacks are also listed, with a note to the effect that this habitat is more frequent in the UK than elsewhere, hence the UK has a special responsibility for its conservation (www.jncc.gov.uk, 2005).

Minimum selection criteria for dunes

- The largest examples of each of the following dune types will qualify for selection:
 - Mobile foredune and "white" or "yellow" dune with:
 - marram grass *Ammophila arenaria*;
 - lyme grass *Leymus arenaria* and
 - Sand couch *Elytrigia juncea*;
 - Semi-fixed "grey" dunes;
 - Fixed dune grassland;
 - Dune heath.
- Any dune system greater than 2ha in extent which fulfils at least one of the Priority Sites Criteria listed in Part 3 will qualify for selection, with priority given to sites with good naturalness and representativeness (i.e. those sites with least disturbance, modification and damage).
- Selected areas of dunes should be included with as much adjacent semi-natural coastal habitat as possible, including bare and vegetated shingle banks, strandline and coastal scrub.
- When selecting dune systems, special consideration should be given to the landward boundary, which should encompass sufficient land to allow the dune system to develop onto suitable adjacent habitat where applicable.



Hare's-foot clover *Trifolium arvense*, a protected species, growing on semi-fixed dune at Ramsey Mooragh Shore ASSI

Table 4.6c: Broad categories of dune habitats on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (See Rodwell, 1992 ⁵)	Rarities present?	Protected species present?
<p>Mobile foredune and yellow dune</p> <p>Mobile foredune consists of shifting sand on the seaward side of a dune system, behind which larger and progressively more stable dunes can build up. Sandy strandlines (see 4.6.2 above) may lie adjacent to this habitat, and strandline vegetation can play a part in the formation of embryo dunes. Yellow dune is slightly less mobile, and generally supports a loose mat of plant growth of one or two species such as marram grass or lyme grass. A few specialised invertebrates and a very limited number of plants can survive in these habitats, which are subject to constant change and abrupt re-grading following strong winds and storm events.</p>	No	<p>SD4: <i>Elymus farctus</i> ssp <i>boreali-atlanticus</i> (sand couch – now called <i>Elytrigia juncea</i> ssp <i>boreoatlanticus</i>) foredune One Island NVC sample SD5: <i>Leymus arenarius</i> (Lyme grass) mobile dune community No Island NVC sample SD6: <i>Ammophila arenaria</i> (marram grass) mobile dune community No Island NVC sample</p>	May form part of breeding and/or feeding territory for rare birds (see right)	Not usually, although some bird species such as little tern <i>Sterna albifrons</i> may utilise foredunes as nesting areas.
<p>Fixed and semi-fixed dune</p> <p>Semi-fixed dunes are often referred to as grey dune, and are usually well-vegetated, with more varied vegetation than yellow dunes. This vegetation becomes progressively more consolidated inland, until it grades into fixed dune grassland and/or heath. Dune scrub may also occur, although it is rare on the Island due to the small areas of dune involved.</p>	May be species-rich in both plant and invertebrate communities	<p>SD7 <i>Ammophila arenaria</i> – <i>Festuca rubra</i> (marram grass and red fescue) semi-fixed dune community 2 Island NVC samples SD8 <i>Festuca rubra</i> – <i>Galium verum</i> (red fescue and lady's bedstraw) fixed dune grassland One Island NVC sample SD10 <i>Carex arenaria</i> (sand sedge) dune community No Island NVC sample SD12 <i>Carex arenaria</i> – <i>Festuca ovina</i> –</p>	Rare and/or locally distributed plants and invertebrates are a possibility	A range of orchid species grow on developed dunes, including Twayblade <i>Listera ovata</i> , Autumn lady's tresses <i>Spiranthes spiralis</i> , Pyramidal orchid <i>Anacamptis pyramidalis</i> and early marsh orchid <i>Dactylorhiza incarnata</i> . Larvae of the scarce crimson and gold moth <i>Pyrausta sanguinalis</i> feed on thyme in this habitat.

		<p><i>Agrostis capillaris</i> (sand sedge, sheep's fescue and common bent grass) dune grassland</p> <p>Disturbed areas of fixed dunes may have SD19: <i>Phleum arenarium</i> – <i>Tortula ruralis</i> (sand timothy grass and moss) dune annual community</p> <p>No Island NVC sample</p>		
<p>Dune slacks</p> <p>Damp depressions and hollows in between dune ridges may have a high water table for all or part of the year, allowing marshy vegetation to survive. The degree of brackishness of dune slacks depends on the proximity to the sea and level of exposure to salt spray deposition; in general they are classed as freshwater, with truly salt slacks classed as types of saltmarsh even when on sand rather than mud (see Table 4.6.d). Dune slacks on the Island are rare (restricted to the Ayres) and usually small (under 1ha).</p>		<p>SD15 <i>Salix repens</i> – <i>calliargon cuspidatum</i> (creeping willow and moss) dune slack community</p> <p>SD16 <i>Salix repens</i> – <i>Holcus lanatus</i> (creeping willow and Yorkshire fog) dune-slack community</p> <p>One Island NVC sample</p> <p>Sand sedge (SD10 – see above) may also occur around dune slacks</p> <p>SD17 <i>Potentilla anserina</i> – <i>Carex nigra</i> (silverweed and common sedge) dune-slack community</p> <p>No Island NVC sample</p>	<p>Dune slacks may support very locally distributed invertebrate populations.</p>	<p>Orchids may thrive in dune slacks, notably Northern marsh orchid <i>Dactylorhiza purpurea</i>. Wintering waders such as curlew <i>Numenius arquata</i> and lapwing <i>Vanellus vanellus</i> may be attracted to dune slacks as feeding areas.</p>

4.6.4 Saltmarsh

Description and Distribution

Saltmarsh is a habitat composed of species that can withstand high salinities and regular flooding from the tide. It forms in areas with fine sediments and usually some protection from the worst onslaught of the storm waves. On the Isle of Man it is found fringing tidal rivers and in coastal bays with some protection from the waves.

It can incorporate a number of zones of different vegetation. Seaward there is often a pioneer community which must be tolerant of extended periods of submergence in saltwater and washing by wave and tide. Higher up the beach the communities are more terrestrial in character. These can form a succession through which sediments are gained to the land, or a zonation that is more static but protects the higher areas from erosion by the sea, the roots of the vegetation binding the sediments together and the shoots breaking the force of the waves. This range of communities can be shortened by erosion from the sea, or due to 'coastal squeeze', in which the landward boundary of a saltmarsh is curtailed by a bank or sea wall whilst the marsh moves up against it with a rise in sea level or due to sediment movements.

Saltmarsh communities vary geographically and with local physical conditions (tidal range, salinity of the water and sediment type). Moving northwards in the British Isles, there is a steady decline in the numbers of saltmarsh plant species and particle size tends to increase in a northerly and westerly direction, affecting soil aeration, drainage and nutrient status, with a greater number of species growing on finer soils (Ratcliffe 1977). Northern saltmarshes, such as those found in the Isle of Man, therefore tend to be botanically poorer than those in the south of England.

Saltmarshes often form part of a larger system of coastal habitats, which can be of importance for birds and invertebrates. Associations with brackish habitats, in marshes or creeks, or with terrestrial habitats such as woodlands, can allow a higher diversity of species to live there. Such areas can form localised features that are important or rare in the British Isles. For instance, there are very few places in Britain where a gradual transition from saltmarsh to woodland can be seen (Ratcliffe 1977). On the island, Port Cornaa has a saltmarsh which is botanically poor, yet the proximity of other habitats such as woodland, has benefited the wildlife and Boyce and Fowles (1989) recorded a very good list of invertebrates in this habitat mosaic.

There are areas of coast, mostly too exposed and not silty enough for saltmarsh habitat to form, where saltmarsh species may be found as either strandline vegetation or on salt-sprayed cliffs. Such sites do not fall within this section. For instance, sea arrowgrass is found on salt-sprayed grasslands. It is therefore not a good indicator of the habitat, whereas saltmarsh rush is more specific in its habitat requirements.

Good stretches of saltmarsh vegetation can be found around the fringes of the Irish Sea, with some extensive areas in north-west England and south-west Scotland. However, the Isle of Man has very little area that could be termed 'estuary', where there would be sufficient shelter for fine sediments to fall and saltmarsh to form over an area of significant size. Saltmarsh grows best on fine sediments, but most of the coastal Manx sediment is too coarse. Derbyhaven has a fairly large area of mud but almost no saltmarsh vegetation. Those areas having an estuarine environment have been subject to development and drainage to form the port areas of Douglas, Castletown, Peel and Ramsey. In these areas saltmarsh may have been found in the past but only Ramsey retains this habitat, along the banks of the Sulby River. However, there are other small areas elsewhere on the island in estuaries and sheltered bays.

The Isle of Man Ecological Habitat Survey Phase 1 report recorded only 6.72 ha of saltmarsh in total. There are four main areas: Ramsey, Port Cornaa, Langness and Poyllvaish with small patches of saltmarsh species also present elsewhere. Langness is the largest site.

Saltmarsh sites on the Isle of Man

Langness has the largest patch of saltmarsh on the island, 3.44 ha (phase 1), around Poyll Breinn, the stinking pool. It is considered the best example of saltmarsh on the island (Gubbay 2000) and was grazed (at least parts of it) formally (phase 1 report). The phase 1 survey also shows a tiny patch further south (0.04 ha), another close to the golf course hotel (0.04 ha) and 7 small patches on St. Michael's Island totalling 0.21 ha. All of these are within the Langness, Sandwick and Derbyhaven ASSI, designated on 19th October 2000. It was listed as a Site of Marine Nature Conservation Importance by the Manx Wildlife Trust (Gubbay 2000). The site includes celery-leaved buttercup *Ranunculus sceleratus*, chestnut sedge *Blasmus rufus*, grass-leaved orache *Atriplex littoralis*, sea club-rush *Scirpus maritimus*, sea lavender *Limonium vulgare*, parsley water dropwort *Oenanthe lachenalii* and cord grass *Spartina anglica*.

Visits by Luff and by Boyce and Fowles have produced invertebrate lists. Boyce and Fowles (1989) found the weevil *Polydrosus pulchellus* ab. *insquamosus*, a nationally notable (Category Nb) species, and of a rare form previously found in Devon, Kent and Gloucestershire. This site is the northernmost recorded locality of the centipede *Schendyla peyerimhoffi*. The saltmarsh and adjoining wrack beds and pools are an important feeding ground and roost for ducks and waders. In the winter wigeon, teal, shelduck, curlew and golden plover are common and in summer lapwing nest here. Redshank were previously recorded nesting but have not returned in recent years.

Port Cornaa has a small saltmarsh of 0.72 ha (phase 1) lying behind the high shingle bar, astride the river. This saltmarsh is listed as a Site of Marine Nature Conservation Importance in the Isle of Man by Gubbay (2000). Formerly grazed, it has also often been flooded by fresh water when the river has been dammed for safe bathing (Garra 1972). The river Cornaa runs through it and it is backed by marshy grassland, with woodland close behind and an area of alder carr linked to the marsh. The saltmarsh drains through a creek system, leaving open mud at low tide. Some erosion has been noted by Spencer (pers. comm.) recently and some minor bank works may have been done to prevent the river from eroding the southern marsh.

Areas are dominated by either red fescue *Festuca rubra* or creeping bent *Agrostis stolonifera* with abundant sea arrowgrass *Triglochin maritimum* and sea plantain *Plantago maritima*. Sea rush *Juncus maritimus* is locally abundant and saltmarsh rush *Juncus gerardii* locally frequent. Glasswort *Salicornia europaea* agg. has also been recorded (Allen 1984). Parsley water-dropwort *Oenanthe lachenalii* is recorded, and two uncommon sedges, distant sedge *Carex distans* and false fox-sedge *C. obtrubae*. The area to the east of the river is more complex and supports an abundance of sea rush, saltmarsh rush and rock samphire. A water crowfoot *Ranunculus trichophyllus* characteristic of brackish water has been recorded here, considered very local by Allen (1984).

Boyce and Fowles (1989) considered this area to be of considerable invertebrate interest, due to the close proximity of saltmarsh, pasture, river shingle and woodland, forming a mosaic of habitats, each perhaps small but adding greatly to the value of the others by their close proximity. A high species abundance should result. The saltmarsh, though low in plant diversity, had a significant invertebrate interest, the sheltered conditions and shallow silt and gravel channels providing habitat for ground-active species. Two ground beetles found here are scarce saltmarsh species with a restricted distribution in the British Isles: *Aepus marinus* (nationally notable Nb, widespread but local in the British Isles in intertidal habitats) and *Bembideon maritimum* (Nb in Hyman 1986, but not included in the revision of Parsons 1992).

At **Ramsey**, saltmarsh is stretched along both sides of the tidal Sulby River, including the largest area, at Poyll Dhooie, which used to be grazed. The alteration of the river mouth may have affected the distribution and extent of saltmarsh significantly. At Poyll Dhooie a tip created in the 1940s resulted in the loss of some of the saltmarsh (Spencer pers. comm.). Despite this, the area as a whole remains the second largest area of saltmarsh on the island. Most of the areas are dominated by red fescue. At the White Bridge reed swamp of *Phragmites australis* merges with the red fescue saltmarsh, the reeds continuing upstream in fresher water, lining the banks, as typical of an upper estuary situation. Sea aster *Aster tripolium*, grass-leaved orache *Atriplex littoralis* and long-spiked glasswort *Salicornia dolichostachya* are all recorded. There is an area with brackish pools fringed with saltmarsh rush *Juncus gerardii*. Here rabbits graze the sea arrowgrass *Triglochin maritima* and creeping bent *Agrostis stolonifera*.

Poyllvaish has scattered areas of ungrazed saltmarsh on carboniferous limestone mixed with strandline vegetation, shingle and scattered saltmarsh plants. It is dominated by red fescue *Festuca rubra*, with locally abundant common saltmarsh grass *Puccinellia maritima*. Sea aster *Aster tripolium*, glasswort *Salicornia europaea* agg. and sea purslane *Halimione portaculoides* (very local, Allen 1984) are found here (phase 1). The oyster plant *Mertensia maritima* (very rare; Allen 1984) was once recorded here c. 1866 but crushed by a passing cart and not seen since. To the south, one patch of the rare reflexed sea meadow grass *Puccinellia distans* was recorded in Allen (1984) by Poyllvaish Farmhouse, and celery-leaved buttercup *Ranunculus sceleratus* on the shore between this site and Ronaldsway.

Classification of saltmarsh

Saltmarsh on the Island may be classified broadly as outlined in **Table 4.6.d**.

Saltmarsh grows on fine sediments which settle out in sheltered areas of salty or brackish water. Because the sediment type, slope and level of brackishness are important, the changing physical nature of a site can affect the saltmarsh growing on it. On the Island, saltmarshes grow in three physiographic situations, estuarine (Sulby River), lagoonal (Cornaa), and beach plain (Langness and Poyllvaish).

Rodwell (2000) identified four saltmarsh NVC communities on the Isle of Man, of which two fall within the Atlantic Salt Meadows protected in Europe under the Habitats Directive. From the extreme lower end of the transition of communities comes the common eel grass *Zostera marina* which is sometimes washed up on Manx beaches around the island (Allen 1984; Rodwell 2000). *Zostera* beds (Rodwell's NVC SM1 community) are known to grow off St. Michael's Island / Langness and are likely to be found elsewhere by diving survey. However *Z. marina* is mainly found on sites constantly submerged by the sea (and some lagoons elsewhere), from just above the spring low water mark, down to 4m below. On the Isle of Man it is not recorded as a continuing seaward extension of saltmarshes and is likely to form separate, sub-littoral stands which may be best considered under the criteria for Marine Nature Reserves.

In the lower marsh SM13c *Puccinellietum maritimae*, *Limonium vulgaris* - *Armeria maritima* sub-community is characterised by Rodwell as having *Puccinellia maritima* as a constant constituent. In the mid-upper marsh red fescue saltmarsh SM16 *Juncetum gerardi* is characterised by the constants *Festuca rubra*, *Plantago maritima* and *Glaux maritima*. Higher still, the upper marsh community SM24 *Atriplici* - *Elymetum pycnanthi* has been identified, but this is invariably dominated by sea couch *Elytrigia atherica* (form. *Elymus pycnanthus*) though this has not been recorded on the Island.

A third characterisation of saltmarshes lies in their management. Grazed marshes form a different structure to the rank ungrazed marshes, resulting in a different invertebrate community and frequently different birds. Continuing traditional management may be important to the retention of some of the wildlife interest; a change in management can result in the development of a different community. Port Cornaa, Langness and Poyll Dhooie were formerly grazed but none of the sites were grazed when the phase 1 report was written.



Lesser sea-spurrey *Spergularia marina* on saltmarsh at Langness ASSI

Table 4.6.d: Broad categories of saltmarsh on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (See Rodwell, 1992 ⁵)	Rarities present?	Protected species present?
<p>Sea-clubrush swamp</p> <p>Typical of more or less still, brackish or salt water alongside estuaries, lagoons and the deeper areas of saltmarshes. Uncommon.</p>	No.	<p>S21 - <i>Scirpus maritimus</i> (sea club-rush, now known as <i>Bolboschoenus maritimus</i>) swamp No Island NVC sample</p>	Provides good bird cover, so may attract nesting or passing rarities.	
<p>Lower saltmarsh. This vegetation generally colonises the muddier, wetter parts of saltmarshes, and is regularly covered at medium to high tide. It often forms important feeding sites for wading birds.</p>	Not rich in plant species; may however support a diverse invertebrate population and associated feeding waders.	<p>SM8 Annual <i>Salicornia</i> (glasswort) saltmarsh community No Island NVC sample</p> <p>SM9 <i>Suaeda maritima</i> (annual sea-blite) saltmarsh community No Island NVC sample</p> <p>SM10 Transitional low-marsh vegetation with <i>Puccinellia maritima</i> (common saltmarsh-grass), annual <i>Salicornia</i> (glasswort) species and <i>Suaeda maritima</i> (annual sea-blite). No Island NVC sample.</p> <p>SM13 <i>Puccinellia maritima</i> (common saltmarsh-grass) community, esp. 13c <i>Limonium vulgare</i> - <i>Armeria maritima</i> (Sea lavender – thrift) sub-community One Island NVC sample.</p> <p>SM14 <i>Halimione portulacoides</i> (sea-purslane) saltmarsh community No Island NVC sample</p> <p>SM15 <i>Juncus maritimus</i> – <i>Triglochin maritimum</i> (saltmarsh rush – sea arrowgrass) saltmarsh No Island NVC sample</p>	Yes; several lower saltmarsh plant species are rare or uncommon on the Island, e.g. the more unusual <i>Salicornia</i> glasswort species.	<p>Protected plants may include <i>Halimione portulacoides</i>, <i>Eleocharis uniglumis</i> slender spike-rush, <i>Limonium vulgare</i> common sea-lavender, <i>Seriphidium maritimum</i> sea wormwood, <i>Oenanthe lachenalii</i> parsley water-dropwort.</p> <p>Protected bird species that may feed on lower saltmarsh include curlew <i>Numenius arquata</i> and shelduck <i>Tadorna tadorna</i>.</p>

<p>Middle saltmarsh. These usually resemble muddy, more or less closed grasslands, often dominated by red fescue growing as a flattened mat, or by tufts of saltmarsh rush. Can be grazed, although no middle saltmarsh on the Island is currently grazed.</p>	<p>Yes; may support a range of plants, invertebrates and associated bird life.</p>	<p>SM16 <i>Festuca rubra</i> (red fescue) saltmarsh community One Island NVC sample</p> <p>SM18 <i>Juncus maritimus</i> (saltmarsh rush) saltmarsh No Island NVC sample</p> <p>SM19 <i>Blysmus rufus</i> (saltmarsh flat-sedge) community No Island NVC sample</p>	<p>Yes; several unusual species occur in mid-range saltmarsh; <i>Blysmus rufus</i> is itself a rarity on the Island, and rare invertebrates and feeding birds are also a possibility.</p>	<p>Protected plants may include <i>Halimione portulacoides</i>, <i>Oenanthe lachenalii</i> (parsley water-dropwort), and <i>Seriphidium maritimum</i> (sea wormwood). Protected bird species that may feed on middle saltmarsh include geese and shelduck.</p>
<p>Upper saltmarsh. This tends to have a grassy cover, suitable for grazing – although there is no grazed saltmarsh on the Isle of Man, and grazed sites are becoming rarer throughout the UK. This is not a well-developed habitat on the Island, and upper saltmarsh species may occur only as a fringe amongst rocks and built-up areas on the landward edge of the marsh.</p>	<p>May be species-rich, but is not usually extensive enough on the Island to develop its full potential. Presence of wet hollows and pools can increase the diversity.</p>	<p>SM24 <i>Elymus pycnanthus</i> (sea couch) saltmarsh community One Island NVC sample – but note, <i>E. pycnanthus</i> (now re-named <i>Elytrigia atherica</i>) does not appear to have been recorded on the Island! In other respects the habitat does match this common NVC community.</p> <p>SM28 <i>Elymus repens</i> (common couch) saltmarsh community No Island NVC sample. Note: <i>E. repens</i> has been renamed <i>Elytrigia repens</i>.</p>	<p>Yes; some unusual plant species may occur, especially where pools are present - notably coastal sedges such as <i>Carex otrubae</i> false fox-sedge, <i>C. extensa</i> long-bracted sedge, and <i>C. distans</i> distant sedge and <i>Puccinellia distans</i>. Reflexed saltmarsh grass</p>	<p>Protected plants may include <i>Oenanthe lachenalii</i> (parsley water-dropwort), <i>Atriplex littoralis</i> (grass-leaved orache) and <i>Seriphidium maritimum</i> (sea wormwood). Protected bird species that may feed on lower saltmarsh include lapwing <i>Vanellus vanellus</i>, ringed plover <i>Charadrius hiaticula</i>, and starlings <i>Sturnus vulgaris</i>.</p>

Saltmarsh status

Wider context

Saltmarshes are a priority habitat for conservation in the EU (Annex 1 of the Habitats Directive: Atlantic salt meadows considered by the JNCC to be equivalent to SM10 to 20) and as a result, form parts of a number of Special Areas of Conservation in the UK. They are also listed as a priority habitat in the UK, resulting in the provision of a habitat action plan. There are more than 29,000 ha of saltmarsh in the UK (Saltmarsh Database, JNCC website 2003). About 80% of British saltmarsh is designated as SSSI (50% in north-west Scotland). In Northern Ireland 5 of 7 estuaries with saltmarsh are ASSI. Through the habitat action plan the UK aims to create 100 ha per year to maintain the total area (about 100 ha per year are lost) plus a further 40 ha per year to replace the 600 ha lost between 1992 and 1998.

Regionally (Colwyn Bay to Stranraer, and Isle of Man), Manx saltmarshes are tiny compared with the extensive saltmarshes of Lancashire, Cheshire and Cumbria. Less than 1% of the Manx coast is saltmarsh but this is 46% in Lancashire, 45% in Cheshire and 30% in Cumbria (Barne et al. 1996). A notable feature is the high proportion of mid to upper marsh communities remaining, whereas elsewhere in Britain many have been drained and improved for agriculture or developed. Most of the region's saltmarsh is concentrated in the Dee and Ribble Estuaries, Morecambe Bay and inner Solway Firth and much of it is grazed, grassy turf.

The plant species diversity is lower than that for south-east England. The typical zonation (Barne et al. 1996) is from a *Spartina* pioneer community to a low-mid marsh with a *Puccinellia maritima* community and red fescue saltmarsh (*Juncetum gerardii* community) in the mid to upper marsh. Sea purslane *Halimione portulacoides* saltmarsh is less widespread than on the south and east coasts due to grazing. The driftline is dominated by either couch *Elymus repens* or sea couch *Elymus pycnanthus* saltmarsh. At the Solway there is a change from western to Scottish-type marshes, with upper marsh transitions to grassland and brackish fen, and several species reach or approach the northern limits of their west coast distributions here: sea purslane, sea wormwood *Artemisia maritima*, sea lavender *Limonium vulgaris*, lax-flowered sea lavender *Limonium humile*, sea couch, hard-grass *Parapholis strigosa* and strawberry clover *Trifolium fragiferum*.

Invasion by cord grass *Spartina anglica* has caused concerns at some sites in the UK. This species (a polyploid hybrid arising in southern England) has been planted in the UK to make use of its success in binding sediment and thereby stabilising it. It naturally appeared on the Isle of Man in the mid-1970s but appears to have spread little (Allen 1984 in Barne et al. 1996) and was not found at Langness in 2004-2005 (Spencer, pers. comm.). This occurs at only a few sites possibly because the saltmarshes overlie rock with relatively little mud (Barne et al. 1996).

Manx saltmarshes are not considered to be of international importance because they are extremely small and have relatively low diversity. However, they are of importance to wildlife conservation in the Manx context, as they are highly restricted, form a distinctive habitat with a very different biota to other habitats on the island, they could be lost very easily through a change in land use, whether drainage, development or coastal protection works, or as a result of physical changes in the environment, such as a rise in sea level squeezing the marsh against harder inland boundaries which can prevent its inland advance and restrict its continued effect of coastal protection and land claim.



Sea milkwort *Glaux maritima*, growing on saltmarsh at Fort Island, Langness ASSI

Minimum selection criteria for saltmarsh

Saltmarsh quality should be assessed by size, species richness and the presence of rare species. When comparing sites, those showing the full range of saltmarsh communities or a transition or mosaic between saltmarsh and other habitats, such as woodland or grassland, should be considered a priority for conservation. Grazing can modify the plant communities, so management must be considered in comparisons.

1. Include all areas of significant size, i.e. above 0.1 ha (total area to include separate patches on sites with discontinuous patches of saltmarsh);
2. Include smaller sites with populations of rare species (eg. plants found at three sites or less on the island) or community types not represented in larger sites. Every saltmarsh sub-community should be represented, preferably by the best example.
3. Include smaller sites where they form a transition to or mosaic with other eligible habitats or sites adjacent to eligible habitats.

Boundaries

Saltmarshes form a habitat which lies on the boundary of the sea and the land. Some constituent communities are essentially marine or sediment pioneers, whereas others are really terrestrial. These constituent parts shift with the sediments on which they depend, which may move seaward or landward depending on the local erosion or accretion of sediments. Site boundaries should reflect this by using a variable lower boundary such as the astronomical low tide line and an upper boundary high enough to take in all of the present saltmarsh habitat and allow for an expected rise in sea level.

4.6.5 Brackish pools

Description and distribution

Whilst some ponds which lie in close proximity to the coast may have a slightly brackish ecology (due to sea spray periodically reaching the water), true brackish pools are extremely rare on the Isle of Man. They represent the Island's only representative of an internationally rare habitat, **saline lagoons**. Saline lagoons are essentially bodies, natural or artificial, of salt water partially separated from the adjacent sea. They retain a proportion of their seawater at low tide and may develop as brackish, full saline or hyper-saline (i.e. saltier than sea water) water bodies. The largest lagoon in the UK is in excess of 800 ha (Loch of Stenness) although the rest are much smaller and some may be less than 0.1ha (UK BAP, UK Marine SACs Project, 2005).

Saline lagoons sometimes contain invertebrates rarely found elsewhere, and also provide important habitat for waterfowl, waders and seabirds. The flora and invertebrate fauna present can be divided into three main components: those that are essentially freshwater in origin, those that are marine/brackish species and those that are more specialist lagoonal species. On the Isle of Man, species typical of saline lagoons may colonise brackish pools, forming small populations of plants and animals that are rare or absent from anywhere else on the Island. As with larger saline lagoons, the species composition of brackish pools depends a great deal on the type of saline input and the stability of the environment. On the Island, brackish pools take several forms:

- Depressions in dune systems may permanently or seasonally fill with brackish water, forming **dune slacks** – for details of these and related dune habitats, see Section 4.6.3, above.
- Small, isolated, brackish pools notably occur amongst the limestone and volcanic rock formations at Scarlett, on the Island's south coast. They receive salt water from spray and high tide events, resulting in variable salinity depending on levels of salt and rainwater input, and rates of evaporation. They represent some of the most base-rich water bodies on the Island, with the closest ecology to saline lagoons in the UK. They support a characteristic charophyte, or stonewort, flora along with macro-algae such as

Enteromorpha sp., and pondweeds typical of eutrophic, relatively base rich water, such as fennel pondweed *Potamogeton pectinatus*, an uncommon, protected species on the Island.

- Small brackish “trapped” rockpools also occur at places such as Peel Castle old quarry, where a natural or man-made “sill” of rock traps a mix of sea- and rainwater, allowing only periodic salt water inputs at high tide and during storms. The salinity of such pools may vary sharply, with evaporated pools in hot, sunny weather reaching hypersaline concentrations. Such environments are very harsh, and only highly specialised fish and invertebrates are known to survive their demanding fluctuations in conditions.
- Cliff-top ponds may receive small inputs of sea water during storm event, and often remain slightly brackish in nature throughout the year, at least following inputs of high sea-spray and low rainfall. They are less likely to support specialised rare species than true saline lagoons, but may still support unusual plants such as brackish water-crowfoot *Ranunculus baudotii*.

In each of the above cases, the level of salinity in the pool will be heavily influenced by rainfall levels; even specially-adapted flora and fauna may struggle to survive during extended periods of high evaporation and low rainfall.

Classification of brackish pools

Brackish pools on the Island are too small and fragmented to match the usual vegetation community classifications for saline lagoons, but they may show clear characteristics of open water NVC types such as A11 – *Potamogeton pectinatus*-*Myriophyllum spicatum* (fennel pondweed and spiked water-milfoil) community and A21 – *Ranunculus baudotii* (brackish water-crowfoot) community. Saltmarsh plants such as saltmarsh rush *Juncus gerardii* and sea arrow-grass *Triglochin maritimum* may also occur in and around shallow brackish pools.

Status of brackish pools

True brackish lagoons are only known to occur at two places on the Island, at Scarlett Point and behind Peel Castle. They amount to scarcely 0.05ha – too small an area to be covered by the DAFF Phase 1 Habitat Survey. Only those at Scarlett support a diverse vegetation community; the pool at Peel Castle is notable for the presence of sticklebacks, which are generally able to survive brackish conditions. Other instances of small brackish pools - such as regularly “trapped” rock-pools and ponds near the edge of cliff-tops - are thought to amount to less than 1ha in total, comprising many small examples around the rockier parts of the Island’s coastline. Keehan (1999) states that “*all saline lagoons (as defined by Barne J.H. et al, 1996)*” should be considered for selection as Manx Wildlife Sites.

In the European context, coastal lagoons of any size are a very rare habitat type, associated with rare plant and invertebrate species. For this reason Coastal lagoons are a Priority Annex 1 Habitat under the EU Habitats Directive, and Saline lagoons are a UK Biodiversity Action Plan Priority Habitat. Most of the saline lagoons covered by SAC or SSSI designations are greater than 1ha; however, some of the same ecological issues (and rarities) are also associated with smaller brackish pools.

Unless they occur in very inaccessible locations, brackish pools are vulnerable to a range of threats due to human activity. Their vulnerability is heightened by the fact that their very nature does not allow for regular flushing and replacement of water; thus, any water-borne contaminants may become dangerously concentrated as water evaporates between inputs of rainfall or sea-water, threatening the survival of any plants and animals present. This is further complicated by the tendency of lagoons to accumulate very fine bottom sediments. Certain introduced contaminants, such as phosphorus and organic material, will bind to, or be stored within, fine sediments such as mud more than other substrata such as sand or rock, potentially heightening both the severity and longevity of pollution incidents (UK Marine SACs project, 2005).

Other potential problems for brackish pool species include increased turbidity, which can stir up fine sediments and choke or inhibit delicate species, and oil pollution and any associated dispersant chemicals, both of which can jeopardize brackish water invertebrates by affecting the oxygen availability and surface tension properties of the pool. In all these cases, it may be difficult for brackish pools to recover quickly from contamination, due the factors mentioned above. The precautionary approach to impact

assessment is therefore thought to be especially important where water quality in brackish pools is thought likely to be under threat (UK Marine SACs project, 2005).

Minimum criteria for selection

- Boundaries for wider coastal ASSIs should be drawn so as to include any brackish pools along with at least 5m of buffer zone around each pool (or its known widest extent, if fluctuating), along with the immediate sources of both salt and freshwater input – e.g. adjacent beaches, on the seaward side, and freshwater creeks and/or springlines on the landward side.
- All examples of permanent brackish pools, including sites where the pools fluctuate in size either seasonally or occasionally, should be considered for ASSI selection where they fulfill at least one of the Priority Site Criteria listed in Part 3; pools will normally be included within larger coastal sites, but where this is not appropriate it is recommended that brackish pool ASSIs should cover at least 0.25ha including the pools, even where vegetation around the pools is of low species diversity;

particular priority should be given to sites which support any of the following:

- native fish populations;
- specialist brackish water invertebrates;
- plants which are not known to occur at more than three sites on the Island



Stonewort *Chara vulgaris* growing in brackish pools on the Island's south coast

4.6.6 Sea cliffs, caves and slopes

Description and Distribution

The interface between land and sea is a harsh environment for both plants and animals: maritime cliffs, caves and slopes are exposed to the full might of the sea and the weather, and present many challenges as a habitat. The biggest challenge for most plants is simply that of getting a foothold, as hard cliffs may have few suitable crevices in which to take root, whereas soft cliffs may crumble so quickly that few plants can grow to maturity before the substrate shifts and collapses against the oncoming tide and weather. Other challenges for plants and animals include finding enough food and water, coping with salinity, and surviving storm events.

Despite the challenges, there are advantages for species which can make a home on the most exposed parts of the coast: frosts are rare so close to the sea, there is often little competition from other species, and the range of potential predators is also relatively limited. Thus, seabird colonies thrive on steep and rugged cliffs, and some of the rarest and most competition-sensitive native plants in the British Isles are restricted to cliff tops and crevices (Ratcliffe, 1977). The lack of grazing by large herbivores is also an advantage to cliff-top plants: for example, rare, relict, native oak woodland on the Island is thought to be restricted to inaccessible locations, away from livestock and direct human influence (Garrad, 1972).

Coastal slopes

The softer and more weathered coastline around the Island supports **coastal slope** vegetation, which benefits from the accumulation of soil-forming material on the relatively gentle gradient of the slope compared with cliffs. Heavy inputs of salt from sea-spray, coupled with guano deposits from sea-birds, can lead to relatively base- and nutrient-rich soils on coastal slopes. Ruderal vegetation often results, which may grow up into coastal scrub if left undisturbed. At its most well-developed, coastal scrub may support a wide range of plants, invertebrates and breeding birds see **Chapter 4.1: Woodland and scrub** for a more detailed treatment of this habitat. The mild temperatures and high humidity next to the sea can result in woodland ground flora species thriving on coastal slopes even where there is no tree or shrub cover – notably bluebells *Hyacinthoides non-scriptus* and common dog's-violet *Viola riviniana*, both of which thrive on coastal "brooghs" either amongst bracken cover or rough grassland. Such areas are important for the conservation of Dark green fritillary butterflies, which rely on violets as larval food-plants.

Where coastal slopes are grazed, the resulting grassland is likely to resemble sub-maritime, coastal grassland as described in **Chapter 4.2: Lowland grasslands**, with species typical of both neutral grassland and cliff-tops. A colourful, short, herb-rich turf can result which is of value to invertebrates, particularly on south-facing slopes which offer warm, sunny habitats for moths, butterflies and other flying insects.

Cliffs

The Phase 1 habitat definition of hard and soft maritime cliffs is that they have less than 10% vegetation cover. As such, they are both essentially bare rock habitats, where fauna is at least as important a point of scientific interest as flora. In most other respects, however, hard and soft coastlines vary sharply in the ecosystems they support. The question of whether a stretch of cliff has a predominantly hard or soft substrate is therefore a key point to consider at the start of an assessment.

Soft coastlines change readily in response to physical processes, forming beaches, dunes, soft cliffs or unstable coastal slopes, and hence a mosaic of interrelated and interdependent habitats. **Soft cliffs** on the Island generally consist of muddy and sandy sedimentary rocks, and are subject to continual erosion, particularly on the northwest of the coast, where the effects of the prevailing wind and currents are strongest. Lamb (1999) identified several parts of the coast which were eroding, in some cases quite rapidly, notably at Kirk Michael, Glen Wyllin and Ballaugh Shore (The Cronk). However, he also notes that "*the eroding cliffs are of British if not European importance for their geomorphological exposures and are also important for a number of rare and scarce invertebrates (see Boyce and Fowles, 1989, and JNCC 1996).*" For coastal plants and animals, the instability of a soft coastline is partly offset by its ability to absorb some of the stresses of wind and wave pressure and provide an easy substrate for burrowing and taking root. Hard coasts tend to have a more stable substrate, with simple zones of animal and plant communities dictated largely by proximity to the sea.

Hard cliffs around the Island consist of Manx slate, with localised stretches of red sandstone cliff (at Peel) and limestone (at Castletown). None of these is as hard as the hardest type of cliff found in the British Isles, and nearly all have some cracks, fissures and broken up surfaces which may support plant as well as animal life. Most cliffs around the Island are less than 20m in height; however, some hard Manx cliffs are amongst the taller cliffs in the British Isles, reaching over 300m in south west of the Island - for comparison, the tallest UK cliff faces are 430m (granite, St Kilda), with the tall, Old Red Sandstone cliffs at Foula and Orkney reaching around 370m and 335m respectively (Ratcliffe, 1977). Collapse of hard cliffs occurs less often than collapse of soft cliffs, but can be quite catastrophic and sudden, resulting in abrupt loss of marginal habitat. Examples of this may be seen along the far south-west coast of the Island.

Hard cliffs around the Island support some of the Island's most significant seabird colonies, including guillemot, razorbill, black guillemot, fulmar and puffin. These are notably concentrated in the south west and north east of the Island, but may occur at scattered hard cliff sites elsewhere. Raven, peregrine and chough also benefit from hard cliffs (including suitable caves and disused mine adits) both for safe, inaccessible breeding sites and for hunting territory. Near-vertical cliff faces offer an undisturbed, ungrazed refuge for plants as well as animals. Whilst birds are a visible and much-recorded cliff-dwelling group, the inaccessibility of cliffs means that other groups are less well known; for example, a full lichen survey has not been carried out on Manx cliffs, and non-flying invertebrates are under-recorded.

Caves

Unlike most soft cliffs, hard cliffs may offer an additional element of habitat: caves. Whilst the surging tidal currents in sea-caves scour the rocks and make for a harsh and inhospitable environment, caves can also offer an element of stability in an otherwise turbulent coastline. For example, the roof space of large sea caves may offer an unusually mild and frost-free, sheltered area for roosting birds or bats, or for sensitive ferns such as maidenhair fern *Adiantum capillis-veneris*. The pebble beaches at the back of some caves also offer good pupping sites for grey seal *Halichoerus grypus*, as they are sheltered, easily-guarded, and relatively safe from predators.

Whilst coastal slopes can often be surveyed on foot with care, there is great difficulty involved in surveying many cliff habitats and caves, some of which are simply too dangerous to explore without specialised equipment. This is especially the case where they lie underwater or within the intertidal zone (see also: Section 4.6.1: Intertidal habitats). Cliffs and caves, and particularly their marine fauna, are consequently an under-recorded habitat on the Island.



The chough *Pyrhocorax pyrrhocorax*, a familiar coastal protected species which breeds on inaccessible sea cliffs, has been immortalised on the Island's stamps and a coin

Coastal flushes and springs

Freshwater elements such as streams, flushes and seepage lines on and adjacent to cliff faces may not be large enough to qualify for ASSI selection on their own, but are a valuable element of habitat diversity on cliffs, often with associated plant and invertebrate communities. Tufa forms on some of the Island's western cliffs where freshwater flushes and seepages deposit a lime-rich crust on cliffs. This is colonised by mosses and small plants such as brookweed *Samolus valerandi*. Other flushed cliff faces support colonies of royal fern *Osmunda regalis* and hemp agrimony, *Eupatorium cannabinum*. Locally, small reed beds may become established on the soft cliffs, where the cliff vegetation merges with wet coastal grassland. These offer a sheltered habitat for birds such as sedge warblers, which have a limited range of available habitats on the Island.

Classification of sea cliffs, caves and slopes

Vegetation on sea cliffs and slopes may be classified broadly as outlined in **Table 4.6.e**.

Table 4.6.e: Broad categories of maritime cliff vegetation on the Isle of Man

Type and Description	Species rich?	Closest NVC categories (See Rodwell, 1992 ⁵)	Rarities present?	Protected species present?
Hard cliff rock crevices	Not usually, but may support rarities (see right); some communities of annual plants (e.g. MC5) are relatively species-rich	MC1 – <i>Crithmum maritimum</i> - <i>Spergularia rupicola</i> (rock samphire/ rock sea-spurrey) maritime rock-crevice community No Island NVC sample MC5 – <i>Armeria maritima</i> - <i>Cerastium diffusum</i> ssp <i>diffusum</i> (thrift/ sea mouse-ear) maritime therophyte (annuals) community No Island NVC sample	Provides good bird cover, so may attract nesting or passing rarities; some rare and specialised invertebrates may occur (currently an under-recorded group), including the Pod lover moth <i>Hadena perplexa capsophila</i> , whose larvae feed on sea and bladder campions.	Rock sea lavender <i>Limonium binervosum</i> agg. may occur; Schedule 1 birds may nest and/or feed in hard cliff crevices, including chough <i>Pyrhocorax pyrrhocorax</i> , raven and peregrine; Schedule 5 animals may occur, including grey moths, whose larvae which feed on sea campion.
Sea bird colonies These areas of cliff attract numbers of breeding sea-birds, which affect the vegetation and invertebrate communities in turn; e.g. colonies of burrowers such as puffin and Manx shearwater are associated with NVC type MC7, whereas steep cliffs of breeding guillemots and gulls are commonly associated with type MC6.	No, but may attract rare birds and/or specialised invertebrates	MC6 – <i>Atriplex prostrata</i> - <i>Beta vulgaris</i> ssp <i>maritima</i> (common orache/sea beet) sea-bird cliff community One Island NVC sample MC7 <i>Stellaria media</i> - <i>Rumex acetosa</i> (chickweed/sorrel) sea-bird cliff community Not mapped in NVC (too scattered).	Rare breeding birds may occur (see right)	Schedule 1 protected birds including chough, peregrine, Manx shearwater and shag – also, see below sand martins on soft cliffs).
Soft cliffs The upper landward edge of soft cliffs may slump to form friable, shifting coastal slopes, with a distinct “undercliff”, usually vertical, on the seaward side. Undercliffs and other vertical, soft cliffs formed when soft coastlines collapse can be valuable for insects such as mining bees, and	No.	Soft coastlines are often too friable and unstable to support an established vegetation cover; however, scattered plants may get a foothold, forming loose communities of species commonly found on either strandlines and shingle or foredunes (see Sections 4.6.2 and 4.6.3, above)	Soft cliffs can support a specialised invertebrate fauna where the conditions are suitable. Sites that are not eroding too quickly, and allow vegetation to form, or have freshwater influence, have the greatest potential. Sand martins favour soft	Sand martins nest here in colonies; protected plants such as oysterplant <i>Mertensia maritima</i> and Viper’s bugloss <i>Echium vulgare</i> may occur at the base of soft, sandy cliffs, especially where they meet the strandline. Lizards may occupy sunny slopes.

may offer suitable nesting cliffs for sand martins. The base of soft cliffs may also support unusual vegetated strandline plants such as oysterplant and purple fumitory.			cliffs for nesting. Scattered rare ephemeral plant species such as purple ramping-fumitory <i>Fumaria purpurea</i> may occur.	
Cliff top vegetation	This kind of vegetation, where it does not fall into the Sea bird colonies category (above), generally forms grassland and/or scrub, and has been treated as Coastal scrub or Coastal grassland; see Chapter 4.1 – Woodland and scrub , and Chapter 4.2 – Lowland grasslands .			



Sea ivory *Ramalina siliquosa* (centre of picture) is well-known, but most coastal rock face lichens are likely to be under-recorded on the Island

Status of sea cliffs, caves and slopes

The Terrestrial Survey of the Coast of the Isle of Man (Lamb, 1999) was carried out in a series of 58 coastal sections of approximately equal length. The survey revealed the rugged nature of the Island's coastline, with hard maritime cliffs a major component of several stretches, mostly around the east and south-west edges of the Island, and a minor component of 22 sections. Soft cliff was less common, recorded in seven sections, as a major component in each case, in the northwest and northeast of the Island. Lamb (1999) also noted that soft cliffs were exclusively a maritime habitat, whilst hard cliffs could be found inland, e.g. in ravines or quarries. The small section of limestone rock in the Castletown area has the rarest cliff and slope types on the Island, but supports a disproportionately high proportion of the Island's cliff and rock crevice plant biodiversity (BDIS). In addition to cliff and slope, 34 sections of the Survey had boulders and/or rocks above the high tide mark, both from soft and hard rock types.

The value of cliffs and slopes for fauna in particular is recognised in the **Manx Wildlife Sites** Handbook (Keehan, 1999), which states that all stretches of maritime soft cliff that are shown to be of high invertebrate value, and all stretches of maritime hard cliff that support important colonies of seabirds, should be considered for selection as Wildlife Sites.

The **UK** supports a significant proportion of EU sea cliff vegetation (www.jncc.gov.uk). *Maritime Cliff and Slopes* is a UK BAP Priority Habitat, and a high proportion of the hard cliff coast in England has been notified as SSSIs - in areas such as the south-west of England almost the whole cliff-lined coast has been notified. Notification of soft cliffs has been less extensive, but areas such as north-west Norfolk and the Isle of Wight have a high proportion of their soft cliffs notified. In Wales approximately half of the total maritime cliff resource has been notified as SSSIs, but as yet only a small proportion has been notified as ASSIs in Northern Ireland (www.Ukbap.org.uk).

In the **European context**, '*Vegetated sea cliffs of the Atlantic and Baltic coasts*' is listed in Annex 1 of the EC Habitats Directive. Under this Directive, nine lengths of coastline in the UK have been nominated as candidate Special Areas of Conservation (SAC) for their cliff features (two of which include substantial representation of soft cliffs). The importance of cliffs for seabirds has also been recognised: under the EC Birds Directive, 38 Special Protection Areas (SPA) in the UK have been designated which include cliff sites - these comprise 30 sites in Scotland, 5 in Wales, 2 in England, and 1 in Northern Ireland.

Whilst cliffs often provide habitats which are unusually free from direct human influence, cliff and slope communities still face a number of **threats**, including:

- *Erosion.* High rates of erosion do not imply a loss of the cliff resource, either in geological or biological terms. Cliff face communities are able to retreat with the cliff line, and erosion is vital for constantly renewing geological exposures and recycling the botanical succession on soft cliffs. However, cliff-top vegetation may be destroyed where it is squeezed between a receding cliff face and cultivated land (UK BAP) and the rate of erosion of a soft cliff can affect the ability of a diverse invertebrate fauna to form.
- *Built development.* Built development near the edge of cliffs can prevent cliff-top biological communities from retreating in response to cliff erosion, subjecting them to a form of 'coastal squeeze'. Developments may also result in decreased habitat naturalness and representativeness, either from the need for enhanced coastal protection, or from increased littering and introduction of new species (see below), e.g. from newly adjacent gardens.
- *Coastal protection.* Coastal protection systems are still relatively limited on the Island, but may play a stronger part in coastal defences in the future. Attempts to protect property from the effects of erosion can result in cliff faces being re-profiled and/or sown with hardy grasses of little value for nature conservation. All such works have the effect of stabilising the cliff face, resulting in geological exposures being obscured, bare soil and early pioneer stages becoming overgrown, and wet flushes drying out.
- *Agriculture.* In traditional low-intensity grazing systems, livestock are grazed on cliffs and slopes, maintaining the open maritime grassland vegetation. Intensification of agriculture can lead to maritime grassland on more level terrain being ploughed out, while that on sloping ground may be abandoned and, where not maintained by exposure, is frequently overgrown by scrub. Localised eutrophication can be caused by fertiliser run-off from arable land above and this encourages coarse, vigorous 'weed' species at

the expense of the maritime species. Agricultural land drains discharging on the cliff face may cause local acceleration of erosion.

- *Introduced species.* Predators, such as cats and rats, can have a significant impact on populations of cliff or burrow nesting seabirds, particularly on isolated sites such as the Calf and small islets. Other introductions include feral goats and stray sheep; these are not thought to cause much damage to coastal vegetation communities in general, but their presence as nimble grazers could affect the survival success of rare cliff crevice plants.

The spread of certain alien, invasive plants, especially members of the flowering plant family Aizoaceae such as the hottentot fig *Carpobrotus edulis*, can have a devastating impact on indigenous maritime plant communities (UK BAP). This species has yet to colonize Manx coastlines as it has in Cornwall and elsewhere in the British Isles, but the Atlas 2000 (BSBI, 2000) shows it to have been recorded in five of the 10km squares that cover the coastline. Its emergence as an invasive species should not therefore be ruled out as a future problem. Other species which may invade cliff and slope vegetation and displace native species *en masse* include Montbretia *Crocasmia x crocosmiiflora*, cotoneaster *Cotoneaster spp.*, Spanish bluebell *Hyacinthoides hispanicus* (and its hybrid with the native bluebell) and Japanese knotweed *Fallopia japonica* (which prefers disturbed areas).

Minimum criteria for selection

Selection on species grounds

The importance of maritime cliffs for breeding birds means that some cliff sites are likely to fit the criteria for selection on bird criteria alone - see **Chapter 4.9: Fauna**. As a priority, the best site for any species of conservation concern should be put forward for designation.

Selection on habitat grounds

- The best examples of uninterrupted soft cliff and slope coastline should each be selected from the northwest and northeast Manx coastlines; sites should fulfil at least one of the Priority Site Criteria listed in Part 3, and priority should be given to the sites with the most diverse ecology, in terms of vegetation cover, adjacent habitat types (including intertidal habitats), and fauna.
- Stretches of hard maritime cliff will be considered for selection where they are known to fulfil at least one of the Priority Site Criteria as listed in Part 3. As a minimum, site boundaries should include the shoreline, cliff, and any semi-natural cliff-top vegetation.
- With all cliff, slope and cave assessments, priority should be given to sites which are known to support rare plants and/or animals, in particular those which depend exclusively on highly undisturbed and inaccessible habitats, including at least one example of each of the following:
 - core breeding sites for dark green fritillary butterfly;
 - good examples of native woodland and scrub on cliffs and slopes;
 - key pupping sites for grey and/or common seal;
 - cliff crevice communities with good plant diversity, especially rarities;
 - cliff and cave communities with good plant diversity, especially ferns;
 - known bat roosting caves;
 - core breeding sites for Schedule 1 protected birds
 - sites on limestone rock, which is the rarest cliff type on the Island, and which supports a high proportion of the Island's cliff and rock crevice plant biodiversity (BDIS);
- Freshwater elements such as streams, flushes and seepage lines on and adjacent to cliff faces should be included within designation boundaries as a valuable element of habitat diversity.
- In all cases, where there is a natural transition between maritime communities and inland grass or heath, the site boundary should be drawn to encompass both interests where practicable.

N.B.: maritime cliffs, caves and slopes are amongst the least well-recorded habitats on the Island, but this should not be allowed to prejudice their selection as ASSIs; where there is a dearth of biological information the precautionary principle should be applied to site selection, with appropriate surveys made a priority pending boundary confirmation and management planning.

4.7 Uplands

Introduction

This chapter concerns the selection of ASSIs which lie wholly or partly above the usual altitude at which fields are enclosed and cultivated. Throughout the British Isles the open, unenclosed nature of upland vegetation is distinctive, a familiar landscape resulting from necessarily extensive farming practices. The resulting mosaic of habitats is widespread in the highest and most exposed parts of the UK, Ireland and the Isle of Man, and in each case represents the largest single category of semi-natural habitat. Paradoxically, this terrain supports some of the rarest and most vulnerable habitats in the world, namely blanket bog and temperate heather moorland – both of which are protected under the EU Habitats Directive. Countries within the British Isles are therefore considered to have a special responsibility for the conservation of these habitats.

Whilst onlookers usually have little difficulty in identifying an upland landscape by appearance, there is no established scientific definition of an upland as opposed to lowland habitat, and deciding on appropriate boundaries for upland ASSIs is notoriously difficult (Ratcliffe, 1977). It has been noted elsewhere that a simple altitude “cut-off point” is not always the best way of defining uplands, and that the distinction between upland and lowland is an arbitrary one (NCC 1989). This is partly due to the varying degrees of latitude and coastal exposure in the British Isles, with some northern sites exhibiting the same vegetation communities at much lower altitudes than their southern counterparts, particularly on exposed coastland. This gradient of variation is recognised as an important feature of habitats in the Atlantic biogeographic region (which includes all of the British Isles), particularly where typically upland and lowland vegetation both lie on a continuous ecological gradient.

Manx uplands occupy an unusual position, being more or less at the northern limit of typical English lowland heathland, and the southern limit of the type of wet, bryophyte-rich moorland commonly found in north-west England and western Scotland (see Rodwell *et al.*). In most cases, the transition from lowland to upland vegetation on the Island occurs at around 250m, i.e. lower than the 300m which is commonly used to define the same threshold in England. Whilst this is a useful starting guide for the identification of potential upland ASSIs, it will not be taken as an absolute cut-off point, as many upland habitats grade into adjacent habitats at lower altitudes, and the resultant gradient of ecological characteristics may be an important feature of the site.

Upland habitats on the Isle of Man are almost entirely semi-natural in origin, and contribute greatly to the wild appearance of the Manx landscape, which is often praised for its attractiveness. The uplands are not, however, wilderness in any sense, having received centuries of settlement, mining, quarrying and farming which have shaped the nature and extent of the habitats present. These include a range of mires, acid grasslands, heaths and streams. Native woodland is restricted to natural colonisation of sheltered river valleys and rills, and the more sheltered hillsides where gorse and bracken have been left ungrazed and native scrub has started to colonise it. Elsewhere, the tree cover is limited to plantations, mostly of conifers but with a more recent emphasis on mixed and broadleaved planting. It should be noted that all of the uplands receive some degree of human influence, either from grazing or burning (deliberate or not), walking, forestry, shooting or in some cases direct cultivation. There has also been widespread agricultural improvement, particularly on the lower fringes of the uplands which can provide productive fields if drained and limed.

Description and distribution

European uplands are commonly defined as being either montane or sub-montane. Montane habitats occur at high altitudes, above the natural tree-line. They support low-growing communities of specialised plants and invertebrates which are strongly frost-hardy. Whilst the altitude at which montane vegetation occurs can vary with latitude and degree of exposure, in the central British Isles it is commonly defined as starting at 600m. This altitude only occurs in one place on the Isle of Man, the summit of Snaefell, and the montane limit is borne out by the presence of a tiny community of arctic/alpine vegetation – notably least willow *Salix herbacea* – on the summit. Aside from this, upland habitats on the Island can be said to fall clearly within the sub-montane category.

The fact that this is varied by the presence of small pockets of arctic/alpine plants is of interest in itself, offering the potential for a biological indicator of whether the climate on the Island is changing.

A topographical map of the Isle of Man shows three broad areas of sub-montane upland. The highest land is often referred to as the Northern Uplands. It consists of several high hills including Snaefell at 620m, and a long, hilly ridge culminating in North Barrule at 565m. There is also a Central Uplands, which is linked to the Northern Uplands by a series of low hills. The Southern Uplands are the lowest in altitude, centering upon South Barrule (483m) and extending to the south-west coastline of the Island. They are divided from the central and northern hills by the Central Valley. Other high ground around the Island occurs in association with high cliffs, notably at Maughold Head and Spanish Head. However, neither of these clifftop areas rises to much over 100m, or exhibits typical upland vegetation.

Manx uplands contain most of the biological features which are identified in the UK *Guidelines for Selection of Biological SSSIs* as being of special importance. These being:

Acidophilous dwarf shrub heath and scrub

Acidic and calcicolous grassland

Fern- and bryophyte-rich vegetation

Montane plant communities generally

Blanket bog

Phytogeographical interest

Ornithological features

Other faunal features eg relict assemblages of invertebrates (NCC, 1989)

Of the above, all the habitats mentioned occur within the Manx uplands except extensive montane vegetation and exceptionally species-rich fern and bryophyte communities – although limited elements of both do occur. All the non-habitat-specific features are also known to occur:

- **“Phytogeographical interest”** is a frequent, and important, feature of many uplands in the Atlantic fringes of the British Isles, and refers to the unusual combination of species representing different geographical elements of flora – for example, Southern Atlantic, Mediterranean and Northern Continental – within the same area (Matthews, 1955 *per* NCC, 1989). An example on the Island might be the proximity of hare’s-tail cotton-grass *Eriophorum vaginatum* (a boreo-arctic montane species) to cross-leaved heath *Erica tetralix* (a predominantly suboceanic temperate species) and devil’s-bit scabious *Succisa pratensis* (a Eurosiberian or North Continental temperate species), all of which commonly occur together in the more species-rich wet heaths on the Manx upland fringes (Preston *et al* 2002).

- **Ornithological features.** The Manx uplands are of importance for birds. Cullen and Williams (*per* MNCT, 1997) identified the importance of the Manx uplands for those upland birds which benefit from open, heathy habitats – notably the hen harrier *Circus cyaneus*. Despite the relative scarcity of woodland and wetland habitats in the Manx uplands, a wide range of typical moorland breeding species has been recorded, including meadow pipit *Anthus pratensis*, wren *Troglodytes troglodytes*, stonechat *Saxicola torquata*, wheatear *Oenanthe oenanthe*, lapwing *Vanellus vanellus*, curlew *Numenius arquata*, short-eared owl *Asio flammeus*, skylark *Alauda arvensis*, raven *Corvus corax*, red grouse *Lagopus lagopus scoticus*, linnet *Carduelis cannabina*, redstart *Phoenicurus phoenicurus* and twite *Carduelis flavirostris* although the last two species have not been recorded breeding for many years. Merlin *Falco columbarius*, peregrine *Falco peregrinus* and reed bunting *Emberiza schoeniclus* are also recorded as rare breeders in this habitat. Upland grassland, including agriculturally improved areas, additionally provides valuable winter feeding areas for the Island’s important chough *Pyrrhocorax pyrrhocorax* population, and has the potential to support breeding corncrake *Crex crex*. Cullen and Williams (*per* MNCT, 1997) identify several conservation priorities amongst upland birds: hen harrier, corncrake, merlin, skylark, linnet, reed bunting, peregrine, chough and red grouse. Most of these are known to be of conservation concern either globally (in the case of corncrake) or within the context of the British Isles.

- The presence of **relict faunal features**, in this case invertebrates, is likely but has yet to be fully investigated, the one major study of Manx invertebrates having missed out the uplands (Boyce & Fowles, 1989). However, Luff (*per* MNCT, 1997) identified “at least 76 species” as possible upland ground beetles, of which 11 occur only, or typically, in the uplands. Luff also notes that this is over half the 19

upland species known to occur in northern England. It might be implied from this that the Manx uplands merit further study of this and other invertebrate groups.

Much of the background information on habitats such as mires and watercourses has been detailed elsewhere in the Criteria. The following chapters in particular are of relevance:

4.3 Lowland heathland

4.4 Bogs, fens and flushes

4.5 Open water and swamp

All of the above habitats may occur next to, and in an ecological gradient with, upland habitats. In such cases it is desirable to protect the range of the semi-natural habitat rather than impose an artificial limit on the altitudinal boundaries of the site see (**Recommendations**, at the end of this chapter). The chapters on **Woodland and scrub** and **Lowland grassland** may also have a bearing on upland habitats. Sections **4.7.1 – 4.7.3**, below, therefore concern those elements of Manx habitat which occur solely, or most typically, within the uplands.

Assessing uplands

The UK Guidelines for assessment of complex upland areas

The problem of how to select protected sites from extensive areas of semi-natural habitat with no obvious boundaries has been identified in the UK Guidelines for Selection of Biological ASSIs:

"Uplands tend to be the most difficult cases, with blanket bogs the most problematical of all, because they usually lack conspicuous edge features along which to draw boundaries".
(NCC 1989 p35)

This difficulty applies as much in the Isle of Man as elsewhere in the British Isles. The solution in the UK was to apply a set of guidelines to aid decision-making in areas with poorly-defined boundaries. Subject to considerations of scale within the smaller island context, these guidelines are also applicable to the Manx ASSI selection criteria. They are as follows:

"There are five simple guidelines which help to reach decisions on these complex and extensive habitats, though they also apply more generally to site selection.

- *At least one example must be included of every biotic feature (except recent artefacts) present - habitat and vegetation types and species of plants and animals. While special consideration should be given to inclusion of rare habitats and species, the common and widespread need to be represented in proportion to their relative extent and abundance. The total animal community "pyramid" should be represented, from "top" predators to the bottom of the food chain.*
- *The full complement of abiotic features should also be included, over as complete as possible a range of the variation present in each (e.g. in underlying geology, soil type, slope, aspect, elevation, macro- and micro-topography and hydrological regime). The amount of each site component to be included should be sufficient to ensure the viability of that component or of any other component which is functionally dependent on it. There is usually a correspondence between biotic and abiotic diversity, but this is not always fully understood and sometimes the abiotic features are more readily identified and mapped.*
- *Where the conservation of an important habitat or species depends on the maintenance of a functional relationship with some peripheral habitat or environmental condition, sufficient of the latter must be included to satisfy the needs of the former. Examples are hydrology of adjoining areas and feeding range for some animals.*
- *Juxtapositions between site components should be treated as further components in their own right, in so far as they reflect varied biotic and abiotic interactions. For instance, where plant community X adjoins communities Y and Z at one place but adjoins communities A and B at another place, the two groups XYZ and XAB should be treated as separate entities and both included within the site boundary.*
- *The total amount and distribution of the ecosystem concerned, as the sum of all the SSSIs representing that particular type, must be sufficient to ensure viability of all component communities and species in case all remaining areas of such habitat outside these SSSIs become subjected to land-use change. This is a matter not only of size but of spacing between sites, to maximise the probability of exchange of mobile species between them. They must also be sufficient to satisfy the national and international conservation needs..."*
(NCC 1989 p35)

Achieving a similar level of protection in the Isle of Man

In 2002, DAFF commissioned research into the selection of protected upland areas, based on computer modelling of the biodiversity resource in the Manx uplands as a whole (Balson, 2002). The objective was to help achieve the aims listed above by providing a scientifically sound basis for the selection of large, upland ASSIs. The research utilised **Marxan**, a piece of software designed to produce theoretical series of sites, dependent on the value attached to key biological features. A large amount of the most up-to-date biological data available was entered into the program, including plant and bird species distribution, protected species locations, data from the DAFF Hill Land Survey and habitat distributions from the DAFF Phase 1 and 2 Habitat Surveys. Elements of the data were assigned a value according to conservation importance – for example, RSPB “red list” birds were given a higher data value than “green list” species, and internationally important habitats and species were also prioritised.

The Marxan program uses optimisation algorithms to give an indication of the “hotspots” that would need to be included in an upland ASSI series if all the biological features mentioned in the guidance (as quoted above) were to be protected. To maximise the usefulness of the project, all hilly areas – including those Southern Upland areas which fall below 250m – were included.

Results of computer modelling of upland “biodiversity hotspots”

The computer modelling approach was partly aimed at providing quantitative confirmation of earlier recommendations made prior to the development of a detailed biological database – in particular, the key uplands which were identified in the report *Nature Conservation on the Isle of Man* (Pritchard 1975). These were:

- Southern Uplands - includes South Barrule and Stony Mountain
- Slieu Ruy Uplands - includes Slieau Ruy heath, grassland and mire, Colden and the upper reaches of the Blaby River
- Bein-y-Phott Uplands - adjacent to the above, including extensive grassland, mire and heath
- Snaefell Uplands - includes the whole of Snaefell and surrounding grassland, and the only arctic-alpine vegetation on the Island
- North Barrule Uplands - contains the largest proportion of high-altitude land on the Island, and supports extensive wet acid upland pasture with heath elements and strict bog plants such as cranberry *Vaccinium oxycoccus*.

The Marxan program is flexible in terms of data input. Hypothetical reserve networks could thus be generated for a range of parameters according to “maximised coverage”. Maximised coverage applies a pre-set limit on cost and/or area, and aims to maximise the conservation features possible within the constraints. This “working backwards” approach is useful where it is desirable to protect a minimum proportion of a large complex of habitats which contain a broad distribution of many individual features of conservation importance. The drawback to this approach is that some features may have to be “sacrificed” in order to gain the maximum overall advantage within the constraints applied. However, this can be corrected by adding to the network any omitted sites which are known to have just one or two unique features of high importance rather than a high-scoring overall diversity.

Three Marxan “trials” were run, aimed at selecting either 30%, 41% or 50% in total of the 1km² units that make up the total upland area. These percentages were chosen to reflect the high importance of the uplands in terms of total contribution to semi-natural habitat area on the Island, and their international importance. Depending on the total area of sites eventually selected, the mid-range of 41% would be more or less in line with the overall long-term aim of ASSI protection for 10% of the Island (see **Section 1.4.3 What proportion of the land should be designated?**). A similar list of priority areas for upland conservation on the Isle of Man was produced by each trial, varying only in extent, indicating a clear series of “hotspots”. This extends, supports and qualifies the recommendations of ITE 1975. The areas in question are listed in Table 4.7a, and form part of the recommendations of the Balson report:

Table 4.7.a – Manx upland conservation “hotspots”

(Table adapted from Balson, 2002 p37)

1km grid square	Area description
Southern Uplands (including Bradda Coast)	
SC1869, SC1870	West of Bradda West
SC2071, SC2072, SC2073	Eastern portion of Bradda Hill and northeast of Fleshwick
Note: these included the cliff-top coastal heathland of the Bradda Head area. Due to a catastrophic fire in October 2004, much of that area has now lost its important heathland communities. Whilst the vegetation is starting to regenerate at the time of writing (2006), the previous level of heathland diversity may not return for many years, increasing the relative importance of other Southern Upland habitats compared with the original analysis.	
SC2175, SC2176	Portion nearest the coast of the Manx National Heritage site at Eary Cushlin/Glion Moar
SC2376	Glen Rushen, immediately northwest of Glen Rushen plantation
SC2378	Doarlish-Cashen, southwest of Arrasey Plantation
SC2475, SC2476	Immediately south (east and west) of Glen Rushen plantation and northwest of Cringle plantation. Includes the Round Table area.
Central Uplands	
SC3184	South of Earyglass, on the lower north slope of Beary Mountain, near Blaber River
SC3182, SC3282, SC3283, SC3284, SC3381, SC3385	Southern slopes of Beary Mountain, also Slieau Ruy summit and northwest slopes, and Lhargee Ruy's western slopes and south of the plantation at Glion Gill
SC3584	North of Colden Plantation and west of Injebreck Plantation
Northern Uplands	
SC3586, SC3484, SC3587, SC3687, SC3787, SC3886, SC3887, SC3888, SC3889, SC3788, SC3990, SC3991, SC4090, SC4089, SC4088, SC4187, SC4186, SC4286, SC4386	A large block surrounding the Sulby reservoir that stretches directly east from the northeastern slopes of Sartfell (southwestern slopes of Slieu Freoghane) to the lower southwestern slopes of Snaefell. This area also stretches northward, taking in the summit of Slieau Managh and surrounding slopes, but not including the summit of Snaefell or its direct northern or southern slopes. However, the northeast slopes of Snaefell are included in this area, as are the top and northern side of the Laxey Valley.
SC3781	A portion of the southwestern slope of Slieau Ree
SC3982, SC4082, SC4083, SC4084	East of Kate's Cottage, west of Conrhenny plantation, and north and south west of Glenroy
SC4289, SC4389	The valley west from Corran to Cornaa
SC4391	The northwestern slopes of North Barrule (southeast of the Brookdale plantation)
SC4292, SC4192, SC4092	Northeast of the Slieau Managh Plantation, and stretching east to the upper reaches of Glen Auldyn
SC3791, SC3692	The northern half of, and stretching north and west from, the Manx Natural Heritage site at Sulby towards (although not including) Cashtal Lajer (roundhouse).

The above units contain the maximum overall upland biodiversity within a set maximum area of the Island's uplands, according to the values assigned to each species and habitat type (see Balson, 2002 for more details). It may be inferred that, without a representative ASSI from each of these hotspot areas, the full range of Manx upland biodiversity would not be protected by the ASSI system without having to increase the proportion of uplands protected. **To the list should be added any 1km squares for which the overall amount of data is relatively low, but there is known to be a specific and/or unique interest: for example, the arctic-alpine vegetation on the summit of Snaefell.**

The final recommendation of the Balson report relates to the hotspots identified above:

"2 The Island's conservation goals for the uplands would best be met by designating a series of areas corresponding to the "conservation hotspots", opting for the largest sites possible." (Balson, 2002)

Whilst computer modelling does not enable the precise selection of individual site boundaries, it follows that a representative site from each of the unit areas listed above should be included in any series of upland ASSIs which aims to fulfil the same quality of criteria as those used in the UK.

Assessment of upland ASSI quality

All of the points on the Checklist of Priority Sites Criteria listed in **Part 3** are relevant, with the following criteria of especially high importance:

Size

The need to represent topographic, rather than artificially enclosed, units means that most upland areas which are worthy of protection are necessarily large. Where possible, they should include catchments, and all the semi-natural habitats that follow a natural gradient from high to low altitude. In practical terms it may be necessary to draw boundaries along the "hill wall" which separates enclosed from unenclosed land; however, where semi-natural habitats of high conservation importance are contiguous with upland habitats they should be included with upland ASSIs (as is the practice in the UK), as the ecological gradient from higher to lower altitudes represents an important biodiversity resource. It is thus preferable to select a small number of large, complete sites rather than pick out artificially-defined boundaries from parts of larger ecosystems (see NCC, 1989 pp181-188).

It is accepted that the very large designated areas recommended for the UK (upwards of 4,000ha for some upland habitats) are not applicable in the context of the Isle of Man, but the need to consider designating relatively large areas of the most important uplands should still be taken into account when assessing potential ASSIs.

Location

Much of the variation in upland habitats is due to the amount of peat present, and whether it is accumulating, static or diminishing in quantity. This is heavily dependent on the location of the habitats in question, particularly in relation to surface hydrology. Sites where blanket bog is "active", i.e. where peat is actively being formed, are recognised as internationally important. These sites are irreplaceable in terms of their dependence on a particular type of location.

The location of the higher altitude vegetation on the Island is evidently irreplaceable should the current land use change; it is therefore important that the ASSI series covers the summits of important hills. The prospect of climate change makes these important not just for current biodiversity, for which there are no alternative sites, but for the study of changes in the future, as cold-tolerant species at higher altitudes are often sensitive to changing climatic profiles, and may offer an early indication of permanent change.

Diversity

The presence of a full range of successional stages is important for the conservation of species which depend on upland habitats, particularly invertebrates (see English Nature, 2001 p6.11; Ratcliffe, 1977; NCC, 1989). This is related to the importance of structural diversity, which results from a mixed pattern of grazing, burning and unmanaged areas. Thus, areas of unburnt heather are vital for the conservation of slow-growing lower plant groups and overwintering invertebrates, whilst some vascular plants may benefit from the patchy habitats that result from rotational burning. A mixed moorland which exhibits a range of structural stages is therefore preferable to a "monoculture" of burned or very old heather.

Rarity, fragility and vulnerability

The presence of internationally rare Habitats Directive Annex 1 Priority Habitats on the Island's uplands makes them a priority for conservation. In addition, active blanket bog is a rare and locally distributed habitat on the Island in its own right, and vulnerable to a range of threats. ASSI selection should prioritise these habitats, especially those which show a full range of sub-types, successional stages and ecological gradation.

Rare breeding birds and animals, and protected plant species, should also be prioritised, especially those which are solely dependent on the uplands for their continued survival on the Island. The use of the Marxan programme, which utilises a weighted data value approach, has ensured that the conservation "hotspots", listed in Table 4.7a, above, should encompass the majority of rare and vulnerable features in the uplands within a minimum "cost" in terms of area. However, the comment which follows Table 4.7a, regarding the inclusion of any "single interest" sites not included in the table, should be borne in mind when looking at protection of rare upland species.

Ecological position

The boundaries of upland ASSIs should follow natural topographic units wherever possible, especially where wetland is included that depends on catchment hydrology. When selecting watercourses (see **Chapter 4.5 Open water and swamp**) effort should be made to include at least one watercourse from an upland source for its full length, in order to protect the range of ecological variation from upland to lowland. Hill land ASSIs should include contiguous lowland semi-natural habitats wherever possible – particularly heathland, where there may be no clear distinction between upland and lowland vegetation, but a gradually changing community which is dependent on both upland and lowland elements for its continued biodiversity. See **Recommendations**, below.



The Raven *Corvus corax* is a characteristic bird of steep upland and coastal areas

Dominant upland habitat types

Whilst a wide range of minor habitat elements may occur in the uplands, all of which add to their diversity and wildlife value, there are three clearly dominant habitat types: **acid grassland**, **heath**, and **bog**. All three have lowland counterparts, which are detailed in **Chapters 4.2, 4.3 and 4.4** respectively. Those elements which occur solely or typically in the uplands are described below, and summarised in **Table 4.7.b. The minimum criteria for selection of each of these habitat types should be used in conjunction with Table 4.7a, above, in order to select a series of upland ASSIs which collectively represent the full range of upland biodiversity on the Island – see Recommendations, below.**

4.7.1 Acid grassland

Description and Distribution

The origins of acid grassland in the hills lie in the clearance of upland woodland prior to the onset of modern patterns of livestock farming. Regular grazing of cleared uplands suppresses the natural tendency towards growth of dwarf shrub heath, resulting in extensive, open grassland of relatively low species diversity (Drewitt & Manley, 1977). As with lowland examples, drier acid grassland in the uplands may be subject to agricultural improvement such as drainage, liming and re-seeding, whilst wetter and more acid areas may still receive basic drainage and control of unwanted species. In both cases the dominance of grasses is maintained by grazing, as acid grassland on the Island tends towards heathland if left unmanaged, and is prone to encroachment from European gorse and bracken. For these reasons the term “unimproved upland acid grassland” is possibly a misnomer, as upland grassland usually represents an agriculturally improved state on sites that would quickly revert to heather and bilberry cover if farming activities were reduced or removed.

Rainfall in the Manx uplands is heavy and frequent, averaging over 1524mm (60 inches) a year in the hills (compared with under 760mm (30 inches) per year in the south of the Island around Port St Mary) (Birch 1960 *per* Allen, 1984). This can result in peaty “mor humus” topsoil with very low levels of nutrients available to plants, and a pH as low as 3.2 (Fullen, Harris & Kear 1999), i.e. almost as acidic as vinegar and hence inhospitable to all but the most specialised plants. Upland pasture on this kind of soil is frequently interspersed with bog and/or heath, with a full spectrum of degrees of variation in between. A complex, interrelated mix of wet and dry acid grassland, bog, flush, wet and dry heathland and stream valleys is thus strongly characteristic of the Manx uplands. Collectively these make up over half of all the semi-natural habitats on the Island, and support a significant number of protected species.

Despite its lower species diversity compared with some lowland grassland types, upland acid grassland complements other upland habitats, and constitutes a vital part of the ecosystem for some upland birds. For this reason, the assessment process should aim to recognise good examples of patchy upland grassland mosaic within areas of varied habitat, rather than pure acid grassland sward with no heath or wetland elements.

Classification of upland acid grasslands

Upland acid grassland on the Island may be classified broadly as outlined in **Table 4.7.b.**

Upland acid grassland status

Acid grassland occurs throughout most of the Isle of Man, making up around 7% of the Island’s land use total. This equates to over a quarter of all the semi-natural habitat present, and nearly half of all the grassland (Sayle *et al*, 1995). Whilst this includes lowland as well as upland categories, the majority lies within the uplands, often in extensive tracts intermingled with other habitats such as heath and bog.

Acid grassland in the uplands of the British Isles is generally treated as less of a conservation priority than heath or blanket bog, as it is very widespread and often results from the modification of heath and bog by human activities (i.e. livestock management). There are some British types of upland grassland which exhibit considerable species diversity, and are home to a range of rare and delicate plants. However, these habitats are associated with more base-rich soils than are found on the Island, usually over siliceous bedrock substrates and in very high and/or exposed conditions. These upland grasslands of high

conservation priority do not occur in the Isle of Man.

Despite the relative lack of diversity in upland grassland types, some of the Manx uplands support a relatively species-rich pasture due to the mixture of bog and heath plants present, which remain as a minor element throughout due to a light grazing regime. These more diverse areas may support rarities such as mountain pansy *Viola lutea* and ivy-leaved bellflower *Wahlenbergia hederacea*. The very highest grasslands, which fall within montane grassland categories, are not as diverse but represent a vulnerable and restricted habitat, and the only available sites for the Island's small number of arctic-alpine species, notably least willow *Salix herbacea*. Both the more diverse upland grasslands, and the high-altitude types, should be given priority when considering how much grassland to include within an ASSI.

The Manx Wildlife Sites Handbook states that all examples of unimproved upland acid grassland of 2 or more hectares in extent should be considered for Wildlife Site selection if they occur in association with other semi-natural habitats of Wildlife Site quality.

Minimum selection criteria for upland acid grassland

Upland pasture

U4 - Sheep's fescue/common bent grass/ heath bedstraw *Festuca ovina*- *Agrostis capillaris*-*Galium saxatile* grassland - upland examples;

U5 - Mat grass/heath bedstraw *Nardus stricta*-*Galium saxatile* grassland

U6 - Heath rush/sheep's fescue *Juncus squarrosus*-*Festuca ovina* grassland.

Upland examples of U4, U5 or U6 grassland or similar communities will qualify for selection if they are over 10ha in total extent and form part of mosaic of semi-natural habitats. Priority should be given to sites with a range of upland grassland types, rather than one NVC category alone.

U20 – Bracken/heath bedstraw *Pteridium aquilinum*-*Galium saxatile* grassland will not normally qualify for selection unless it forms part of a larger area of eligible upland habitats which would be fragmented or otherwise reduced in value for wildlife without the inclusion of the U20 habitat. However, special consideration should be given to including bracken/heath bedstraw grassland where it lies adjacent to or forms part of an area of regenerating native woodland and scrub and/or territory for birds of conservation concern.

Montane grassland

U16 – Greater woodrush/bilberry *Luzula sylvatica*-*Vaccinium myrtillus* grassland

Upland examples of U16 grassland or similar communities will **all** qualify for selection where they form part of mosaic of eligible semi-natural habitats. Priority should be given to sites which support rare species.

4.7.2 Upland heathland

Description and Distribution

Heather moorland is a striking and evocative feature of the Manx landscape, forming large areas of attractive purple hillside in early autumn, interspersed with bright, golden-yellow Western gorse *Ulex gallii* – which is so locally distinctive as to be generally known as Manx gorse. Despite the appearance of wildness, this is a man-made landscape, resulting from the clearance of sub-montane woodland and the deliberate maintenance of a low-growing shrub layer suitable for livestock grazing. Traditional upland heath management includes extensive grazing, occasional cutting and rotational burning. Other factors that are known to have affected the development of heathland in the past include accidental fires, and cutting of peat-rich upland turf for fuel and potash (Garrad 1972).

Upland heathland lies between the uppermost enclosed land and the onset of montane vegetation; in both cases there may be a gradient of vegetation types, but for the purposes of this chapter of the Criteria the altitude range is considered to be 250m to 600m. Within this range, heathland usually occurs wherever there is a thin to moderate layer of peat. Very deep peat (over 0.5m) is less common, and is

more likely to be associated with bog habitats, although some more typically heathland species may also occur within bogs. Examples of upland heathland may be found throughout the Island, on the hill-land in both the Southern and Northern Uplands. Due to heavy sheep grazing, the highest peaks (e.g. Snaefell and North Barrule) do not always support the best examples of upland heather heath, as the grazing regime favours grasses over dwarf shrubs. Better examples of upland dwarf shrub heath occur on the slightly lower slopes of Slieu Freoghane, Greeba Mountain, Cronk ny Arrey Laa and South Barrule (BDIS).

The Island's upland heathland vegetation is almost all of the type referred to by ecologists as dwarf shrub heath. It is usually defined as having at least 25% cover of dwarf shrubs, principally heather (or ling) *Calluna vulgaris*, bell heather *Erica cinerea* and Manx gorse *Ulex gallii*. Compared to lowland sites, upland heathland also tends to have a higher proportion of bilberry *Vaccinium myrtillus* – although poorly-managed burning and heavy grazing pressure can produce a similar proportion of bilberry in lowland heaths as well (www.Ukbap.org.uk, 2005). Other typically upland elements include crowberry *Empetrum nigrum*, deergrass *Trichophorum cespitosum*, and a small number of rare or locally-distributed plants such as cranberry *Vaccinium oxycoccus*, cowberry *Vaccinium vitis-idaea*, common cow-wheat *Melampyrum pratense* and fir clubmoss *Huperzia selago*. Of these, cranberry and deergrass are typical of wetter areas, fir clubmoss grows on rocky outcrops and short, exposed turf, and crowberry favours places where it cannot be out-competed by heather. The presence of a varied microtopography and hydrology is thus an important influence on the diversity of species present in an upland area.

Although neither dry nor wet upland heathland is noted for its great botanical diversity, these habitats are both associated with important bird populations. Species of note in the Manx uplands include hen harrier *Circus cyaneus*, curlew *Numenius arquata*, merlin *Falco columbarius* and short-eared owl *Asio flammeus*, whilst breeding golden plover *Pluvialis apricaria* are exceedingly rare. The significance of the uplands for birds relates to seasonal as well as year-round activities, hence upland moorland and grassland may provide winter feeding sites for birds such as chough *Pyrrhonorax pyrrhonorax*, which breed elsewhere in summer. Other fauna of interest in the uplands includes those invertebrates which prefer the habitat provided by dwarf shrub and lower plant cover. The uplands have not been subjected to extensive invertebrate survey; more general surveys, such as the DAFF Phase 1 Habitat Survey, indicate that upland moorland may be of importance for the Island's dragonfly and damselfly population, especially where it occurs in conjunction with upland streams and bog pools.

The usage, and hence the extent and vegetation structure, of upland heath on the Island has clearly varied according to human needs. Garrad, 1972 lists previous hill-land livestock as including both dairy and beef cattle, sheep, goats and pigs, with some collection of elm and other tree leaves for feed as well as dwarf shrubs. Following a gradual change to heavily sheep-dominated grazing, Garrad comments that " ... the vegetation is now monotonous heath with variety only in the wettest areas and the few really steep faces where even the most persistent and agile sheep could find no foothold". In the same chapter Garrad also states that "The monotony of the vegetation is largely due to a combination of prolonged over-grazing and repeated burning". In the decades since these opinions were published, the intensity of farming in the uplands appears to have changed, with some areas receiving less grazing pressure than previously, and some receiving more. Whilst there are still large areas of homogeneous heather, the regularity of heather burning has diminished in some parts of the Island, resulting in localised stands of older heather, taller gorse and occasional emergent tree growth. Marginal "intact" land has been allowed to become part of the adjacent open heath in some areas, and incorporated into improved and cultivated fields in others, as the pattern of need for supplementary hill-land grazing areas has changed. This dynamic pattern of change and adaptation is typical of upland areas in the British Isles as a whole (see Anderson, 2003 Chs 3-5).

Classification of upland heathland

Wet and dry heathland types known to occur in the uplands are listed on **Table 4.7b**; it should however be noted that some of these vegetation types may also occur in exposed conditions at lower altitudes, and/or in gradient with more typically lowland habitat types. Table 4.7b should therefore be considered alongside Tables 4.3a (Broad categories of dry lowland heathland on the Isle of Man) and 4.3b (Broad categories of wet lowland heathland on the Isle of Man), in Chapter **4.3 Lowland heathland**.

Status of upland heathland

Dry upland heathland

The DAFF Phase 1 Habitat Survey does not distinguish between dwarf shrub heath on lowland as opposed to upland areas, hence figures for each category are approximate. What is clear is that overall, dry heathland is a common habitat on the Island, over 90% of which occurs in upland areas. Unlike dry lowland heathland, upland heath tends to occur in large, unenclosed areas rather than small fragments. Site boundaries may not be well-defined unless a clear hill-wall is present, and many other habitats may occur with or adjacent to the heather. The Manx Wildlife Sites Handbook states that all areas of dry dwarf shrub heath greater than 2ha in extent, or occurring in association with other habitats of Wildlife Site quality, will qualify for inclusion as Wildlife Sites.

Dwarf shrub heaths as a whole are recognised as being of international importance because they are largely confined within Europe to the British Isles and the western seaboard of mainland Europe. The importance of heathland is recognised in the UK Biodiversity Action Plan, which lists both Lowland and Upland heathlands as Priority Habitats. The global and regional scarcity of heather moorland has led to its inclusion on Annex 1 of the EU Habitats Directive, under the general headings of *European Dry Heaths*, *Northern Atlantic wet heaths with cross-leaved heath *Erica tetralix**, and *Alpine and Boreal heaths*. Both the first two categories, and elements of boreal heath, occur on the Isle of Man. Of these, **European dry heaths** are found in every EU Member State except for Greece, but are only extensive in the western oceanic fringes of Europe. A high proportion of the EU resource of European dry heaths occurs in the UK, although this proportion is not as high as that of the other Annex 1 habitat, Northern Atlantic wet heaths with *Erica tetralix*. Dry heaths in the UK exhibit exceptional diversity in comparison with examples found elsewhere in the EU. (www.JNCC.gov.uk, 2005).

Wet upland heathland

Wet heath is much less common on the Isle of Man than dry heath – around 310ha in total, or approximately one twentieth of the total non-coastal heath on the Island. Unlike dry dwarf shrub heath, which is predominantly upland, around half of the wet heath recorded on the Island lies within small, fragmented and scattered lowland sites (BDIS). Lowland examples are more likely to be species-rich, but upland wet heath is significant for feeding upland waders and the small number of rare, specialised plants which can tolerate exposed, wet, acid conditions. The Manx Wildlife Sites Handbook states that all areas of wet dwarf shrub heath greater than 1ha in extent, or occurring in association with other habitats of Wildlife Site quality, will qualify for inclusion as Wildlife Sites.

Wet heath is extensively developed in western Britain and there are large areas of poor-quality habitat, almost all of which is in upland areas. One exception is **Northern Atlantic wet heaths with *Erica tetralix***, which are an Annex 1 habitat under the EU Habitats Directive. They are restricted to the Atlantic fringe of Europe between Norway and Normandy. A high proportion of the EU resource occurs in the UK, where it is restricted to highly localised sites in parts of southern and central England (JNCC). Small fragments of this habitat type occur on the Isle of Man, usually in conjunction with valley mires and fens. This habitat type is most commonly found on upland margins (BDIS), but may occur on hillsides wherever conditions are suitable.

Wet heath is an important habitat for a range of vascular plant and bryophyte species of an oceanic or Atlantic distribution in Europe, several of which have an important part of their EU and world distribution in the UK. The UK is one of the most important parts of the world for Atlantic bryophytes, and this is the most important habitat for many of these oceanic species of restricted world distribution (www.JNCC.gov.uk, 2005). The location of the Island would suggest that it may contribute to the UK's high proportion of bryophyte-rich wet heath; however, the full extent of bryophyte diversity in wet heaths on the Isle of Man is not well known at present. It is therefore an area which requires further study on both upland and lowland sites.

Upland heathland as a whole

Much of the Island's hill land is owned and leased by the Department of Agriculture, Fisheries and Forestry. Smaller areas are owned and managed by Manx National Heritage; the remainder is privately owned. As with Isle of Man, substantial areas of UK heathland are also publicly owned or managed, for

example by Scottish Office Agriculture, Environment and Fisheries Department (SOAEFD), the National Parks, and Forest Enterprise, and many areas lie in countryside designated as National Park, National Scenic Area (NSA) or Area of Outstanding Natural Beauty (AONB). The status of heather moorland means that the date and conditions under which heather can be burnt are defined by law throughout Ireland, the UK and the Isle of Man. Codes of good practice are published by the relevant Agricultural departments and take into account biodiversity, as well as agricultural and landscape, issues.

Within the UK large tracts of upland heath are notified as SSSIs/ASSIs, including at least 42,000 ha in England, 34,000 ha in Wales, 7,000 ha in Northern Ireland and 152,000 ha in Scotland. The UK Government has put forward areas that it considers qualify as Special Areas of Conservation (SAC) for these habitats. Some areas of upland heath are also classified as Special Protection Areas (SPA) under the EC Birds Directive, including the Bowland Fells and the North Pennines. Upland heathland is also represented within the UK National Nature Reserve (NNR) network.

Threats to upland heathland

The importance of heather moorland throughout the British Isles has led to considerable analysis of the threats posed both to its extent and quality. In general, the reduction in structural diversity and the trend towards acid grassland – both resulting from heavy grazing, sometimes in conjunction with burning – are considered to lead to a deterioration away from favourable conservation status. The UK Biodiversity Action Plan for Upland heathland states:

"There have been considerable losses of heather moorland in recent times. For example, 27% of heather moorland is estimated to have been lost in England and Wales between 1947 and 1980. On the Berwyn mountains in north-east Wales there was a 44% decline in the extent of heather-dominated vegetation between 1946 and 1984, whereas other upland sites in Wales have shown much smaller losses over similar periods. An estimated 18% was lost in Scotland between the 1940s and 1970s and the trend continued throughout the 1980s with a further estimated loss of 5%. Much of this loss is attributed to agricultural land improvements, heavy grazing by sheep (and, in certain areas, red deer and cattle), and afforestation.

It has also been estimated that 440,000 ha of land in the uplands in England and Wales have less than 25% cover of heather (ie grassland containing suppressed dwarf shrubs). There is likely to be further significant loss of heather moorland to acid grassland if current grazing levels and pressures continue. However, the conversion of heathland to acid grassland is not a purely recent phenomenon. On some sites in Wales (and elsewhere in UK) the major decline in heathland cover probably took place in the 19th century or even earlier." (<http://www.ukbap.org.uk>)

Associated threats to UK upland heathland are then identified as follows:

"Current factors affecting the habitat

- *Although upland heathland is dependent on management of grazing and burning to prevent succession to scrub or woodland, many upland heaths suffer from overgrazing and environmentally damaging burning regimes. Some of the problems caused by these management practices are described below. As well as direct loss of habitat, many areas of upland heathland are characterised by limited structural diversity with few natural transitions from open heath into scrub and woodland. Current management and other factors are continuing to prevent development of these features in most areas. Undergrazing occurs but is generally only of local significance.*
- *Agriculture is the dominant land use in the uplands. Most of the upland heathland resource falls within the Less Favoured Areas (LFAs). Livestock subsidies have led to a substantial increase in stocking rates (especially of sheep) in many upland areas environmental conditions are now also attached to all the main livestock subsidy schemes, although overgrazing is still a problem and losses of upland heathland habitat continue.*
- *High stocking levels of sheep, and to a lesser extent cattle, lead to heavy grazing of heather and other dwarf shrubs. High numbers of red deer *Cervus elaphus* are a problem in parts of the Scottish Highlands. Inappropriate methods of supplementary feeding and the absence or minimal use of shepherding also contribute to the problem of overgrazing.*

- *Heavy grazing by sheep, cattle or deer can prevent regeneration by native woodland and scrub, notably along upland heathland margins and stream sides where such habitat additions would be likely to enhance biodiversity value.*
- *Difficulties in negotiating agreements with commoners are hampering take-up of agri-environment schemes on common land.*
- *Conversion to grassland occurs through ploughing, reseeding, liming and fertilisation for agricultural purposes, particularly at lower elevations. Drainage and moorland 'gripping' also reduce the interest of wet heath. These factors have become less significant over the past ten years.*
- *Afforestation (mainly by non-native conifers) leads to direct loss of dwarf-shrub habitat, although temporary and permanent areas of heathland are now being created within some existing forests by restructuring after the first rotation.*
- *Poorly managed muirburn (ie large-scale and too frequent in operation) reduces the habitat quality of upland heath by causing a simplification of structure, loss of lower plant assemblages and erosion of peat.*
- *Encroachment by bracken can lead to a loss of biodiversity; this is a significant problem in some upland areas.*
- *Quarries, windfarms, communication masts, access tracks and certain other planning developments can impact directly on wildlife interest.*
- *Acidification, tropospheric ozone and nitrogen enrichment caused by atmospheric deposition can lead to vegetation changes including a reduction in the lichen and bryophyte interest. Nitrogen deposition can increase the likelihood of insect defoliation of upland heathland.*
- *Climate change could potentially lead to changes in vegetation composition and structure, although any increase in temperature may also be accompanied by possible increases in rainfall and wind speeds. The future position is still unclear but one of the dominant heathland species, heather, does have a relatively wide tolerance of temperature and rainfall, providing the overall climate remains oceanic. It is likely that within the time span of this plan other factors, notably high stocking levels, will have by far the greatest impact on upland heathland vegetation and species.*
- *Localised damage and threats from other forms of land use in the uplands, such as military use and recreation, are a concern.*
- *The interaction of two or more of the factors listed above often greatly increases the overall impact on upland heathland vegetation. For example, poorly managed burning followed by heavy grazing will result in the loss of dwarf shrubs more rapidly than would either factor in isolation. It is possible that grazing pressures interact with pollution to influence vegetation change." (<http://www.ukbap.org.uk>)*

With the exception of Commoners' rights issues and the impact of military use, the above threats have the potential to apply equally to Manx upland heathland. In particular the grazing levels in parts of the uplands have led to the development of extensive acid grassland on areas otherwise suitable for heather moorland (see p11 above, re. Garrad 1972). The risk of this is likely to increase where higher altitudes and more exposed conditions result in low vegetation recovery rates and a consequent high impact of overgrazing.

Minimum selection criteria for upland heathland

It is expected that upland sites will encompass a range of habitats, of which dwarf shrub heath is one element. However, due to its globally restricted status:

- all wet dwarf shrub heath over 4ha in extent, and
- all dry dwarf shrub heath greater than 8ha in extent,

which fulfil at least one of the Priority Site criteria listed in Part 3, will qualify for selection.

Where small sites are selected, priority should be given to those which exhibit the best diversity of structure and which support key populations of location-dependent rare species.

In addition to this:

- Wet dwarf shrub heath should always be included where it occurs in or adjacent to an area selected as an ASSI under other habitat criteria.

4.7.3 Blanket bog

Description and Distribution

The British Isles are uniquely suited to the formation of blanket bog, which “blankets” the less steep areas of hill and mountainsides in the wetter parts of the UK, Ireland and the Isle of Man. The high rainfall and relatively mild, humid conditions allow waterlogged, moss-dominated vegetation to form on undulating ground, building up into layers of peat which remain waterlogged as bog communities develop on top. Bogs where peat is accreting are known as “active”.

*“Active’ is defined as supporting a significant area of vegetation that is normally peat-forming. Typical species include the important peat-forming species, such as bog-mosses *Sphagnum* spp. and cottongrasses *Eriophorum* spp., or purple moor-grass *Molinia caerulea* in certain circumstances, together with heather *Calluna vulgaris* and other ericaceous species. Thus sites, particularly those at higher altitude, characterised by extensive erosion features, may still be classed as ‘active’ if they otherwise support extensive areas of typical bog vegetation, and especially if the erosion gullies show signs of recolonisation.” (www.JNCC.gov.uk, 2006).*

Peat accumulates very slowly under these conditions:

“Blanket bog peat accumulates in response to the very slow rate at which plant material decomposes under conditions of waterlogging. It is not, however, confined to areas of poor drainage but rather can cloak whole landscapes, even developing on slopes of up to 30°. The period over which blanket peat has been accumulating and the depth it can attain are very variable and not necessarily related. Studies indicate that most blanket peat development began 5000-6000 years ago, but the range extends from 9000 - 1500 years ago. There is evidence to suggest that some areas of blanket bog began to form following clearance of the original forest cover by early man, but the relative significance of this activity and changing climate on the historical and contemporary extent of the resource has yet to be determined. Peat depth is also very variable, with an average of 0.5-3 m being fairly typical but depths in excess of 5 m not unusual. There is no agreed minimum depth of peat which can support blanket bog vegetation.” (www.Ukbap.org.uk)

Active blanket bog is rare in the global context, and a Priority Habitat under the EU Habitats Directive. In the EU, **Blanket bogs** are found primarily in the UK and Ireland, but the extent of surviving habitat is now much reduced in Ireland. Blanket bogs are found in the north and west of the UK, extending from Devon in the south to Shetland in the north. *Scirpus – Eriophorum* mire predominates in the west, particularly at lower altitude, while *Calluna – Eriophorum* mire is particularly abundant in the east and at higher altitudes. *Erica – Sphagnum* mire is widely but patchily distributed (www.Ukbap.org.uk). Despite the apparent suitability of the Manx countryside for the formation of blanket bog, past habitat surveys have shown it to be quite limited in extent – in contrast to UK uplands, where it is one of the most extensive semi-natural habitat types. Rather than blanketing whole hillsides and mountain-tops, blanket bog on the Isle of Man usually occurs in patches on the less steep sides of the higher peaks in the Central and Northern hills – for example, Slieu Meayll in the Central Uplands. Smaller areas may be indistinguishable from localised hillside flushes, and true blanket bog, as opposed to flushes, only makes up around one quarter of upland bog on the Island (Sayle *et al*, 1995).

Whilst some groundwater-fed fen vegetation may occur within upland areas, most boggy wetland is reliant on rainfall and poor or impeded drainage to maintain the uniquely acidic and nutrient-poor conditions. The structure and vegetation composition of blanket bog is thus highly dependent on the underlying topography, and the extent to which it allows water to remain in the bog and/or peat to build up (JNCC, 1994). Human activities such as drainage, peat-cutting and burning can affect this process, particularly where smaller areas of blanket bog and flush lie adjacent to burnt and drained heather moorland. An altered or agriculturally improved hillside on deep peat is likely to support dry modified bog, which has some characteristics of blanket bog – for example, abundant cotton-grass *Eriophorum* spp. - but little in the way of *Sphagnum* mosses and no active peat generation. Such areas are usually dominated by heather *Calluna vulgaris* or cross-leaved heath *Erica tetralix*.

Blanket bog as it occurs in the UK, in very extensive areas, may exhibit a wide range of surface patterning such as series of bog pools and channels, tussocky as opposed to smooth vegetation, rills, flushes and erosion features:

"An important element in defining variation is the relative proportion of pools on the bog surface. In general, the proportion of surface patterning occupied by permanent pools increases to the north and west, although the precise shape and pattern of pools appears to depend on local topography as well as geographical location. Variety within the bog vegetation mirrors this pattern and is also affected by altitude. Similarly, the number of associated habitats and communities, such as springs, flushes, fens and heath, is greater in the milder, wetter and geologically and topographically more complex north and west."
(www.JNCC.gov.uk, 2006).

Blanket bog may also be sub-categorised according to morphology, e.g. saddle mires, watershed mires. The lack of very extensive blanket bog on the Island makes these features fewer and less obvious, but structural diversity of upland bogs is still an important consideration when assessing the diversity of a potential upland ASSI. In particular the presence of bog pools and other wetland features should be considered a priority for site selection.

As in the rest of the British Isles, Manx blanket bogs offer a store of archaeological and palaeological records, including preserved buried human artefacts, foundations of features such as round-houses, and a fossilised pollen record. Blanket bogs as a whole are increasingly recognised as "carbon sinks", which store atmospheric carbon in the preserved peat layer – a function which is lost where peat is removed faster than it can regenerate.

When assessing a potential upland ASSI with blanket bog, the full hydrological catchment on which the bog is dependent should be included within the ASSI boundary. This is especially important where the bog is thought to be active, i.e. where peat is still building up. Other habitats which depend on the same hydrological environment, such as wet heath, bog pools, streams, springs, fens and flushes, are important elements of biological and structural diversity and should always be included.

Classification of blanket bog

Blanket bog types known to occur in the uplands are listed on **Table 4.7b**; it should however be noted that some of these vegetation types may also occur in exposed conditions at lower altitudes, and/or in gradient with more typically lowland habitat types, which frequently occur on upland margins. Table 4.7b should therefore be considered alongside Tables 4.4a (Broad categories of bog on the Isle of Man) and 4.4b (Broad categories of valley mires and flushes on the Isle of Man), in Chapter **4.4 Bogs, fens and flushes**.

Status of blanket bog

Despite the apparent suitability of the climate and terrain, blanket bog is a rare habitat in the Isle of Man, even when modified bog and flushes are taken into account. The DAFF Phase 1 Habitat Survey found just 106ha of true blanket bog, and 455ha of flush and spring. Blanket bog thus makes up 0.64% of the total semi-natural habitat, or 3.41% if flushes are also included (Sayle *et al*, 1995). Dry modified bog makes up around 0.41% of the total semi-natural habitat, although it should be stressed that all three categories are fairly broad, often occur in complex associations with each other, and may be hard to differentiate at times. Likewise, the precise proportion of active bog is difficult to be sure of and unlikely to constitute all of the total recorded area of blanket bog. What is clear is that the habitat is limited on the Island, and tends to be fragmented at least in comparison with similar habitats in the UK and Ireland. The importance of bog in the uplands is recognised in the Manx Wildlife Sites Handbook, which states that the following will qualify for Wildlife Site status:

- *All areas of flush which are greater than 1ha in extent, or are part of a mire complex*
- *All areas of active blanket bog*
- *All areas of dry modified bog, except severely degraded examples*
- *All areas of wet modified bog, except severely degraded examples*
- *All areas of valley mire and basin mire, except severely degraded examples*

The total extent of blanket peat in the UK amounts to just under 1.5 million ha. There is no agreed figure for the extent of blanket bog vegetation. In terms of national cover of blanket peat soil (in the main >0.5 m deep) England supports some 215,000 ha, Scotland approximately 1,060,000 ha, and Wales has around 70,000 ha. Northern Ireland has approximately 140,000 ha of blanket bog vegetation. Significant

proportions of peat soil, probably in excess of 10%, no longer support blanket bog vegetation (www.Ukbap.org.uk). In recognition of the high proportion of the habitat type which occurs in the UK, and the UK's special conservation responsibility, blanket bog is a Biodiversity Action Plan Priority Habitat.

The global scarcity of active blanket bog has led to its inclusion on Annex 1 of the Habitats Directive as a Priority Habitat – with the caveat that the more species-poor, modified and inactive blanket bog that is frequent throughout much of the British uplands is not a priority. Protective measures in the UK include ASSI (Northern Ireland), SSSI, Ramsar Site, Special Area of Conservation and Special Protection Area designations, often for upland areas which include blanket bog as part of a wider mosaic of interrelated upland habitats. A small number are also managed as National Nature Reserves.

For parties to the Ramsar Convention there has been a recent move towards prioritising wet peatland habitats, which are currently under-represented in the Ramsar Site series.

Threats to blanket bog

The question of climate change, and how it is, and will, impact on this very climate-dependent habitat should be borne in mind whilst considering other threats.

Detailed data on the main threats to blanket bog on the Isle of Man are lacking, although it is clear that agricultural activities have a part to play in restricting the suitability of some hills for blanket bog – especially where hillsides have been subject to drainage and other agricultural improvement. Comprehensive data for changes to the total UK resource are similarly lacking, but studies in Scotland (where most of the resource lies and where it accounts for some 13% of the land area) suggest a 21% reduction in the extent of blanket mire between the 1940s and the 1980s. The greatest single cause of this reduction (51%) is afforestation, and substantial losses to forestry are reported from Wales. Further losses of extent and condition can be attributed to drainage and heavy grazing, peat cutting and atmospheric pollution, resulting in significant habitat change in, for example, mid and south Wales and the Pennines (www.Ukbap.org.uk). As with the other, related, globally-scarce upland habitat, heather moorland, the UK Biodiversity Action Plan lists many potential threats to blanket bog, the majority of which are of relevance on the Island:

"Current factors affecting the habitat

- The potential threat from climate change could over-ride many of the following factors. However, it is precisely because of the unknown effects climate change could have that it is important that as much of the resource as possible, representing its full biogeographical extent, is brought into, or maintained at, favourable condition. Pollution, from sulphate and nitrate deposition, may also be significant in certain areas, such as the Southern Pennines.*
- Drainage - extensive tracts of blanket bog have been drained in the past in attempts to improve the quality of the grazing. New drains continue to be dug and old drains cleaned in some areas. Even without maintenance most drains continue to lower the adjacent water table and some initiate erosion.*
- Heavy grazing (by sheep, red deer, cattle and horses) - especially if accompanied by supplementary feeding, burning, fencing and drainage, has a significant impact on vegetation. This is a particular concern on common land.*
- Burning - agricultural and sporting management both involve the use of fire to modify moorland vegetation for the benefit of livestock, grouse and deer in particular. Poorly managed and/or accidental fires can be particularly damaging to blanket bog.*
- Forestry - although new planting may be relatively small scale, some existing plantations are having an impact on the hydrology and species composition of adjacent areas of blanket bog, notably as the trees mature. Aerial application of fertilisers and pesticides can also result in drift on to adjacent bog.*
- Peat extraction - commercial extraction, though relatively limited in extent (some 2000 ha in Scotland), can have important local effects. Domestic cutting, most of which occurs on common land, is locally extensive (some 50,000 ha in Scotland) and, particularly where mechanical methods are employed, can have a significant impact, especially in Northern Ireland where the distinction between commercial and domestic activity can be difficult to determine.*
- Agricultural improvement - drainage, fertiliser application and conversion to pasture has occurred frequently in the past and can be of local significance.*

- *Recreation - many popular walking routes, some of which are also used by cyclists and horse-riders, traverse blanket bog areas which are very sensitive to such pressure. The increased use of all-terrain vehicles for recreational, agricultural and sporting activities can also result in local erosion.*
- *Planning developments - wind farms and communication masts, together with their associated infrastructure, are increasingly being proposed on areas of blanket bog, especially those at high altitude. There are also threats from hydro-electric schemes in Scotland.*
- *Erosion - high altitude bogs in particular, especially those in the Pennines and south Wales, are losing habitat through constant erosion of the peat mass. Some of this may be due to natural processes.*
- *Water course liming - where lime is added to lochs, lakes and rivers as a treatment for acidification, there may be detrimental implications for adjacent areas of blanket bog. Sometimes the bogs themselves have had lime applied.*

Minimum selection criteria for blanket bog

- Blanket bog should always be included where it occurs in or adjacent to area selected as ASSIs under other habitat criteria. This includes modified bog and flush.
- It is expected that upland sites will encompass a range of habitats, of which blanket bog is one element. However, due to its globally restricted status, all active blanket bogs over 2ha in extent which fulfil at least one of the Priority Site criteria listed in Part 3 will qualify for selection. Where small sites are selected, priority should be given to those which exhibit the best diversity of structure and morphology, in addition to their biodiversity.



The soft beds of *Sphagnum* moss on a blanket bog can support other plants growing on and through them, such as round-leaved sundew *Drosera rotundifolia* (red plants on bottom left of picture). However, the habitat is delicate: bike tracks such as those shown above may persist for months or even years.

Table 4.7.b: Broad categories of upland habitat on the Isle of Man

Type and Description	Species rich?	Closest National Vegetation Classification (NVC) categories	Rarities present?	Protected species present?
GRASSLANDS				
<p>Montane grassland</p> <p>The tops of high peaks such as North Barrule, Clag Ouyr and Bein y Phott may share some characteristics of montane vegetation communities, but true "arctic/alpine" vegetation occurs only above 600m, i.e. on the summit of Snaefell.</p>	<p>No. High-altitude habitat on the Island is low in species-richness, but represents a locally rare habitat which is very vulnerable to climate change.</p>	<p>U16 - <i>Luzula sylvatica</i>-<i>Vaccinium myrtillus</i> grassland Occasional on wet, peaty mountain tops where grazing is limited, notably Snaefell. No Island NVC samples</p>	<p>The tops of the highest peaks may support montane species which are rare on the Island, notably Cranberry <i>Vaccinium oxycoccus</i>, Alpine clubmoss <i>Diphasiastrum alpinum</i> and Least willow <i>Salix herbacea</i>.</p>	<p>May form part of the territory of protected bird of prey species.</p> <p>Several protected plant species occur on the highest peaks – see left.</p>
<p>Upland pasture (Sub-montane grassland)</p> <p>Frequent, generally above 250m. Always grazed - will revert to heathland and bog if left ungrazed.</p>	<p>No, although all types of upland pasture may occur within a diverse heath/grass/bog mosaic which is species-rich overall.</p>	<p>U4 - <i>Festuca ovina</i>- <i>Agrostis capillaris</i>-<i>Galium saxatile</i> grassland One of the commonest grassland types, often covering extensive areas especially on upland fringes. See also Section 3.1. One Island NVC sample.</p> <p>U5 - <i>Nardus stricta</i>-<i>Galium saxatile</i> grassland No Island NVC samples</p> <p>U6 - <i>Juncus squarrosus</i>-<i>Festuca ovina</i> grassland. Common, especially in the wettest and most exposed areas. No Island NVC samples.</p>	<p>Yes; a range of plant and animal rarities occur throughout the uplands, almost always in habitat mosaics where upland pasture is one element of an interrelated complex of habitats (BDIS).</p>	<p>May form part of the territory of protected bird of prey species; may be used by lapwings <i>Vanellus vanellus</i>, although not normally for breeding. May form part of golden plover <i>Pluvialis apricaria</i> habitat, and chough <i>Pyrrhocorax pyrrhocorax</i> winter feeding areas. Several protected plant species occur on upland pasture, including mountain pansy <i>Viola lutea</i>, Stag's-horn clubmoss <i>Lycopodium clavatum</i>, Ivy-leaved bellflower <i>Wahlenbergia hederacea</i>, Orchid</p>

		U20 - <i>Pteridium aquilinum</i> - <i>Galium saxatile</i> grassland very common: bracken "invaded" areas, often on margins of U5 grassland, eg. next to plantations. One Island NVC sample.		spp., Pale butterwort <i>Pinguicula lusitanica</i> , and Common cow-wheat <i>Melampyrum pratense</i> .
HEATHLAND				
Dry dwarf shrub heath May be dominated by heather, Manx gorse or bilberry. Often found in conjunction with other habitats such as wet heath, bog, and acid grassland. Marginal dry upland heathland may be subjected to agricultural improvement such as liming and drainage. Heavy grazing, particularly with frequent burning, can reduce the proportion of dwarf shrubs present until the habitat becomes entirely dominated by acid grassland species.	Pure stands of heather are not species-rich; upland margins and more sheltered sites may be relatively species-rich if a mosaic of habitat types is present.	H8a &b <i>Calluna vulgaris</i> – <i>Ulex gallii</i> (heather and western or "Manx" gorse) heath Six Island NVC samples, all within the species-poor or <i>Danthonia decumbens</i> (heath-moss) sub-communities. H10 <i>Calluna vulgaris</i> – <i>Erica cinerea</i> (heather and bell heather) heath One Island NVC sample Especially characteristic of the western margins of the British Isles. H12 <i>Calluna vulgaris</i> – <i>Vaccinium myrtillus</i> (heath and bilberry) heath One Island NVC sample. H18 <i>Vaccinium myrtillus</i> – <i>Deschampsia flexuosa</i> (bilberry and wavy hair-grass) heath No Island NVC sample.	Yes, may attract rare birds (see right). Rare invertebrates and lower plants are a possibility on this habitat, but both groups are data-deficient on the Island. Rare plants do not usually occur, but diverse sites may support unusual clubmosses such as fir clubmoss <i>Huperzia selago</i> and stag's-horn clubmoss <i>Lycopodium clavatum</i> . The best examples may support protected plants (see right) and locally-distributed species such as cowberry <i>Vaccinium vitis-idaea</i> , and crowberry <i>Empetrum nigrum</i> .	Yes. Protected bird species are likely, including hen harrier, kestrel, merlin and short-eared owl. May attract feeding croug especially during the winter. Lizards possible. Protected plants may include orchid species, mountain pansy <i>Viola lutea</i> , common cow-wheat <i>Melampyrum pratense</i> , lesser twayblade <i>Listera cordata</i> and stag's-horn clubmoss <i>Lycopodium clavatum</i> .

<p>Wet dwarf shrub heath</p> <p>Cross-leaved heath wet heath Both NVC types listed here are a type of mire; other mire types with less heathy characteristics may occur nearby (see below).</p>	<p>May be species-rich, especially compared with dry dwarf shrub heath of limited structural diversity.</p>	<p>M15 <i>Scirpus cespitosus</i> – <i>Erica tetralix</i> (deergrass and cross-leaved heath) wet heath No Island NVC sample Especially characteristic of the western margins of the British Isles. May occur with blanket bog.</p> <p>M16 <i>Erica tetralix-Sphagnum compactum</i> (cross-leaved heath and <i>Sphagnum moss</i>) wet heath No Island NVC sample</p>	<p>May support rare invertebrates and/or plants</p>	<p>Protected birds of prey such as peregrine, hen harrier and short-eared owl may use as feeding territory, the latter two forming roosts in wet areas of heathland/mire; also a possibility of lapwing. Lizards possible. A likely habitat for cranberry <i>Vaccinium oxycoccus</i>, heath-spotted orchid <i>Dactylorhiza maculata ssp ericetorum</i> and pale butterwort <i>Pinguicula lusitanica</i>.</p>
<p>BLANKET BOG</p> <p>Occurs rather sporadically throughout the Manx uplands, usually as a complex of varying vegetation communities amongst wet heath. Some spring-fed fens and flushes may also occur, which are not strictly blanket bog, but the two are treated together here as they are often contiguous and the habitat is ecologically very similar.</p> <p>Bog pools may occur, especially on hill tops, and are usually very small with limited associated vegetation. This is an under-researched habitat on the Island.</p>	<p>Not generally rich in vascular plants, although structurally diverse sites may support unusual species, and the more nutrient-rich sites (e.g. those which occur in conjunction with fens and flushes) may be more diverse than adjacent heath.</p>	<p>M1 <i>Sphagnum auriculatum</i> (bog moss) bog pool community. No Island NVC sample.</p> <p>M3 <i>Eriophorum angustifolium</i> (common cotton-grass) bog pool community No Island NVC sample</p> <p>M15 <i>Scirpus cespitosus</i> – <i>Erica tetralix</i> (deergrass and cross-leaved heath) wet heath – see wet heath, above.</p> <p>M17 <i>Scirpus cespitosus-Eriophorum vaginatum</i> blanket mire One Island NVC sample</p> <p>M18 <i>Erica tetralix-Sphagnum papillosum</i> raised and blanket mire No Island NVC sample</p> <p>M19 <i>Calluna vulgaris-Eriophorum vaginatum</i> blanket mire No Island NVC sample</p>	<p>May support rare invertebrates and/or plants, and attract feeding upland birds.</p>	<p>Protected birds of prey such as hen harrier and peregrine may use as feeding territory; also a possibility of lapwing. Lizards possible.</p> <p>A likely habitat for cranberry <i>Vaccinium oxycoccus</i>, pale butterwort <i>Pinguicula lusitanica</i> and dioecious sedge <i>Carex dioica</i>, in addition to Orchid species.</p>

		M25 <i>Molinia caerulea-Potentilla erecta</i> mire 1 Island NVC sample; mostly found on upland margins.		
OTHER HABITATS – see relevant chapters 4.1 – 4.5				
Valley mire, basin mire, fens and flushes	Yes, good examples may be relatively species-rich, especially when compared with blanket bog and structurally-poor heath.	These habitats occur sporadically, usually in very small areas, throughout the uplands. They are common to uplands, upland margins and unimproved lowlands. A minor but distinctive feature of upland margins is the presence of small springhead rills which support a vegetation community similar to M35 <i>Ranunculus omiophyllus</i> – <i>Montia fontana</i> (round-leaved water crowfoot and blinks) rill No Island NVC samples. See Chapter 4.4 Bogs, fens and flushes for details of commonly-distributed mire types that may also occur in upland areas, particularly upland margins.	Yes. May support rare invertebrates, birds and plants – including lower plants such as mosses. More research is needed on the significance of valley mire for invertebrates and lower plants on the Island.	Yes. These habitats may support a range of rare wetland plants, including good populations of orchids. They also attract wetland birds such as Snipe <i>Gallinago gallinago</i> , Curlew <i>Numenius arquata</i> and Lapwing <i>Vanellus vanellus</i> .
Open water	No, due to very low nutrient levels and an often dystrophic status. However, birds and invertebrates of conservation importance may rely on these areas.	Streams, ditches and small pools occur throughout the uplands, which form the catchment for the Island's rivers. See Chapter 4.5: Open water and swamp for details of more commonly-distributed open water and swamp types that may also occur in upland areas, particularly upland margins.	Not usually; the fast flow of upland streams militates against emergent vegetation growth. However, upland pools may support rare invertebrates – currently an under-recorded group.	Upland pools may attract waders such as lapwing, curlew and snipe.

Recommendations for selection of upland ASSIs

1. Upland ASSI selection should reflect the extensive nature of the uplands

As in adjacent countries, it is likely that upland ASSIs will make up a high proportion of the total protected area in the Isle of Man.

"The total area of upland (including blanket bogs) will be by far the largest of all habitats in the SSSI series: this is entirely appropriate, and in proportion to the extent of upland in Britain." (NCC 1989, p188)

2. Select a full range of representative upland mosaic areas

The **best example*** should be selected from within each of the "biodiversity hotspots" identified in Table 4.7a, using hydrological catchment, rather than hill-wall or administrative boundaries, as a guide to the extent of the protected area. Site quality assessment should follow the guidelines above (**Assessing uplands**, p3), including the key Priority Site Criteria listed.

*As a general guide, **best examples** will be:

- the most favourable according to the points listed in the Priority Site Criteria in Part 3
- and
- offering the most intact, unmodified and comprehensive inclusion of the hydrological catchment of a peatland, where applicable
- and
- forming a large ecological unit with a good range of associated upland habitats such as wet and dry dwarf shrub heath, bog, mire, rocky outcrops, springs, rills and flushes.
- and/or
- of critical importance for one or more populations of rare or vulnerable species.

3. Include best examples of each individual habitat type in the uplands

Upland areas which are important for the conservation of particular species or habitat sub-types but which are too narrow in overall biodiversity to have been identified as "hotspots" by the Balson, 2002 report should be selected according to the **minimum selection criteria** for each of the habitat types as described in sections **4.7.1 – 4.7.3** above.

4. Integrate with lowland ASSI selection wherever possible

The selection of upland ASSIs should relate to lowland ASSIs as follows:

- Eligible lowland habitats which occur in an ecological gradient with upland habitats – e.g. lowland heath, watercourses and mires – should be included in the boundary of upland ASSIs where they are ecologically contiguous. Thus, although the minimum altitude of 250m is taken as a guideline to the definition of an upland area, ASSI boundaries should not cut off important habitats just in order to create a purely upland site.
- The criteria for selection of flowing water (see Chapter 4.5: Open water and swamp) indicate that at least one river should be selected from source to sea. As the source of most Manx rivers lies in the uplands, this criterion should be borne in mind when selecting upland ASSIs.

4.8 Flora – sites of exceptional importance for plant conservation

Notes:

The species scoring system is based on the rarity/significance of the species, using lists which may themselves be updated at intervals. This avoids having to review the species criteria scoring lists separately. To qualify under plant species criteria the plant in question should normally have been confirmed as present within the 12 months before designation.

For scoring purposes a species may only be included **once**, at its highest-scoring category (e.g. a Schedule 7 Red Data Book species scores 10, not 12).

International significance is denoted by inclusion of naturally resident species which are listed on Appendix I of the Bern Convention and/or Annexes II, IV or V of the EC Habitats and Species Directive.

British Isles significance refers to inclusion of species in the British or Irish Red Data Book as *Extinct*, *Endangered* or *Vulnerable*.

Plants listed on Schedule 7 of the Wildlife Act 1990 are considered nationally important in the Manx context, in particular those which are known to occur at three sites or fewer. In the case of Schedule 7 species which are listed as a group - e.g. all orchids - the score applies separately to each species present.

Priority sites for plants are those with a score of 10 or higher.

CRITERIA	SCORE
International significance	10
British Isles significance	10
Schedule 7 species excluding those present on > 3 sites	2
Schedule 7 species occurring on > 3 sites.	1

Examples:

A site with 5 Schedule 7 plants found within the last 12 months, all of which are known to occur at three or fewer sites, would score **10** and be a **priority site** for ASSI selection;

A site with any British Isles Red Data Book species known to occur within the last 12 months would also score **10** and be a **priority site** for ASSI selection.

A site which harboured three Schedule 7 plants, one of which occurs at more than three sites Island-wide, would score **5** points, and would **not** be considered a priority site for ASSI selection solely on the basis of the plant species present.

4.9 Avifauna – sites of exceptional importance for birds

General

Birds can be used in site assessment in two ways. Firstly, the presence of a particular bird community in a specific habitat can indicate that a site is of high quality and worthy of designation as ASSI. Under the site quality guidelines in 3.2, the naturalness and diversity of a site can be indicated by the presence of a typical assemblage for the habitat. This will include some birds that rarely aggregate in large numbers but may rely on particular habitats to sustain them, eg. merlins on moorlands, or corncrakes in hay fields. It is important that such habitats are maintained in order to sustain such species on the island in the future. The scores for Breeding Bird Assemblages (criteria below) may be used in site comparison. For instance, a woodland with a score of 16, will not qualify for its breeding bird assemblage alone, but will be considered of greater interest for its habitat-specific birds than a site with only a score of 10. In this situation, where more than one of the important habitats is included in a site, the score may be raised by further species of interest in the second habitat, but double scoring is not applicable. The thresholds will not be relevant in this situation.

Secondly, sites may be considered of interest because of their bird interest itself, due to the regular presence of concentrations of a species. Such sites are easy to pick out for colonial breeders but some species tend not to congregate, so areas with the highest densities or linked to other interest should be put forward.

The intention is to protect populations of all of the regularly occurring species on the island within the ASSI series. Common species are likely to be included, by default, within ASSIs designated for other reasons and would not normally provide the only interest of a site. As a priority, the best site for any species of conservation concern, should be put forward for designation.

There may also be sites that are of particular importance because of their scientific value, such as bird observatories and long-term survey areas. A long history of recording at a site can provide very important information on changes in the fauna and in the environment, that may be impossible to detect in other ways. Although not an eligibility criterion in itself, such importance may form a reason for choosing one eligible site over another.

Birds populations can be divided into breeding and non-breeding populations. Some species are sedentary and some migratory at least to some extent. In order to protect a population of birds, it is important to ensure that habitats remain available for feeding birds all year round. Whereas some species move up and down the hills, others cross international borders in the search for a year-round food supply (see below). Also, breeding sites will be required and some species require undisturbed areas where they can roost in relative safety when their feeding areas are unavailable, such as waders during high tide.

The proportion of a population found on a site in a particular season has been accepted as a criterion for assessment. This not only shows that a site is important for the maintenance of the population, due to its holding a significant proportion of it, but is also self-adjusting for rarity, because more sites will become eligible for species with small populations than those that are common and widespread. There is therefore an in-built bias towards the protection of rare species.

International obligations

The Isle of Man has obligations under certain international conventions for the conservation of wildlife. These include the Convention on the Conservation of Migratory Species of Wild Animals (Bonn Convention) and the Agreement on the Conservation of African-Eurasian Migratory Waterbirds made under that convention, the Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention) and the Convention on Wetlands of International Importance, especially as waterfowl habitat (Ramsar Convention).

Sites that are recognised as internationally important for birds may therefore be eligible for designation as Wetlands of International Importance, for waterfowl under Ramsar, or Areas of Special Conservation Interest, for birds of European conservation concern under Bern. The latter is equivalent to an EU Special Protection Area, as SPAs are accepted as the EU's ASCIs. Such sites should receive legal protection by designation as ASSIs, and the criteria have been written to allow this. Sites of international importance for birds will also be considered as of national importance.

Migratory birds are of particular concern. Some move around a series of sites during the non-breeding season, searching for winter food along a traditional route and frequently relying on particular sites for their survival when moving along the chain of feeding areas. Areas used by significant numbers of wintering and migrating waterfowl must therefore receive protection. In the Isle of Man the numbers of birds at such sites is relatively small in comparison with north-west England, but this is partly because of the lack of suitable habitat on the island, which lacks any large, productive mudflats. However, over a year many birds will pass through the island, stopping to take on food to burn on the next stage of the journey. Manx sites, though holding relatively small numbers of birds at any one time are important in the maintenance of an international network of sites for such species.

Important Bird Areas (IBAs) are sites that support internationally important assemblages of birds. Together they form a coordinated international network of sites, the protection of which relies on cooperation between nations. Sites in the Isle of Man are listed in Pritchard *et al.* (1992). These were selected for their importance in the context of the European Community or globally. Five sites have been recognised: the Isle of Man Sea Cliffs, Calf of Man, The Ayres, Ballaugh Curraghs and The Isle of Man Hills. The sites are recognised on the basis of criteria used by Ramsar, BirdLife International (as the ICBP) and the then Nature Conservancy Council. They were designed to enact requirements under the Ramsar Convention and the EU Birds Directive. Although the Isle of Man has no responsibilities under the Birds Directive, it does have a responsibility to protect sites of international importance for birds. The Birds Directive arose from the Bern Convention to which the island is a party. The listed sites are therefore of a high priority for conservation and will be considered priority areas for designation as ASSI.

Arable fields and highly improved grasslands will not normally be included as feeding and roosting sites, due to their wide availability, unless they are put forward due to particular aspects of their ecology or location which makes them special. Seasonally flooded grasslands, however, can be especially important for birds and would therefore receive consideration, but such areas are not common on the island.



Shy breeding birds such as the curlew *Numenius arquata* (centre of picture) benefit from large protected areas such as the Ballaugh Curragh ASSI and Ramsar Site

Sites of British importance

Regular status assessments of bird populations are made at the British level, such as the list of Birds of Conservation Concern for Britain, including the Isle of Man but excluding Ireland. An accepted level for national importance in Britain is 1% or more of a population.

Sites of Manx importance.

Some species that are common in Britain are rare on the island and at risk of loss as breeding species or are retained only as visitors, having bred previously. Sites of importance for maintaining the presence of species on the island should be protected as ASSI. Sites should be regularly used (see below), based on recent information.

Site information

Where possible site selection should be made on good data, ideally from 5 consecutive years. However, the records do not always allow this. The publication of the new atlas will provide a view of the use made of the island by each species in detail, as a snapshot in time. Some detailed site recording may, however, be necessary to show that the expected bird numbers are seen consistently.

Some species of particular interest in Britain and the Isle of Man, are counted fully at intervals, but not annually, such as seabird colonies and hen harrier nests. Three such counts, which could span a period of thirty years, are acceptable and show the continued use of a site over an extended period. Mean numbers can be compared but sites with high numbers of breeders or with a high productivity of chicks will be favoured over declining sites.

Minimum selection criteria for sites of high importance for birds

1. Sites of British importance: any site holding at least 1% of the population of a species in Britain and the Isle of Man, either for a breeding or non-breeding population.

For breeding sites of rare species it is important that the site is regularly used (see Site Information above) and the site boundary should normally include all the land areas required for the survival of the bird during breeding.

2. Sites of Manx importance for birds of local conservation concern: for any species listed on Schedule 1 of the Wildlife Act or the Red List of Birds of Conservation Concern (for Britain or for Ireland), the best site providing for its needs in each relevant season will qualify.

A site may be rated highly as having the largest proportion of the Manx population on a single site or having the highest density of the species on the island. If a species has populations that rely on separate, regularly used sites in different seasons, the important sites for a particular population may be chosen, rather than simply the best wintering and breeding areas, to ensure the protection of an important population in all seasons.

3. Smaller seabird sites and heronries: for any heron or seabird, except herring gull, if no site has been chosen for this interest under (1) or (2), then the best breeding site with at least 10 breeding pairs regularly seen, should be picked.

This may be justified on the basis of the number of chicks produced on a site each year, or if such data is not available, the number of pairs nesting. This criterion caters for species that aggregate in colonies but not necessarily in large numbers that would exceed the 1% threshold. Seabird populations in the British Isles are of special international importance so this criterion will ensure that the significant sites are eligible.

4. Breeding bird assemblages: sites with a very good range of the birds peculiar to a particular habitat are eligible.

The habitats given below have been chosen because they have a particular bird fauna. These vary in the numbers of species that they support, so for each habitat a list of typical species has been provided, with scores rated by the rarity of the species in Britain (see ratings by British breeding population, Nature Conservancy Council 1989). To assess a site, the scores are summed for each listed species of bird recorded as a probable breeder in a majority of recent years for which information is available.

Species regularly using a site for essential activities may be included, even if the nest is outside of the site. Any site with a score equal to or exceeding the stated threshold is eligible for designation. The species and scores are given in Table 4.9a.

Where substantial areas of more than one of these habitats is included in a site, the site could become eligible if one or more of the thresholds is exceeded, the whole site might then be considered as an ecological unit if applicable (see guidelines on boundaries), or the wider site could be eligible if the score lists are summed for each major habitat and exceed the sums of the thresholds (double counting for a species is allowed).

The Manx thresholds were adapted from the Nature Conservancy Council figures by first summing the scores for Manx species that breed in the relevant habitat on the island, giving a potential maximum (scores of 5 or 6 were not included in this maximum). The thresholds were set at about ½ of this maximum score, and adjusted to a level appropriate to make eligible the best quality sites on the island. An exception was woodland, in which the threshold is set at 2/3 due to the large number of species found in many woods.

If a rare species begins to breed regularly at a site, having an index listed by the NCC under a similar habitat, the score given there may be used to add value to the total for a site. The threshold would not be reassessed unless major changes occur or an overall revision is necessary. Sites with very rare breeders, establishing themselves as regular, will therefore have a particularly high score.

Habitat	Threshold score
Lowland damp grasslands	11
Lowland fen without open water	11
Open waters and their margins	22
Upland moorland and grassland with water bodies	18
Upland moorland and grassland without water bodies	16
Scrub (excluding heath)	9
Woodland	18
Hard coastline	21
Soft coastline	21



Upland moorland sites demand a higher threshold score than lowland grassland and fen

Table 4.9a: Criterion 4 scores for habitatsLowland damp grasslands

Shelduck	2	Curlew	2
Teal	3	Cuckoo	2
Shoveler	4	Whinchat	2
Quail	5	Grasshopper warbler	2
Corncrake	6	Sedge warbler	1
Lapwing	1	Reed bunting	1
Snipe	2		

Lowland fen without open water

Little grebe	2.5	Cuckoo	2
Teal	3	Whinchat	2
Shoveler	4	Grasshopper warbler	2
Water rail	3	Sedge warbler	1
Snipe	2	Reed bunting	1

Open waters and their margins

Little grebe	2.5	Common tern	3
Mute swan	3	Cuckoo	2
Shelduck	2	kingfisher	3
Teal	3	Grey wagtail	2
Shoveler	4	Grasshopper warbler	2
Tufted duck	3	Sedge warbler	1
Water rail	3	Reed bunting	1
Ringed plover	3	Oystercatcher	2
snipe	2	Curlew	2

Upland moorland and grassland with water bodies

Hen harrier	4	Grey wagtail	2
Merlin	4	Whinchat	2
Peregrine	4	Stonechat	2
Red grouse	1	Wheatear	2
Snipe	2	Chough	4
Curlew	2	Raven	3
Short-eared owl	3		

Upland moorland and grassland without water bodies

Hen harrier	4	Short-eared owl	3
Merlin	4	Whinchat	2
Peregrine	4	Stonechat	2
Red grouse	1	Wheatear	2
Snipe	2	Chough	4
Curlew	2	Raven	3

Scrub (excluding heath)

Cuckoo	2	Garden warbler	1
Long-eared owl	3	Blackcap	1
Whinchat	2	Linnet	1
Stonechat	2	Redpoll	1
Grasshopper warbler	2		
Whitethroat	2		

Woodland

Grey heron	3	Chiffchaff	1
Sparrowhawk	2	Goldcrest	1
Woodcock	2	Spotted flycatcher	1
Stock dove	1	Long-tailed tit	1
Cuckoo	2	Coal tit	1
Long-eared owl	3	Treecreeper	1
Redstart	1	Raven	3
Garden warbler	1	Redpoll	1
Blackcap	1		
Wood warbler	2		

Hard coastline (including rocky cliff and associated pebble beach)

Fulmar	1	Black guillemot	2
Manx shearwater	1	Razorbill	1
Cormorant	3	Puffin	1
Shag	2	Rock pipit	2
Peregrine	4	Stonechat	2
Shelduck	2	Wheatear	2
Eider	2	Raven	3
Oystercatcher	2	Chough	4
Ringed plover	3		
Kittiwake	1		
Great black-backed gull	2		
Lesser black-backed gull	2		

Soft coastline (including soft cliff, saltmarsh, sand dune and associated sand or pebble beach)

Fulmar	1	Arctic tern	2
Shelduck	2	Little tern	3
Eider	2	Cuckoo	2
Oystercatcher	2	Sand martin	1
Ringed plover	3	Rock pipit	2
Lapwing	1	Stonechat	2
Snipe	3	Wheatear	2
Curlew	2	Grasshopper warbler	2
Black-headed gull	1	Sedge warbler	1
Great black-backed gull	2	Linnet	1
Lesser black-backed gull	2	Reed bunting	1
Common tern	3		

5. Species rich sites: sites with at least 39 breeding species or 83 species on passage, or wintering, in recent years.

This criterion is especially appropriate for very varied sites with habitat mosaics and for passage/wintering sites, either of which may hold large numbers of species but small populations of each. These thresholds have been produced by reducing the equivalent UK thresholds (Nature Conservancy Council 1989) by the difference in the size of the British and Manx Species Lists for categories A and C (see Manx Ornithological Society and British Ornithologists' Union), i.e. by 55%, because there are fewer species recorded on the Isle of Man. The categories of passage and wintering have been combined because recent information has shown that these cannot be separated in a simple way. The higher passage threshold has been adopted as, at the time of writing, diverse sites in the Isle of Man have greater significance as a passage stop-over than as a site for longer-term over-wintering species. Only highly diverse sites are likely to qualify but these should include the top passage/wintering sites that many small populations may rely on.

4.10 Fauna – sites of exceptional importance for animals

Notes:

The species scoring system is based on the rarity/significance of the species, using lists which may themselves be updated at intervals. This avoids having to review the species criteria scoring lists separately. To qualify, the species should normally have been confirmed as present within 3 years.

For scoring purposes a species may only be included **once**, at its highest-scoring category (e.g. a Schedule 5 Red Data Book species scores 10, not 12).

International significance is denoted by inclusion of naturally resident species which are listed on Appendix II or III of the Bern Convention, Appendix I of the Bonn Convention and/or Annexes II, IV or V of the EC Habitats and Species Directive.

British Isles significance refers to inclusion of species in the British or Irish Red Data Book as *Extinct*, *Endangered* or *Vulnerable*.

Animals listed on Schedule 5 of the Wildlife Act 1990 are considered nationally important in the Manx context, in particular those which are known to occur at three sites or fewer. In the case of Schedule 5 species which are listed as a group - eg. Vespertilionid bats - the score applies separately to each species present.

Note: Bat roosts will only be considered for designation if this is considered appropriate.

Priority sites for animals are those with a score of 10 or higher.

CRITERIA	SCORE
International significance, excluding bats	10
British Isles significance, excluding bats	10
Schedule 5 species, excluding widespread species such as common pipistrelle, common frog and common lizard. Mobile species should spend significant amounts of time at the site*	2
Widespread species on Schedule 5	1

* Significant amounts of time may refer to bat roosts, regular observations of feeding bats, or seal pupping or haul-out areas.



Common frog *Rana temporaria*, a protected species, photographed by a pond in Ballaugh Curragh ASSI

Part 5: Bibliography and References

- Allen D E (1978) "The present day fauna and flora of Man as indicators of the date of the Flandrian severance" in *Man and Environment in the Isle of Man*, ed Davey P, 9-13 Oxford: British Archaeological Reports Series 54 (i).
- Allen, D. E. 1984. *Flora of the Isle of Man*. Manx Museum and National Trust, Douglas, Isle of Man.
- Anderson, P (ed.) 2003. *Upland ecology, tourism and access Proceedings of the 18th Conference of the Institute of Ecology and Environmental Management*. (IEEM, Winchester)
- Asher, J., Warren, M., Fox, R., Harding, P., Jeffcoate, G., & Jeffcoate, S. 2001. *The Millennium Atlas of Butterflies in Britain and Ireland*. (Oxford University Press, Oxford)
- Balson, J. 2002. *Modelling a network of candidate conservation areas in the uplands of the Isle of Man, using optimisation algorithms*. (MSc project, DAFF/UCL)
- Barne, J. H., Robson, C. F., Kaznowska, S.S., Doody, J. P. & Davidson, N.C. (eds.) 1996. *Coasts and seas of the United Kingdom Region 13 Northern Irish Sea Colwyn Bay to Stranraer, including the Isle of Man*. (JNCC, Peterborough)
- Biodiversity The UK Action Plan* 1994. (HMSO, London)
- Biological Data Information System (BDIS) – DAFF's biological records system, which incorporates a fully-digitised Phase 1 and 2 habitat map of the whole Island on ArcView GIS, and biological records held on a Recorder 2002 database, from which records are linked to the maps.
- Blackshall, J., Manley, J., & Rebane, M. 2001. *The upland management handbook*. (English Nature, Peterborough)
- Boyce, D. C. & Fowles, A. P. 1989. *Invertebrate conservation in the Isle of Man – An assessment of selected sites of ecological interest 30 April – 5 May 1989*. (NCC, Aberystwyth)
- Callister, J "Dog", 2001: *Manks flower names* (unpublished list)
- Clement, E. J. & Foster, M. C. 1994. *Alien plants of the British Isles a provisional a provisional catalogue of vascular plants (excluding grasses)*. (BSBI, London)
- Cottrell, J. E., Munro, R.C., Tabbener, H. E., Gillies, A.C.M., Forrest, G.I, Deans, J. D. & Lowe, A. J. 2002. *Distribution of chloroplast DNA variation in British oaks (Quercus robur and Q. petraea): the influence of postglacial colonisation and human management*. *Forest Ecology and Management* 156 (2002) 181-195. (Elsevier)
- Crofts, A. & Jefferson, R. G. (eds.) 1999. *The Lowland Grassland Management Handbook 2nd Edition*. (English Nature/The Wildlife Trusts)
- Cullen, J. P. & Jennings, P. P. 1986. *Birds of the Isle of Man*. (Bridgeen Publications, Douglas)
- DEFRA, June 2000. *Water Quality, a guide to water protection in England*. (DEFRA, information leaflet; available at <http://www.defra.gov.uk/environment/water/quality/guide/water.htm>)
- Drewitt, AL & Manley, VJ 1997. *The vegetation of the mountains and moorlands of England National assessment of significance* (English Nature Research Report No. 218, English Nature, Peterborough).
- English Nature 2002. *Lowland Heathland a cultural and endangered landscape* (booklet - an EN External Relations Team publication, English Nature, Peterborough)
- English Nature 2001. *The Upland Management Handbook* (English Nature, Peterborough)
- Falk, Steven 1991. *Research & survey in nature conservation No. 35 A review of the scarce and threatened bees, wasps and ants of Great Britain*. (NCC, Peterborough)
- Falk, Steven 1992. *Research & survey in nature conservation No. 39 A review of the scarce and threatened flies of Great Britain (part 1)*. (JNCC, Peterborough)
- Ford, Trevor D. 1993. *The Isle of Man Geologist's Association Guide No. 46*. (The Geologists' Association)
- Fullen, M. A., Harris, J. & Kear, B. S. (Gen. ed. P. J. Davey) 1996. *Soils of the Isle of Man Centre for Manx Studies Research Report 5*. (CMS, Douglas)
- Fullen, M., Harris, J. & Kear, B. 1999. *Soil-forming factors on the Isle of Man* (article in Geography Review, September 1999)
- Garrad, Larch S. 1972. *The Naturalist in the Isle of Man*. (David & Charles, Newton Abbot)
- Garrad, Larch S. 1990. *Lateglacial and Postglacial environmental history*. (in Robinson & McCarroll, 1990).
- Garrad, Larch S. 1990. *Nature conservation*. (in Robinson & McCarroll, 1990).
- Garrad, Larch S. 2003. *The Manx Garden "A History of Manx Gardens" updated*. (Manx Heritage Foundation, Douglas)
- Harris, J., Fullen, M. A., & Hallett, M. D. (Gen. ed. P. J. Davey) 2001. *Agricultural soils of the Isle of Man CMS Research Report 9*. (CMS, Douglas)
- JNCC – see key websites, below.
- JNCC 1994. *Guidelines for selection of biological SSSIs: bogs*. (JNCC, Peterborough)
- JNCC 1996. *Guidelines for selection of biological SSSIs: intertidal marine habitats and saline lagoons*. (JNCC, Peterborough)
- Hawthornthwaite, D. L. (ed.) 2001. *The Changing Wildlife of Great Britain and Ireland*. (Taylor & Francis, London)
- Hodgetts, N. G. 1996. *The conservation of lower plants in woodlands*. (JNCC, Peterborough)
- Hopson, A & Lamb, J (eds.) 1995: *Manx Hill-land Report*. (MNCT)
- Keehan, B. 1999 (revised by Dullaghan, P., 2000). *Manx Wildlife Sites Handbook*. (MWT, St Johns, Isle of Man)

- Lack, Peter 1992. *Birds on Lowland Farms*. (HMSO)
- Lamb, J. 1999. *A terrestrial survey of the coast of the Isle of Man*. (MWT, St. Johns, Isle of Man)
- MNCT (now MWT) (various authors) 1995: *Proceedings of the Manx Hill-land Seminar 11th – 13th April 1995* (MNCT)
- MNH 1998: *Ballaugh Curragh Management Plan draft edition 1*. MNH, Douglas, Isle of Man.
- NCC 1989. *Guidelines for selection of biological SSSIs*. (NCC, Peterborough)
- Osborn, T 2001: *Climate Research Unit Information Sheet 15: The changing intensity of rainfall over Britain* (CRU, University of East Anglia, Norwich)
- Parsons, M. S. 1993. *UK Nature Conservation No. 11 A review of the scarce and threatened pyralid moths of Great Britain*. (JNCC, Peterborough)
- PPP (Precautionary Principle Project) 2005: *Guidelines for applying the precautionary principle to biodiversity conservation and natural resource management*. (www.pprinciple.net)
- Preston, C. D, Pearman, D. A., Dines, T. D. 2002. *New Atlas of the British and Irish Flora*. (Oxford University Press)
- Pritchard, D. E., Housden, S. D., Mudge, G. P., Galbraith, C. A., Pienkowski, M. W., (eds.) 1992. *Important bird areas in the United Kingdom including the Channel Islands and the Isle of Man*. (RSPB, Sandy)
- Pritchard, T. (compiler) 1975. *An Ecological Survey of the Isle of Man Part 1 Nature Conservation Appraisal*. (NCC, Bangor)
- Ratcliffe, D. A. (ed) 1977. *A Nature Conservation Review The selection of biological sites of importance to nature conservation in Britain*. . (Cambridge University Press)
- Robinson, V. & McCarroll, D. 1990 (eds.). *The Isle of Man Celebrating a sense of Place*. (Liverpool University Press)
- Rodwell, J. S. (ed) 1992 ¹⁻⁵. *British Plant Communities*. (Cambridge University Press)
- Volume 1: Woodlands and scrub*
- Volume 2: Mires and Heaths*
- Volume 3: Grassland and montane communities*
- Volume 4: Aquatic communities, swamps and tall-herb fens*
- Volume 5: Maritime communities and vegetation of open habitats*
- Samson, L, 2002. *Ponds and Dubs Survey report*. (DAFF, Isle of Man)
- Sayle, T., Lamb, J., Colvin, A., Harris, B., 1995. *Isle of Man Ecological Habitat Survey Phase 1 Report 1991-1994*. (DAFF)
- Scott, A. 1997. *Road verge survey project report* (DoT, MNCT)
- Stace, Clive 1997. *New Flora of the British Isles Second Edition*. (Cambridge University Press)
- Stewart, A., Pearman, D. A., Preston, C. D. (compilers & editors) 1994. *Scarce Plants in Britain*. (JNCC)
- UK Biodiversity Group, 1998. *Tranche 2 Action Plans: Terrestrial and Freshwater Habitats* (HMSO – also, see key websites, below)
- Wigginton, M. J. (ed.) 1999. *British Red Data Books 1 Vascular plants 3rd Edition*. (JNCC, Peterborough)
- World Wildlife Fund October 2004: *Dead Wood Living Forests*. (<http://www.panda.org>).

Abbreviations:

- BDIS – Biological Data Information System
- BSBI – Botanical Society of the British Isles
- BDIS - Biological Data Information System (DAFF internal biological database)
- CMS – Centre for Manx Studies
- DAFF – Department of Agriculture, Fisheries and Forestry
- DEFRA – Department for Environment, Food and Rural Affairs (UK Govt. Department)
- DOT – Department of Transport (Manx Government, unless otherwise stated)
- HMSO – Her Majesty's Stationery Office (now The Stationery Office)
- JNCC – Joint Nature Conservation Committee
- MNCT – Manx Nature Conservation Trust – now MWT
- MNH – Manx National Heritage (also: Manx Museum, Manx National Trust)
- MWT – Manx Wildlife Trust (previously MNCT)
- NCC – Nature Conservancy Council (now JNCC)
- NVC – National Vegetation Classification (as detailed in Rodwell et al, 1992)
- RSPB – Royal Society for the Protection of Birds
- UCL – University College, London
- UK BAP – UK Biodiversity Action Plan
- WWF – World Wildlife Fund

Key websites:

<http://www.forestry.gov.uk> Forestry Commission – this website includes information on habitats that occur in conjunction with woodland, as well as on Forestry Commission plantations and their management e.g. <http://www.forestry.gov.uk/forestry/Lowlandheath>

http://www.jncc.gov.uk/Publications/JNCC312/UK_habitat_list.asp - JNCC webpage of habitats listed under Annex 1 of the Habitats Directive, with links to habitat descriptions and selection criteria e.g:
UK Marine Special Areas of Conservation (SACs) Project <http://www.ukmarinesac.org.uk/index.htm>

www.rspb.org.uk – bird information e.g. www.rspb.org.uk/birds/guide/c/chough/feeding.asp and related habitat information, e.g.

RSPB: Lowland heathland habitat information factsheet

(http://www.rspb.org.uk/Images/Lowland%20Heathland_tcm5-45464.pdf , 2005)

www.ukbap.org.uk - the website of the UK Biodiversity Action Plan, containing Priority and Broad Habitat descriptions and status accounts, taken from the UK Biodiversity Group Tranche 2 Action Plans - Volume II: Terrestrial and freshwater habitats (December 1998)

January 2008

Compiled and edited: L Moore

Chapter reviews: each chapter has been contributed to and agreed by the Wildlife Committee over the years 2003 – 2007. Dates of Wildlife Committee acceptance of the Criteria are available from DAFF.

Photographs: L Moore, except chough stamp and coin pictures on p99: Isle of Man Post

© DAFF

Areas of Special Scientific Interest

The Government working with landowners to conserve our natural heritage

The Isle of Man contains many areas of particular interest for their wildlife habitats, geological features and landforms. Some are important as wildlife havens, as natural laboratories for science, or as educational resources displaying our geological history and richness of wildlife. The protection of our natural resources is a responsibility that we all share. Areas of Special Scientific Interest (ASSIs) are given legal protection as the best examples of this natural heritage. Some areas are of international importance and we have an added responsibility to maintain their quality.

Site Selection

The selection of ASSIs is based on scientific criteria. They can cover public or private lands. This designation sets in place a statutory consultation process so that the Department of Environment, Food and Agriculture (DEFA) can provide advice and financial support for their protection and management. These areas can be served best by a partnership between the Department and the land owners and managers. By working together we aim to maintain a living and working countryside in which future residents can enjoy the same benefits that we have.

Notification

DEFA, after consulting the Wildlife Committee, can notify sites as ASSIs for their special wildlife or geological interest or their physiographic features under Section 27 of the Wildlife Act 1990. This allows DEFA to work with the owners and occupiers, so that the conservation interest is maintained during the day to day operations that are necessary or occur on the land.

Notification and objection procedures

New sites are notified to the Department of Infrastructure and to every owner or occupier of the land and the legal provisions take effect immediately. A map of the site, a description of the special interest and a list of Operations Likely to Damage the special interest are included with the notification letter. A period of at least three months is then provided to allow representations and objections to be made. These may be resolved by liaison with a DEFA Biodiversity Officer. At the conclusion of this period the Department will consider such replies and within two months will decide whether to confirm, vary or rescind the notification. This decision will be notified to the consultees mentioned above, with reasons.

Ownership

Notification does not affect site ownership.

Farming and land use

Many sites are of importance to wildlife *because* they have been managed or farmed in a particular way for many years so it is vital that such management continues. Some sites may require slight modifications to their management. DEFA will discuss with owners and occupiers the present and future management of the ASSI, which may combine some production with nature conservation.

A management agreement can be proposed at any time, not just when a regime is changed or interest threatened.

Workers on site

Owners and occupiers should advise their employees, contractors and other authorised users of the existence of an ASSI, in order to avoid them unknowingly damaging the site. Statutory bodies such as the public utilities should be aware of them, but owners and occupiers are requested to inform them when approached for permission to carry out work on an ASSI. Owners and occupiers are not liable for the unauthorised action of third parties, such as trespassers and vandals.

Operations Likely to Damage the Special Interest

Once designated, the owner or occupier of a site retains responsibility for their land but they must provide notice to DEFA of specific operations that could affect the conservation interest. This sets off a dialogue with DEFA regarding conservation issues raised by such an operation.

A list of these 'Operations Likely to Damage the Special Interest' (OLDs) is provided with the notice of designation. This comprehensive list includes anything that could conceivably damage the interest. It may appear forbidding but this is not the case. These are not prohibited actions but DEFA **must** be given advance notice of such works and allowed time to respond and offer advice, or come to an agreement before the operation takes place. Where an agreement is offered by DEFA but not accepted, the matter may go to an arbitrator, agreed by both parties. The procedures can be found in Section 27 of the Wildlife Act 1990.

This system ensures that conservation is taken into account during works that might affect the interest of a site. The list usually covers the whole of the site, even though certain operations may only be possible or damaging on part of it. Certain listed operations may be required in specific parts of a site and can be done in a way that will not affect the special interest. By liaison between DEFA and a land owner or occupier, a management agreement may result which would not require the owner/occupier to give further notice for specified operations while the agreement is in effect.

Operations requiring notice may include those intended to conserve wildlife, if they are on the list of OLDs. Such actions can cause significant damage to the existing interest if carried out thoughtlessly or at the wrong time of year. Obtaining the consent of DEFA under the Wildlife Act does not remove the need to meet other legal obligations, such as tree felling licences or planning permission where relevant.

Planning applications

The presence of an ASSI will be taken into account when the Planning Committee considers applications affecting such a site. The outcome will depend on the nature of an application.

Financial provisions

No financial burden will be placed on an owner or occupier by virtue of notification, as no change in land use is required automatically and provisions exist for payments towards conservation management requested by either party.

If DEFA does not give consent for an operation, it may offer a management agreement including payments calculated according to the Financial Guidelines approved by Tynwald in 2001.

Further grants may be available through the Agri-Environment Scheme, for which an increased payment is available for ASSIs.

Access

Access is not normally affected by notification. Notification of a site does not give DEFA officers or anyone else a right of access. Existing rights of way remain and any statutory agencies with rights of

access, such as the public utilities, continue to have access but should be notified of the need to consult DEFA before undertaking any work in an ASSI.

Recreation

It is the responsibility of the owner or occupier to notify DEFA if recreational users are carrying out any of the notified operations. DEFA will, in many cases, permit existing levels of use to continue but will wish to discuss any proposals for new or significantly greater levels of use.

For further information contact:

Email: aline.thomas@gov.im


The Biodiversity Officer (Designations)
Department of Environment, Food and Agriculture
Forestry, Amenity and Lands Division
Thie Slieau Whallian
Foxdale Road
St Johns
IM4 3AS

Tel: 695737

Web address: www.gov.im/defa

This leaflet is produced for public awareness purposes only and is not a statement of law. The legislation covering ASSIs can be found in the Wildlife Act 1990.

The information in this leaflet can be provided in large print or on audio tape, on request.



Statutory Document No. 769/01

WILDLIFE ACT 1990

**AYRES GRAVEL PIT (BRIDE) AREA OF SPECIAL
PROTECTION FOR BIRDS ORDER 2001**

Laid before Tynwald: 15 January 2002

Coming into operation: 1 December 2001

In exercise of the powers conferred on the Department of Agriculture, Fisheries and Forestry by section 3 of the Wildlife Act 1990(a) and of all other enabling powers, and with the consent of the owner and occupier of the land to which this Order relates and after consultation with the Wildlife Committee and having given the relevant local authority the opportunity to submit objections or representations, the following Order is hereby made:-

Citation and commencement

1. This Order may be cited as the Ayres Gravel Pit (Bride) Area of Special Protection for Birds Order 2001 and shall come into operation on 1 December 2001.

Creation of the Ayres Gravel Pit (Bride) Area of Special Protection

2. The area of land (and water) edged with a thick black line on the plan in the schedule to this Order shall be an Area of Special Protection for birds.

3. Subject to section 3(2) and section 4 of the Wildlife Act 1990, any person who, within the area specified under article 2 of this Order intentionally -

- (a) kills, injures or takes any wild bird or game bird;
- (b) takes, damages or destroys the nest of such a bird while that nest is in use or being built;
- (c) takes or destroys an egg of such a bird;

- (c) takes or destroys an egg of such a bird;
- (d) disturbs such a bird while it is building a nest or is in, on or near a nest containing eggs or young; or
- (e) disturbs any nest or egg of such a bird; or
- (f) disturbs dependent young of such a bird,


shall be guilty of an offence under section 3 of the Wildlife Act 1990;

4. Subject to section 3(2) and (3) and section 4 of the Wildlife Act 1990, any person who at any time enters into the area specified under article 2 shall be guilty of an offence under section 3 of the Wildlife Act 1990, with the exception of-

- (a) any person or class of persons authorised in writing by the Department of Agriculture, Forestry and Fisheries;
- (b) persons supplying emergency services;
- (c) persons exercising a power of access or entry conferred by any statutory provision.

5. Where any offence has been committed under section 3 by virtue of this Order, the offender shall be liable to a special penalty.

MADE NOVEMBER 28TH 2001



Hon Alex F Downie MHK
Minister for Agriculture, Fisheries and Forestry

EXPLANATORY NOTE
(This is not part of the Order.)

This Order creates an Area of Special Protection for birds at the Ayres gravel pit, Bride. It protects the wild birds by making it an offence to do certain activities within the site, and by restricting access to specified people.



Schedule

**Ayres Gravel Pit (Bride) Area of Special
Protection for birds**

Site boundary (centre of line): —————

1:10 000

Grid North



Ballachurry Meadows Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys


Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin

Area: 11.9 hectares (29.4 acres)
National Grid Reference (centroid): SC 4025 9624




Ballachurry Meadows Area of Special Scientific Interest

Designated: 11th November 2009

Area: 11.9 hectares (29.4 acres)

Varied and confirmed: 24th May 2010

National Grid Reference (centroid): SC 4025 9624



Text

2009 Aerial Photograph

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Area: Isle of Man

Site name: Ballachurry Meadows

Parish: Lezayre

Status: Area of Special Scientific Interest (ASSI)
notified under Section 27 of the Wildlife Act 1990.

National Grid Reference: SC 4025 9624

Area: 11.9ha (29.4 acres)

Ordnance Survey Sheets: 1:50,000 : Map 95
1:25,000 Isle of Man Public Rights of Way and Outdoor Leisure Map (North)

Date notified: 17th November 2009

Date confirmed: 24th May 2010

Date of last revision: N/A

Purpose

This site is designated for its unimproved neutral grassland of the type, crested dog's-tail and knapweed *Cynosurus cristatus* - *Centaurea nigra* lowland grassland. This site comprises the majority of the 11.08ha (0.02% of the Island) that Sayle *et al* described as unimproved neutral grassland in the Phase 1 Habitat Survey report in 1995.

Description:

Ballachurry Meadows are a series of grasslands ranging from fairly dry hay meadows to very wet sedge-rich rushy meadows. They are on soils formed on glacial deposits. The meadows are species-rich and variously dominated by crested dog's tail *Cynosurus cristatus*, red fescue *Festuca rubra* and sweet vernal grass *Anthoxanthum odoratum*.

The site supports the largest population of greater butterfly orchids *Platanthera chlorantha**** on the Island and is important for pale sedge *Carex pallescens**** which is very rare on the Island. Four other orchids are also found on the site – northern marsh *Dactylorhiza purpurella****, heath spotted *D. maculata* subsp. *ericetorum****, common spotted *D. fuchsii**** and common twayblade *Listera ovata****.

Other floral interest includes whorled caraway *Carum verticillatum*, devil's bit scabious *Succisa pratensis*, common knapweed *Centaurea nigra*, tufted vetch *Vicia cracca*, red clover *Trifolium pratense*, selfheal *Prunella vulgaris*, meadow vetchling *Lathyrus pratensis*, angelica *Angelica sylvestris* and meadow buttercup *Ranunculus acris*. Royal fern (*Osmunda regalis*) has been recorded.

In the wetter meadows there is yellow flag iris *Iris pseudacorus*, marsh marigold *Caltha palustris*, lesser spearwort *Ranunculus flammula*, greater bird's foot trefoil *Lotus pedunculatus* and purple loosestrife *Lythrum salicaria*. There is also a small pond.

A good range of sedges are found in addition to pale sedge. They include carnation sedge *Carex panicea*, common sedge *C. nigra*, oval sedge *C. ovalis* and common yellow sedge *C. viridula* ssp. *oedocarpa*.

Birds either definitely or probably breeding on or very close to the site include Barn Owl (*Tyto alba*)*, Blackbird (*Turdus merula*), Blackcap (*Sylvia atricapilla*), Blue tit (*Cyanistes caeruleus*), Chaffinch (*Fringilla coelebs*), Coal tit (*Periparus ater*), Curlew (*Numenius arquata*)*, Dunnock (*Prunella modularis*), Goldcrest (*Regulus regulus*), Great Tit (*Parus major*), Robin (*Erithacus rubecula*), Song Thrush (*Turdus philomelos*)*, Spotted Flycatcher (*Muscicapa striata*)*, Swallow (*Hirundo rustica*), Whitethroat (*Sylvia communis*), Willow Warbler (*Phylloscopus trochilus*) and Wren (*Troglodytes troglodytes*).

Other information:

Full species lists are available from the Department of Environment, Food and Agriculture.

- * = species on Schedule 1 of the Wildlife Act 1990
- ** = species on Schedule 5 of the Wildlife Act 1990
- *** = species on Schedule 7 of the Wildlife Act 1990

= Red List Bird of Conservation Concern

Site Name: Ballachurry Meadows, Lezayre

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseeded.
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
26	Use of vehicles or craft likely to damage or disturb features of interest.
27	Recreational or other activities likely to damage features of interest or soil, fauna

	and flora.
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, amphibian, bird, fish or invertebrate.

Ballacrye Meadow Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmhtaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin



Department of Agriculture,
Fisheries and Forestry

**BALLACRYE MEADOW
AREA OF SPECIAL
SCIENTIFIC INTEREST**

Area: 0.55ha



Site Boundary

Designated: 7th February 2005


Confirmed: 11th July 2005

The grid lines on this map form
part of the National Grid,
and are spaced at 50m intervals

0 10 Meters



Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Ballacrye Meadow

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Jurby

Planning Authority: Department of Local Government and the Environment

Local authority: Jurby Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC 39SE & SC39NE

National Grid Reference (centroid): SC 38059754

Area: 0.55 ha (1.36 acres)

Date notified: February 7th 2005

Date confirmed: July 11th 2005

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its flora and habitat - species-rich rush pasture of the *Juncus acutiflorus* – *Galium palustre* (Sharp-flowered rush and Marsh bedstraw) type.

Description and reasons for notification:

Ballacrye Meadow is a species-rich rush pasture of the *Juncus acutiflorus* – *Galium palustre* (Sharp-flowered rush and Marsh bedstraw) type. Nearly all of the species typical of the best Manx examples of this habitat have been recorded from this site, giving it an unusually high level of diversity.

The vegetation is characterised by a high proportion of whorled caraway *Carum verticillatum*, which is abundant throughout. The meadow is bounded on two sides by thick hedges of willow, ash and other shrubs, and is consequently a sheltered site ideally suited to whorled caraway, which has a very limited range of ecological preferences. Whorled caraway meadows are a distinctive kind of fen meadow found in southwest England, southwest Scotland, west Wales, western Ireland and the Isle of Man. This meadow, although small in extent, is considered to be one of the best examples of the habitat type in the Isle of Man, both in terms of its composition and its sheltered setting. The thick woody hedges around the edge of the meadow have been included in the site boundary for this reason.

Managed as a hay meadow, the grassland also has some corresponding similarities to other hay meadows on the Island, including species typical of *Cynosurus cristatus* – *Centaurea nigra* (Crested dog's-tail – Black knapweed) neutral grassland, such as Yellow rattle *Rhinanthus minor*. In addition to a high overall diversity, this site is notable for the presence of large numbers of orchids, including Heath spotted-orchid *Dactylorhiza maculata*, Common spotted-orchid *Dactylorhiza fuchsii* and Northern marsh orchid *Dactylorhiza purpurella*, all of which are protected under Schedule 7 of The Wildlife Act 1990. Such species-rich examples of either fen meadows or hay meadows are known to be very restricted and fragmented on the Island, mostly occurring in small field corners or field margins.

Although a small site, this meadow qualifies for ASSI selection because it fulfils the Priority Site Criteria for diversity, naturalness, representativeness and rarity, for the reasons described above.

Other information:

Species-rich rush pastures are internationally threatened, having suffered very heavy losses due to agricultural improvement, drainage and development. Countries within the British Isles are considered to have a special responsibility for the protection of this habitat, which is in danger of extinction if the current European rate of loss continues. In view of this, "Purple moor-grass and Rush Pastures" are now a Priority Habitat in the UK Biodiversity Action Plan, and several key sites have been designated SSSIs and/or SACs in recent years. Examples with whorled caraway as a prominent component are particularly important as this species has a distinct western distribution in the British Isles and is reliant on a very limited range of ecological preferences.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Ballacrye Meadow

Operations likely to damage the special interest of the site

Type of operation

1. Cultivation including ploughing, rotovating, harrowing and reseeding;
2. Introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing) or continuation of grazing known to be damaging;
3. Supplementary stock feeding, where already damaging, or the introduction of supplementary stock feeding and changes in stock feeding practice;
4. Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing);
5. Changes in the application of manure, fertilisers and lime (including introduction of a new regime, and changing existing amounts and timing of application), or continuation of applications which are known to be damaging;
6. Application of pesticides, including herbicides (weedkillers);
7. Dumping, spreading or discharge of any materials;
8. Burning, and lighting of fires;
9. The release into the site of any wild, feral or domestic animals *, plant or seed (excluding domestic dogs);
10. The killing or removal of any wild animal * (excluding recognised game species in season);
11. The destruction, displacement, removal or cutting of any plant or plant remains, including herbs, dead or decaying wood, moss, lichen, fungus, leaf mould or turf, (excluding accepted management, i.e. hay cuts and control of statutory injurious weeds by non-chemical methods);
12. The introduction of tree and /or woodland management and changes in tree and/ or woodland management;
13. Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains);
14. The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes);
15. Infilling or other deliberate modification of ditches, dykes, drains, ponds, pools, marshes or pits;
16. Extraction of minerals including peat, sand and gravel, topsoil and subsoil;
17. Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground;
18. Storage of materials;
19. Erection of permanent or temporary structures, or the undertaking of engineering works;
20. Removal of geological specimens, including rock and soil samples, minerals and fossils.
21. Use of vehicles which are likely to damage or disturb features of interest;
22. Recreational or other activities likely to damage features of interest (excluding walking, with or without dogs).

* "animal" here includes any mammal, reptile, bird or invertebrate.

February 1st 2005

Ballateare Meadow Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.


Department of Environment, Food and Agriculture
Rheynn Chymmhtaght, Bee as Eirinys

This Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



Isle of Man
Government

Reilrys Ellan Vannin



Citation

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST

Site name: Ballateare Meadow (Consortium Field)

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Jurby

Planning Authority: Department of Infrastructure

Local Authority: Jurby Commissioners

National Grid Reference: SC 3627 9718

Area: 0.96 ha (2.37 acres)

Ordnance Survey Sheets: 1:50,000 OS Landranger Map 95 and 1:10,560 sheet SC 39 NE

Date notified: 21st May 2013

Date confirmed: 19th February 2014

Date of last revision: N/A

Purpose

This site is designated for its neutral grassland dominated by sweet vernal grass *Anthoxanthum odoratum* with Yorkshire fog *Holcus lanatus*.

Description

Species-rich hay meadow grades to a wet, sedge-dominated area on soils formed on glacial deposits. The meadow is dominated by sweet vernal grass *Anthoxanthum odoratum* and Yorkshire fog *Holcus lanatus*. The site supports greater butterfly orchids *Platanthera chlorantha****, northern marsh orchids *Dactylorhiza purpurella****, heath spotted orchids *D. maculata****, common spotted orchids *D. fuchsii**** and common twayblade *Listera ovata****.

Other floral interest includes whorled caraway *Carum verticillatum*, adder's-tongue fern *Ophioglossum vulgatum*, common knapweed *Centaurea nigra*, tufted vetch *Vicia cracca*, common vetch *Vicia sativa*, bush vetch *Vicia sepium*, red clover *Trifolium pratense*, common mouse-ear *Cerastium fontanum*, angelica *Angelica sylvestris*, ox-eye daisy *Leucanthemum vulgare* and meadow buttercup *Ranunculus acris*.

Marshy grassland dominated by sharp flowered rush *Juncus acutiflorus* occupies the northernmost quarter and a small area to the southeast. This area supports bulrushes *Typha latifolia*, branched bur-reed *Sparganium erectum*, meadowsweet *Filipendula ulmaria*, common bird's-foot trefoil *Lotus corniculatus* and greater bird's-foot trefoil *L. pedunculatus* together with a good range of sedges including carnation sedge *Carex panicea*, common sedge *C. nigra*, oval sedge *C. ovalis* and star sedge *C. echinata*.

Birds breeding at the site include curlew *Numenius arquata**, sedge warbler *Acrocephalus schoenobaenus*, whitethroat *Sylvia communis*, and willow warbler *Phylloscopus trochilus*.

* = species on Schedule 1 of the Wildlife Act 1990

*** = species on Schedule 7 of the Wildlife Act 1990

List of operations requiring DEFA's consent

The operations listed below may damage the features of interest of **Ballateare Meadow (Consortium Field) ASSI**. Before any of these operations are undertaken you must consult DEFA, and may require our consent.

It is usually possible to carry out some of these operations in certain ways, or at specific times of year, or on certain parts of the ASSI, without damaging the features of interest. If you wish to carry out any of these activities please contact a DEFA Biodiversity Officer who will give you advice and, where appropriate, issue a consent. Please help us by using the form sent to you with the notification documents to ask us for consent to carry out these operations.

In certain circumstances it will not be possible to consent these operations, because they would damage the features of interest. Where possible a Biodiversity Officer will suggest alternative ways in which you may proceed, which would enable a consent to be issued. To proceed without DEFA's consent may constitute an offence. If consent is refused, or conditions attached to it, which are not acceptable to you, you will be provided with details of how you may appeal to DEFA.

Standard reference number	Type of operation
1	Cultivation, including ploughing, rotovating, harrowing and re-seeding.
2	Grazing, the introduction of grazing and alterations to the grazing regime (including type of stock, intensity or seasonal pattern of grazing).
3	Stock feeding, the introduction of stock feeding and alterations to stock feeding practice.
4	Mowing or cutting of vegetation (where already damaging), the introduction of mowing and alterations to the mowing or cutting regime (such as from haymaking to silage).
5	Application of manure, slurry, silage liquor, fertilisers and lime.
6	Application of pesticides, including herbicides (weed killers) whether terrestrial or aquatic, and veterinary products.
7	Dumping, spreading or discharge of any materials.
8	Burning and alterations to the pattern or frequency of burning.
9	Release into the site of any wild, feral, captive-bred or domestic animal, plant, seed or micro-organism (including genetically modified organisms).
10	Killing, injuring, taking or removal of any wild animal (including dead animals or parts thereof), or their eggs and nests, including pest control and disturbing them in their places of shelter.
11	Destruction, displacement, removal or cutting of any plant or plant remains, including tree, shrub, herb, hedge, dead or decaying wood, moss, lichen, fungus, leaf-mould, and turf.
12	Tree and/or woodland management (where already damaging), the introduction of tree and/or woodland management (where applicable) and alterations to tree and/or woodland management (including planting, felling, pruning and tree surgery, thinning coppicing, changes in species composition, removal of fallen timber).
13 a)	Draining (including moor- gripping, the use of mole, tile, tunnel or other artificial drains).
13 b)	Modification to the structure of water courses (e.g. rivers, streams, springs, ditches and drains), including their banks and beds, as by re-alignment, regarding, damming, or dredging.
13 c)	Management of aquatic and bank vegetation for drainage purposes.
14	Alterations to water levels and tables and water utilisation (including irrigation, storage, abstraction from existing water bodies and through boreholes). Also the modification of current drainage practices.
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, hard rock, sand, gravel, topsoil, subsoil, lime, limestone pavement, shells and spoil.
21	Destruction, construction, removal, rerouting, or regrading of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, including soil and soft rock exposures or

	the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures or the undertaking of engineering works, including drilling.
24 a)	Modification of natural or man-made features (including cave entrances) and clearance of boulders, large stones, loose rock or scree.
26	Use of vehicles or craft.
27	Recreational or other activities.
28 a)	Game and waterfowl management and hunting practices (where already damaging), introduction of game or waterfowl management (where applicable) and alterations to game and waterfowl management and hunting practice.
28 b)	Use of lead shot.

Notes:

1. This is a list of operations appearing to DEFA to be likely to damage the special features of this ASSI, as required under Section 27 of the Wildlife Act 1990.
2. Any reference to 'animal' in this list shall be taken to include any mammal, reptile, amphibian, bird, fish or invertebrate.

Date notified: 21st May 2013

Date confirmed: 19th February 2014

Date of last revision: N/A

Views about Management

This is not a legal document.

A statement of DEFA's views about the management of Ballateare Meadow (Consortium Field) Area of Special Scientific Interest (ASSI)

This statement represents DEFA's views about the management of the ASSI for nature conservation. This statement sets out in principle, our views on how the area's special conservation interest can be conserved and enhanced. DEFA does not have a duty to notify the owners and occupiers of ASSI of its views about the management of the land, but DEFA has determined that this would be good practice with ASSI designations.

Not all of the management principles will be equally appropriate to all parts of the ASSI. Also, there may be other activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

This statement does not constitute consent for any of the 'operations requiring DEFA's consent'. The written consent of DEFA is required before carrying out any of those operations. DEFA welcomes consultation with owners, occupiers and users of the ASSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

Continuation of the hay meadow management will protect the existing natural habitats of interest on this site and control scrub encroachment. Late grazing or aftermath grazing by sheep should not cause a problem for the designated interest. Such management, avoiding agricultural improvement, will maintain the diverse grassland flora and suitable soil conditions for the rare plants. Farmyard manure could be applied with prior consent in order to maintain productivity, if necessary, or left off altogether if the aim is simply to manage for the wildlife interest, but artificial fertilisers may be damaging and should be avoided. The application of pesticides should be avoided generally but where necessary the Department will discuss issues arising and the most appropriate way of tackling them within the designated area.

Date notified: 21st May 2013

Date confirmed: 19th February 2014

Date of last revision: N/A

Information Sheet on Ramsar Wetlands (RIS)

Categories approved by Recommendation 4.7 (1990), as amended by Resolution VIII.13 of the 8th Conference of the Contracting Parties (2002) and Resolutions IX.1 Annex B, IX.6, IX.21 and IX. 22 of the 9th Conference of the Contracting Parties (2005).

Notes for compilers:

1. The RIS should be completed in accordance with the attached *Explanatory Notes and Guidelines for completing the Information Sheet on Ramsar Wetlands*. Compilers are strongly advised to read this guidance before filling in the RIS.
2. Further information and guidance in support of Ramsar site designations are provided in the *Strategic Framework for the future development of the List of Wetlands of International Importance* (Ramsar Wise Use Handbook 7, 2nd edition, as amended by COP9 Resolution IX.1 Annex B). A 3rd edition of the Handbook, incorporating these amendments, is in preparation and will be available in 2006.
3. Once completed, the RIS (and accompanying map(s)) should be submitted to the Ramsar Secretariat. Compilers should provide an electronic (MS Word) copy of the RIS and, where possible, digital copies of all maps.

1. Name and address of the Official Respondent:

Joint Nature Conservation Committee

Monkstone House
City Road
Peterborough
Cambridgeshire PE1 1JY
UK

Telephone/Fax: +44 (0)1733 – 562 626 / +44 (0)1733 – 555 948

Email: RIS@JNCC.gov.uk

FOR OFFICE USE ONLY.

DD MM YY

--	--	--

Designation date

--	--	--	--	--	--

Site Reference Number

Name and address of the compiler of this form:

UK Overseas Territories Conservation Forum, 102 Broadway, Peterborough, PE1 4DG, UK

Updated by Elizabeth Charter

Wildlife and Conservation Division

Department of Agriculture Fisheries and Forestry

Knockaloe

Peel

Isle of Man

IM5 3AJ

2. Date this sheet was completed/updated:

Designated: 06 September 2006

3. Country:

UK (Isle of Man)

4. Name of the Ramsar site:

Ballaugh Curragh

5. Designation of new Ramsar site or update of existing site:

This RIS is for: Designation of a new Ramsar site

6. For RIS updates only, changes to the site since its designation or earlier update:

a) Site boundary and area:

**** Important note:** If the boundary and/or area of the designated site is being restricted/reduced, the Contracting Party should have followed the procedures established by the Conference of the Parties in the Annex to COP9 Resolution IX.6 and provided a report in line with paragraph 28 of that Annex, prior to the submission of an updated RIS.

b) Describe briefly any major changes to the ecological character of the Ramsar site, including in the application of the Criteria, since the previous RIS for the site:

7. Map of site included:

Refer to Annex III of the *Explanatory Notes and Guidelines*, for detailed guidance on provision of suitable maps, including digital maps.

a) A map of the site, with clearly delineated boundaries, is included as:

- i) **hard copy** (required for inclusion of site in the Ramsar List): yes ✓ -or- no ☐;
- ii) **an electronic format** (e.g. a JPEG or ArcView image) Yes
- iii) **a GIS file providing geo-referenced site boundary vectors and attribute tables** yes ✓ -or- no ☐;

b) Describe briefly the type of boundary delineation applied:

e.g. the boundary is the same as an existing protected area (nature reserve, national park etc.), or follows a catchment boundary, or follows a geopolitical boundary such as a local government jurisdiction, follows physical boundaries such as roads, follows the shoreline of a waterbody, etc.

The site boundary is the same as, or falls within, an existing protected area.

For precise boundary details, please refer to paper map provided at designation

8. Geographical coordinates (latitude/longitude):

054 19 00 N 004 31 00 W

9. General location:

Include in which part of the country and which large administrative region(s), and the location of the nearest large town.

8 km west of Ramsey

Administrative region: Isle of Man

10. Elevation (average and/or max. & min.) (metres): **11. Area** (hectares): 193.4

Min. 10
Max. 20
Mean 15

12. General overview of the site:

Provide a short paragraph giving a summary description of the principal ecological characteristics and importance of the wetland.

The Ballaugh Curragh consists of a complex mosaic of interrelated peatland habitats dominated by grey willow *Salix cinerea* and birch *Betula pubescens* scrub (a habitat known locally as 'curragh'). Associated wetland habitats include bog pools, wet woodland, man-made ditch systems and fen grassland. The area supports a huge winter roost of hen harriers *Circus cyaneus* - at times the largest recorded in Europe. It has a very high diversity of breeding birds and a range of mire and aquatic plants including local rarities and species protected under Manx law. The peat deposits of the Ballaugh Curragh are also of importance for the historical pollen record that they provide. The biodiversity of the Ballaugh Curragh is known to have contributed to its cultural importance as an early settlement, and hence to its considerable value as an archaeological resource. Parts of the area are still farmed, principally for seasonal rough grazing and traditional hay cropping.

13. Ramsar Criteria:

Circle or underline each Criterion applied to the designation of the Ramsar site. See Annex II of the *Explanatory Notes and Guidelines* for the Criteria and guidelines for their application (adopted by Resolution VII.11).

1, 2, 3, 4

14. Justification for the application of each Criterion listed in 13 above:

Provide justification for each Criterion in turn, clearly identifying to which Criterion the justification applies (see Annex II for guidance on acceptable forms of justification).

1 The site consists of a representative example of a peatland habitat mosaic with a high degree of habitat diversity, dominated by a once-typical but now increasingly rare and fragmented willow scrub habitat, known locally as 'curragh'. Other habitats include wet birch woodland, wet grassland, purple moor-grass-dominated fen pasture with good populations of marsh orchids, floating mats of bog vegetation and well-established man-made ditch systems with good collections of aquatic and emergent species.

2 Corncrake *Crex crex* (SPEC1 – global conservation concern: Vulnerable) settled at Close Sartfield Nature Reserve in the Ballaugh Curragh in 1999, and definitely bred in 2000. The Curragh is known to have been one of the last strongholds for this bird prior to its dramatic population decline throughout the British Isles; the continuing presence of traditionally managed farmland in the area is now important for the return of the corncrake as a Manx breeding species.

The asilid fly *Epitriptus cowini* (RDB2 (Vulnerable), pRDBK) has been recorded from the Ballaugh Curragh area.

3 The Ballaugh Curragh supports the largest hen harrier *Circus cyaneus* winter roost on the Island, with a 5-year mean peak count of 82 (1996/7-2000/01); well over 100 may be seen. This represents a high proportion of all winter roosting hen harriers in the region; in some years it has been recorded as having the highest number in Western Europe.

The Ballaugh Curragh is one of the most biologically diverse sites on the Isle of Man, containing a very high diversity of breeding birds and diverse plantlife. For some rare and threatened plants the Curragh represents the only Manx station; the uncommon habitat types present are in some cases the most diverse examples of their kind on the Island. It is therefore of high importance for the conservation of Manx biodiversity.

4 Regionally important winter roost for hen harrier *Circus cyaneus* – see above.

15. Biogeography (required when Criteria 1 and/or 3 and /or certain applications of Criterion 2 are applied to the designation):

Name the relevant biogeographic region that includes the Ramsar site, and identify the biogeographic regionalisation system that has been applied.

a) biogeographic region:

Atlantic

b) biogeographic regionalisation scheme (include reference citation):

EU Habitats Directive 92/43/EEC (as amended)

16. Physical features of the site:

Describe, as appropriate, the geology, geomorphology; origins - natural or artificial; hydrology; soil type; water quality; water depth, water permanence; fluctuations in water level; tidal variations; downstream area; general climate, etc.

Soil & geology	alluvium; peat
Geomorphology and landscape	No information available
Nutrient status	oligotrophic
pH	acidic
Salinity	fresh
Soil	mainly organic
Water permanence	usually permanent
Summary of main climatic features	Cool oceanic lowland: Average rainfall 866 mm/year; Average min. temperature: 4.8°C Average max. temperature: 17.5°C

General description of the Physical Features:

No information available

17. Physical features of the catchment area:

Describe the surface area, general geology and geomorphological features, general soil types, general land use, and climate (including climate type).

Part of catchment is within cool oceanic upland climate, as the northern lowlands of the Isle of Man are fed by the Northern Hills, which rise to a maximum of 620 m. Water quality is generally good but acid, base-poor; water supply is relatively steady, although the small catchment area may result in temporary droughts in unusually dry summers. Land use in the catchment is overwhelmingly rural, mostly extensive pasture and moorland.

The Curragh represents a large storage area of water coming off the hills, particularly in winter when the wet pastures are generally flooded at least part of the time. The area has a complex hydrology and a full hydrological assessment is not available.

18. Hydrological values:

Describe the functions and values of the wetland in groundwater recharge, flood control, sediment trapping, shoreline stabilization, etc.

Flood water storage / desynchronisation of flood peaks

19. Wetland types:

Inland wetland

Code	Name	% Area
W	Shrub-dominated wetlands	60
U	Peatlands (including peat bogs swamps, fens)	30
4	Seasonally flooded agricultural land	9
9	Canals and drainage channels	1

20. General ecological features:

Provide further description, as appropriate, of the main habitats, vegetation types, plant and animal communities present in the Ramsar site, and the ecosystem services of the site and the benefits derived from them.

Ballaugh Curragh is the largest remaining intact example of a very distinctive Manx habitat which was historically important as a plentiful source of fish, waterfowl and willow. It therefore forms a culturally-defined area of the Island as well as a distinct wetland, and as such has provided valuable archaeological and palynological (historic pollen) records. Although dominated by willow scrub on peat, the Curragh also has open water, damp neutral to acidic grassland, marshy grassland, dry scrub, hedges, birch woodland with heather, modified and wet bog, tall ruderal and swamp vegetation, semi-natural broadleaved woodland and semi-improved pasture. Of particular significance in the regional context is the presence of habitats on Annex I of the EU Habitats Directive (wet bog woodland, natural dystrophic ponds, *Molinia* meadows on peaty soils, and UK Biodiversity Action Plan Priority Habitats (wet woodland, purple moor-grass and rush pastures, fens, species-rich hedges).

The Curragh is well known locally for its wetland habitats and bird diversity. The presence of impressive winter hen harrier numbers and traditionally-managed meadows with summer displays of thousands of orchids attracts visitors both for bird-watching and the popular 'orchid tours' run by the Manx Wildlife Trust.

21. Noteworthy flora:

Provide additional information on particular species and why they are noteworthy (expanding as necessary on information provided in 12. Justification for the application of the Criteria) indicating, e.g. which species/communities are unique, rare, endangered or biogeographically important, etc. *Do not include here taxonomic lists of species present – these may be supplied as supplementary information to the RIS.*

Rare and/or vulnerable plants (protected under Schedule 7 of the Wildlife Act 1990 as amended)

recorded in the past two years:

Common spotted-orchid *Dactylorhiza fuchsii*
 Heath spotted-orchid *Dactylorhiza maculata* spp ericetorum and hybrids
 Northern marsh-orchid *Dactylorhiza purpurella*
 Common Twayblade *Listera ovata*
 Greater butterfly-orchid *Platanthera chlorantha*
 Lesser tussock-sedge *Carex diandra*

Recorded in the past 50 years but not recently:

Pillwort *Pilularia globulifera*
 Bladderwort *Utricularia vulgaris*
 Greater spearwort *Ranunculus lingua*
 Blunt-leaved pondweed *Potamogeton obtusifolius*
 Fragrant orchid *Gymnadenia conopsea*
 Lesser water-plantain *Baldellia ranunculoides*

22. Noteworthy fauna:

Provide additional information on particular species and why they are noteworthy (expanding as necessary on information provided in 12. Justification for the application of the Criteria) indicating, e.g. which species/communities are unique, rare, endangered or biogeographically important, etc., including count data. *Do not include here taxonomic lists of species present – these may be supplied as supplementary information to the RIS.*

Species Information

(Provisional list)

Breeding birds protected under Schedule 1 of the Wildlife Act 1990 (as amended) include:

Water rail *Rallus aquaticus*
 Corncrake *Crex crex* (not regular)
 Northern lapwing *Vanellus vanellus*
 Eurasian curlew *Numenius arquata*
 Song thrush *Turdus philomelos*
 Common grasshopper warbler *Locustella naevia*

Non-breeding birds protected under Schedule I of the Wildlife Act 1990 (as amended) include:

Little grebe *Tachybaptus ruficollis*
 Whooper swan *Cygnus cygnus*
 Northern shoveler *Anas clypeata*
 Northern pintail *Anas acuta*
 Hen harrier *Circus cyaneus*
 Eurasian sparrowhawk *Accipiter nisus*
 Common kestrel *Falco tinnunculus*
 Long-eared owl *Asio otus*
 Short-eared owl *Asio flammeus*
 Common kingfisher *Alcedo atthis*
 Spotted flycatcher *Musciapa striata*
 House sparrow *Passer domesticus*

23. Social and cultural values:

Describe if the site has any general social and/or cultural values e.g. fisheries production, forestry, religious importance, archaeological sites, social relations with the wetland, etc. Distinguish between historical/archaeological/religious significance and current socio-economic values.

Aesthetic
 Archaeological/historical site
 Livestock grazing
 Non-consumptive recreation
 Tourism

b) Is the site considered of international importance for holding, in addition to relevant ecological values, examples of significant cultural values, whether material or non-material, linked to its origin, conservation and/or ecological functioning?

If Yes, tick the box ☐ and describe this importance under one or more of the following categories:

- i) sites which provide a model of wetland wise use, demonstrating the application of traditional knowledge and methods of management and use that maintain the ecological character of the wetland:
- ii) sites which have exceptional cultural traditions or records of former civilizations that have influenced the ecological character of the wetland:
- iii) sites where the ecological character of the wetland depends on the interaction with local communities or indigenous peoples:
- iv) sites where relevant non-material values such as sacred sites are present and their existence is strongly linked with the maintenance of the ecological character of the wetland:

24. Land tenure/ownership:

Ownership category	On-site	Off-site
Non-governmental organisation (NGO)	+	+
Local authority, municipality etc.	+	
National/Crown Estate	+	+
Private	+	+

25. Current land (including water) use:

Activity	On-site	Off-site
Tourism	+	+
Recreation	+	+
Arable agriculture (unspecified)		+
Permanent arable agriculture		+
Rough or shifting grazing	+	+
Permanent pastoral agriculture	+	+
Hay meadows	+	
Non-urbanised settlements		+

26. Factors (past, present or potential) adversely affecting the site's ecological character, including changes in land (including water) use and development projects:

Explanation of reporting category:

1. Those factors that are still operating, but it is unclear if they are under control, as there is a lag in showing the management or regulatory regime to be successful.
2. Those factors that are not currently being managed, or where the regulatory regime appears to have been ineffective so far.

NA = Not Applicable because no factors have been reported.

Adverse Factor Category	Reporting Category	Description of the problem (Newly reported Factors only)	On-Site	Off-Site	Major Impact?

What measures have been taken / are planned / regulatory processes invoked, to mitigate the effect of these factors?
Drainage/land-claim for agriculture - Controlled through designation.

Is the site subject to adverse ecological change? NO

27. Conservation measures taken:

List national category and legal status of protected areas, including boundary relationships with the Ramsar site; management practices; whether an officially approved management plan exists and whether it is being implemented.

a) List national and/or international category and legal status of protected areas, including boundary relationships with the Ramsar site:

In particular, if the site is partly or wholly a World Heritage Site and/or a UNESCO Biosphere Reserve, please give the names of the site under these designations.

b) If appropriate, list the IUCN (1994) protected areas category/ies which apply to the site (tick the box or boxes as appropriate):

Ia ☐; Ib ☐; II ☐; III ☐; IV ☒; V ☐; VI ☐

c) Does an officially approved management plan exist; and is it being implemented?:

Management plan in preparation

d) Describe any other current management practices:

Conservation measure	On-site	Off-site
Site/ Area of Special Scientific Interest (SSSI/ASSI)	+	
Land owned by a non-governmental organisation for nature conservation	+	
Other	+	
Management plan in preparation	+	
IUCN (1994) category IV	+	

b) Describe any other current management practices:

28. Conservation measures proposed but not yet implemented:

e.g. management plan in preparation; official proposal as a legally protected area, etc.

Management plan in preparation - Other parts of the site are subject to management agreements either as part of or comparable to agri-environment scheme management plans; in both cases there is a financial incentive

29. Current scientific research and facilities:

e.g. details of current research projects, including biodiversity monitoring; existence of a field research station, etc.

Nature reserves in the management of the Manx Wildlife Trust are subject to monitoring; the Curragh area is also subject to bird monitoring by the Manx Bird Atlas.

Current level of archaeological research unknown.

30. Current communications, education and public awareness (CEPA) activities related to or benefiting the site:

e.g. visitor centre, observation hides and nature trails, information booklets, facilities for school visits, etc.

A variety of interpretive material is produced by Manx National Heritage and the Manx Wildlife Trust.

There may be further interpretation at the Wildlife Park which has a nature trail through its part of the Curragh.

31. Current recreation and tourism:

State if the wetland is used for recreation/tourism; indicate type(s) and their frequency/intensity.

Limited public access for walking;

Popular annual 'orchid tours' run by Manx Wildlife Trust at its Close Sartfield nature reserve;

Public access and birdwatching hide at Close Sartfield;

Minor roads through the area used for road racing once a year (Manx International Rally).

32. Jurisdiction:

Include territorial, e.g. state/region, and functional/sectoral, e.g. Dept. of Agriculture/Dept. of Environment, etc.

Isle of Man Government

Wildlife and Conservation Division

Department of Agriculture, Fisheries and Forestry

Knockaloe

Peel

Isle of Man

IM5 3AJ

33. Management authority:

Provide the name and address of the local office(s) of the agency(ies) or organisation(s) directly responsible for managing the wetland. Wherever possible provide also the title and/or name of the person or persons in this office with responsibility for the wetland.

Wildlife and Conservation Division

Department of Agriculture, Fisheries and Forestry

Knockaloe

Peel

Isle of Man

IM5 3AJ

34. Bibliographical references:


Scientific/technical references only. If biogeographic regionalisation scheme applied (see 15 above), list full reference citation for the scheme.

Site-relevant references

Fairhurst, J (1998) *Ballaugh Curragh Management Plan*. Manx National Heritage, Douglas (unpublished)

- Falk, SJ (1991) *A review of the scarce and threatened bees, wasp and ants of Great Britain*. Nature Conservancy Council, Peterborough (Research and Survey in Nature Conservation, No. 35)
- Hepburn, I, Oldfield, S & Thompson, K (1992) *UK Dependent Territories Ramsar study: Stage 1*. Unpublished report to Department of the Environment, European and International Habitat Protection Branch, Bristol, from International Waterfowl and Wetlands Research Bureau/ NGO Forum for Nature Conservation in UK Dependent Territories, Slimbridge/ Sandy (Research contract, No. 7/2/126)
- Shirt, DB (ed.) (1987) *British Red Data Books: 2. Insects*. Nature Conservancy Council, Peterborough
- Pienkowski, MW (ed.) (2005) *Review of existing and potential Ramsar sites in UK Overseas Territories and Crown Dependencies*. (Contractor: UK Overseas Territories Conservation Forum, Peterborough.) Final report on Contract CR0294 to the UK Department for Environment, Food and Rural Affairs, Bristol. www.ukotcf.org
- Pritchard, DE, Housden, SD, Mudge, GP, Galbraith, CA & Pienkowski, MW (eds.) (1992) *Important Bird Areas in the United Kingdom including the Channel Islands and the Isle of Man*. Royal Society for the Protection of Birds, Sandy
-

Please return to: **Ramsar Secretariat, Rue Mauverney 28, CH-1196 Gland, Switzerland**
Telephone: +41 22 999 0170 • Fax: +41 22 999 0169 • email: ramsar@ramsar.org



**Isle of Man
Government**

Railtys Elian Vannin

**BALLAUGH CURRAGH
RAMSAR SITE
Site No. UK210001**

SITE PLAN

Date of designation: 6th September 2006

Total area: 193.36 hectares (477.79 acres)

Location (Centroid latitude/longitude):
54d 19m 0s N, 4d 31m 0s W

Altitude range: 5 - 15m above mean sea level

- Line detail**
- Building Outline
 - Building Pecked
 - Parish Boundary
 - District Boundary
 - Island Boundary
 - Electoral Boundary
 - Railway
 - Road Metalling
 - General Detail
 - General Peck Detail
 - Underground Detail
 - Veg/Landform Limit (suppressed)
 - Veg/Landform Limit
 - Overhead Detail
 - Minor Detail
 - Water Feature
 - Mean High Water
 - Mean Low Water
 - Top of Slope
 - Top of Cliff
 - Bottom of Slope or Cliff
 - Ramsar Site boundary

100 0 100 200 Meters



The grid lines on this map form
part of the UK National Grid
and are spaced at 500m intervals

Background map: Crown Copyright
Department of Local Government
and the Environment, Isle of Man
Unauthorised production
infringes copyright



**Isle of Man
Government**
Keillogh Ellen Vannin

**BALLAUGH CURRAGH
RAMSAR SITE
Site No. UK210001**

LOCATION MAP


Date of designation: 6th September 2006

Total area: 193.36 hectares (477.79 acres)

Location (centroid latitude/longitude):
54° 19' 0s N, 4° 31' 0s W

Altitude range: 5 - 15m above mean sea level

-  Ramsar Site boundary
-  Mountains
-  Rivers
-  Towns
-  Roads



Background map: Crown Copyright
Department of Local Government
and the Environment, Isle of Man
Unauthorised production
infringes copyright

Ballaugh Curragh Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmhtaght, Bee as Eirinys

This Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Reillys Ellan Vannin



Department of Agriculture,
Fisheries and Forestry

BALLAUGH CURRAGH AREA OF SPECIAL SCIENTIFIC INTEREST

and PROPOSED
RAMSAR SITE

Area: 193.4ha (\approx 477.9 acres)

Designated: 14th March 2005

Confirmed and varied:
9th September 2005

Ballaugh Curragh ASSI


□ Designation boundary
XX Excluded area

The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

0 100 Meters



Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Area: Isle of Man

Parishes: Ballaugh and Lezayre

Site name: BALLAUGH CURRAGH

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.
Designated as a Ramsar Site on 6th September 2006.

Planning Authority: Department of Local Government and the Environment

Local authority: Ballaugh Commissioners and Lezayre Commissioners

National Grid Reference: SC 367950 (approx. centroid) **Area:** 193.4 ha. (= 477.9 acres)

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95
1:10,560 OS sheets SC 39SE and SC39NE

Date notified: 14th March 2005

Varied and confirmed: 9th September 2005

Date of last revision: 6th September 2006

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its fauna, flora, geology, geomorphology and landscape.

Description and reasons for notification:

The Ballaugh Curragh consists of a diverse mosaic of interrelated wetland habitats which has evolved over a long period of time in response to the gradual colonisation of the site of the ancient lake known as Lake Andreas. The whole area has been the subject of traditional farming and fishing practices since the earliest human settlements on the Island, and the grassland and ditch system ecology reflects this long history. Habitats present include the largest area of wet modified bog on the Isle of Man, the best species-rich ditch system on peat soil, the largest area of continuous wet scrub (curragh), and good examples of wet mesotrophic grassland and traditional hay meadows, wet woodland, fen meadows and bog pools.

The area has long been noted for its diversity of bird life, with at least 39 breeding species. These include RSPB Red List birds such as lapwing *Vanellus vanellus** and corncrake *Crex crex**. The area also supports wintering birds, notably one of the largest wintering hen harrier *Circus cyaneus** roosts in Europe. Other fauna of note include aquatic invertebrates in the bog, pools and drainage ditches, fish including eel *Anguilla anguilla*, and a thriving common frog *Rana temporaria*** population.

The majority of the core Curragh area is dominated by grey sallow *Salix cinerea* scrub, with an understorey of wet woodland groundflora and, in more open areas, wet modified bog with occasional pools of standing water. Further out from the central core of the wetland is a series of traditionally-managed grasslands and hay meadows. Plants of note include good populations of spotted orchids *Dactylorhiza* spp*** and greater butterfly orchid *Platanthera chlorantha****, lesser tussock-sedge *Carex diandra****, bog myrtle *Myrica gale*, hare's-tail cotton-grass *Eriophorum vaginatum*, bog asphodel *Narthecium ossifragum*, whorled caraway *Carum verticillatum* and royal fern *Osmunda regalis*. Many rare and locally occurring plants have been recorded in the area in the past, for which new research is required to confirm their continuing presence on the site; these include pillwort *Pilularia globulifera****, bladderwort *Utricularia vulgaris****, greater spearwort *Ranunculus lingua****, whorled water-milfoil *Myriophyllum verticillatum* and blunt-leaved pondweed *Potamogeton obtusifolius****.

* listed in Schedule 1 of Wildlife Act 1990

** listed in Schedule 5 of Wildlife Act 1990

*** listed in Schedule 7 of Wildlife Act 1990

Other information:

The central core of the Ballaugh Curragh wetland was acquired by the Government in 1963 by means of an Act of Tynwald, the Curraghs Acquisition Act. Much of this area has since passed to the Manx Museum, and is managed not just for its ecological importance but also as a unique cultural and archaeological resource.

This area forms a cohesive ecological unit which is important for its terrestrial ecology and meets the criteria for designation as a wetland of international importance under the Ramsar Convention.

Full species lists are available from the Biological Records Centre, Manx Museum, Douglas.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: BALLAUGH CURRAGH

Operations likely to damage the special interest of the site

Type of operation

1. Cultivation including ploughing, rotovating, harrowing and reseeded;
 2. Introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing) or continuation of grazing known to be damaging;
 3. Supplementary stock feeding, where already damaging, or the introduction of supplementary stock feeding and changes in stock feeding practice;
 4. Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing);
 5. Changes in the application of manure, fertilisers and lime (including introduction of a new regime, and changing existing amounts and timing of application), or continuation of applications which are known to be damaging;
 6. Application of pesticides, including herbicides (weedkillers);
 7. Dumping, spreading or discharge of any materials;
 8. Burning, and lighting of fires;
 9. The release into the site of any wild, feral or domestic animals *, plant or seed (excluding domestic dogs);
 10. The killing or removal of any wild animal * (excluding recognised game species in season);
 11. The destruction, displacement, removal or cutting of any plant or plant remains, including herbs, dead or decaying wood, peat, moss, lichen, fungus, leaf mould or turf, (excluding accepted management, i.e. hay cuts and control of statutory injurious weeds by non-chemical methods);
 12. The introduction of tree and /or scrub management and changes in tree and/ or scrub management, including new planting;
 13. Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains);
 14. Modification of the structure of watercourses, including beds and banks, e.g. by re-alignment, re-grading, and dredging;
 15. Management of aquatic and bank vegetation for drainage purposes (also see 11);
 16. The changing of water levels and tables and water utilisation (including irrigation, increased discharges into ditches and watercourses on the site, storage, and abstraction);
 17. Infilling or other deliberate modification of the structure of watercourses, ditches, dykes, drains, ponds, pools, marshes or pits;
 18. Extraction of minerals including peat, sand and gravel, topsoil and subsoil;
 19. Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground;
 20. Storage of materials;
 21. Erection of permanent or temporary structures, or the undertaking of engineering works;
 22. Removal of geological specimens, including rock and soil samples, minerals and fossils.
 23. Use of vehicles which are likely to damage or disturb features of interest;
 24. Recreational or other activities likely to damage features of interest.
- * "animal" here includes any mammal, reptile, bird, fish or invertebrate.

Central Ayres Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin




**Isle of Man
Government**

Yn Yddir, a' Pen Yn Yddir

Department of Agriculture,
Fisheries and Forestry
**CENTRAL AYRES
AREA OF SPECIAL
SCIENTIFIC INTEREST**

Central Ayres ASSI

 2008 ASSI boundary

Area*: 358.34ha (885.46 acres)

O/S Grid reference: NX430035
(approximate centre of site)

Designated: 1/8/1996

Varied: 1/12/2008

Confirmed 17/10/2008


*Note: the seaward boundary of
the site is here defined as the
Lowest Astronomical Tide
between points A and B.

The grid lines on this map form
part of the National Grid,
and are spaced at 1km intervals

Base Map is

Crown Copyright

Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



VARIATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Central Ayres ASSI

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parishes: Andreas & Bride

Planning Authority: Department of Local Government and the Environment

Local authorities: Bride and Andreas Commissioners

National Grid Reference (centroid): NX430035 **Area:** 358.34ha (885.46 acres) in total, of which the extension to the previously notified area = 98.68ha (243.84 acres).

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets NX40SW & NX40SE

Date notified: 1st August 1996

Date of last revision: 11th February 2008

Date of variation confirmation: 17th October 2008

Purpose

The purpose of this designation is to extend the protection of an area which is important for its coastal habitats: intertidal shingle, gravel and sand, vegetated shingle and decalcified fixed dunes with dune heath and grassland. The area also supports important populations of rare and protected breeding and visiting wild birds, nationally threatened and vulnerable invertebrates¹ and rare and protected plants including one nationally vulnerable species². This Notification applies to the old designation boundary plus an extended area of approximately 98ha, which includes the intertidal zone adjacent to the old boundary, plus contiguous areas of dune heath and grassland. These were not included at the time of the previous designation, but are now known to have a high wildlife importance and form an integral part of the same ecosystem. This is therefore a formal revision, replacing the previous Central Ayres ASSI notification.

Description and reasons for notification:

The Central Ayres Area of Special Scientific Interest comprises the central part of an extensive area of coastal shingle and sand dunes. The dunes nearest to the sea are relatively mobile and rise to 5 metres in height, but have been wind-blown onto the landward side to form flat, fixed dune areas with occasional small depressions, and further back in the heath. Decalcification of the sand on the fixed dune areas has led to a unique vegetation of low herbs and shrubs ranging from almost-bare, lichen-dominated ground through to dwarf shrub heath. To the south of the site are wet hollows with seasonal pools. The extended area of designation comprises the intertidal elements of the shingle and gravel coast adjacent to the existing ASSI, and a larger area of fixed dune grassland and heath on the south-eastern edge of the site.

Intertidal zone and shingle

The main intertidal habitats identified from the site are barren shingle or gravel shore, and burrowing amphipods *Pontocrates* spp. and *Bathyporeia* spp. in lower shore clean sand. Areas of ephemeral algae and edible mussels have also been identified³. "Intertidal edible mussel beds on mixed and sandy sediments" is on the OSPAR Commission list of threatened and/or declining species and habitats.

The importance of the intertidal zone adjacent to the existing designated area relates to its ecological position. In addition to providing the source of sand for the dune system, the intertidal sands and shallow water provide a feeding area for birds that visit and/or breed on the site, including species which rely on the local sand eels, *Ammodytes tobianus* and *Hyperoplus lanceolatus*. Regular bird visitors known to feed on and beyond the intertidal area at high tide include a wide range of coastal species such as gannet *Sula bassana*, shag *Phalacrocorax aristotelis*, cormorant *Phalacrocorax carbo*, red-throated, black-throated and great northern divers *Gavia stellata*, *G. arctica*, *G. immer*, arctic tern *Sterna paradisaea**, sandwich tern *Sterna sandvicensis**, little tern *Sternula albifrons** (see below) and guillemot *Uria aalge*, razorbill *Alca*

torda and waders such as golden plover *Pluvialis apricaria**, curlew *Numenius arquata**, dunlin *Calidris alpina*, sanderling *Calidris alba*, common sandpiper *Actitis hypoleucos** turnstone *Arenaria interpres* oystercatcher *Haematopus ostralegus* and ringed plover *Charadrius hiaticula*. Black-tailed godwit *Limosa limosa*#, whimbrel *Numenius phaeopus*, redshank *Tringa tetanus*, knot *Calidris canutus* and lapwing *Vanellus vanellus** are amongst the more occasional visitors.

The shingle above the high-tide mark supports sparse strandline vegetation including sea sandwort *Honckenya peploides*, sea beet *Beta vulgaris* subsp. *maritima*, common orache *Atriplex patula*, Babington's orache *Atriplex glabriuscula*, spear-leaved orache *Atriplex prostrata*, yellow horned-poppy *Glaucium flavum*, sea-rocket *Cakile maritima*, Ray's knotgrass *Polygonum oxyspermum*, sea holly *Eryngium maritimum* and the nationally scarce, occasional rarity oysterplant *Mertensia maritima****. This area of shingle is of importance for breeding arctic tern *Sterna paradisaea** and little tern *Sternula albifrons**, oystercatcher *Ostralegus haematopus* and ringed plover *Charadrius hiaticula**, as well as providing a feeding area for a range of birds including starling *Sturnus vulgaris*#. The Ayres coast is the only place on the Island where little terns are known to breed regularly and numbers have reached 50 pairs or more, i.e. over 1% of the total British population. The shingle beach is an occasional haul-out site for grey seals *Hallochoerus grypus* ** which are regularly seen feeding in the shallow waters.

Dunes

Mobile dunes on the seaward side have a typical dune flora dominated by marram *Ammophila arenaria* and sand couch *Elytrigia juncea*, with Portland spurge *Euphorbia portlandica****, sea spurge *Euphorbia paralias*, sea bindweed *Calystegia soldanella* and sea holly *Eryngium maritimum*. Newer areas of fixed dune support a lime-tolerant flora including rest-harrow *Ononis repens*, bird's-foot trefoil *Lotus corniculatus*, eyebrights *Euphrasia* spp, sea storksbill *Erodium maritimum*, wild thyme *Thymus polytrichus*, early purple orchid *Orchis mascula*, moonwort *Botrychium lunaria*, mouse-ear hawkweed *Pilosella officinalis* and abundant pyramidal orchids *Anacamptis pyramidalis****.

Further inland the flat area of fixed dunes has an extensive zone of low vegetation and occasional bare ground, with low to short vegetation including wetland species in the wet depressions or "slacks". Dune slack plants of interest include adder's-tongue fern *Ophioglossum vulgatum*, dwarf yellow-sedge *Carex viridula* subsp. *viridula*, northern marsh orchid *Dactylorhiza purpurella****, allseed *Radiola linoides*, chaffweed *Anagallis minima*, twayblade *Listera ovata**** and early marsh orchid *Dactylorhiza incarnata****. This area is a feeding area for wading birds such as snipe in winter.

Fixed dune heath and grassland

Dunes which are well-established and subject to heavy leaching by high rainfall may develop into decalcified fixed dunes, which support a neutral to acid-loving flora and fauna despite their origins in lime-rich shell sand. The heavy rainfall and mild climatic conditions of the Atlantic coasts of Europe give rise to *Atlantic decalcified fixed dunes*, an internationally important Priority Habitat under Annex 1 of the EU Habitats Directive. The well-established fixed dunes at the Central Ayres ASSI largely fit this category, and are notable for a low-growing, wind-pruned dwarf shrub heath with a high proportion of lichen cover. Shrubby areas are dominated by ling *Calluna vulgaris*, bell heather *Erica cinerea*, Western gorse *Ulex gallii* and frequent patches of very low-growing burnet rose *Rosa spinosissima*.

The shortest vegetation on the fixed dune areas is ideal for the growth of lower plants, in particular lichens and mosses. The ground is almost bare in places and is dominated by lichens, in particular the grey lichens *Cladonia portentosa*, *C. arbuscula*, *C. fimbriata*, *C. floerkeana*, *C. uncialis* and *Hypogymnia physodes*. The latter species grows here not on trees but by colonizing other lichens and old heather stalks. The shortest areas are also notable for the presence of abundant "sausage lichen" *Usnea articulata*, which is only locally distributed in the British Isles and usually confined to growing in trees. The proportion of each species of lichen varies across the site, depending on the degree of shading by taller species, level of disturbance, and acidity of the substrate. These lichens have helped to create an exceptionally structurally diverse and species-rich example of the fixed dune habitat. Other lower plants of interest include the moss *Racomitrium ericoides*, and the uncommon winter stalkball fungus *Tulostoma brumale*.

Dune grassland occurs amongst the heath, particularly in fields along the southern edge of the site where there is grazing by rabbits and occasionally livestock. In addition to a diverse range of typical coastal

grassland plants, rarities occur such as Autumn lady's tresses *Spiranthes spiralis**** and field gentian *Gentianella campestris***. These are Near Threatened and Nationally Vulnerable² species respectively, on account of severe population decline throughout the British Isles in recent years. Small ponds occur within the grassland and heath areas, adding to the habitat diversity.

The fixed dunes as a whole offer an excellent habitat for a range of birds and animals, some of which occur nowhere else on the Island. Breeding birds of interest in these dunes and associated scrub include skylark *Alauda arvensis*, eider *Somateria mollissima*, linnet *Carduelis cannabina*#, yellowhammer *Emberiza citrinella***, stonechat *Saxicola torquata* and reed bunting *Emberiza schoeniclus*. Birds which feed over the dune heath and grassland include skylark *Alauda arvensis***, peregrine *Falco peregrinus**, hen harrier *Circus cyaneus**, merlin *Falco columbarius* and long-eared owls *Asio otus** which breed in the small plantation. Animals of note include the common frog *Rana temporaria*** and viviparous lizard *Lacerta vivipara***, and a great diversity of invertebrates including all but one of the Island's 15 native butterfly species. Of particular note are the scarce crimson and gold moth *Pyrausta sanguinalis*** which is classified as a Red Data Book Endangered species¹, and the heath bee-fly *Bombylius minor***, classified as Vulnerable. There are also 26 Nationally Scarce invertebrates recorded on the Ayres.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 5 of the Wildlife Act 1990

*** = Protected under Schedule 7 of the Wildlife Act 1990

= Birds of Conservation Concern Red List species⁴

Other information:

This designation refers to a variation of the original Ayres National Nature Reserve and ASSI boundary, which was drawn up before the present-day ASSI criteria were written, and followed administrative boundaries above the high tide mark as well as habitat considerations. The omission of the intertidal zone – and of areas of dune heath contiguous with the designation boundary – is now seen as a major shortcoming of the original designation. The current policy follows that of the UK⁵, which is to include adjacent related habitat such as the intertidal zone, where that habitat is vital for the support of features for which the site has been designated.

¹ Shirt, D.B. (editor) (1987). British Red Data Books: 2 Insects. Peterborough: Nature Conservancy Council.

² Cheffings, C. and Farrell, L. (Editors), (2005), The Vascular Plant Red Data List for Great Britain, ISSN 1473-0154

³ Holt, TJ (1999), An Intertidal Survey of the Isle of Man. Manx Wildlife Trust

⁴ Gregory Richard D, Wilkinson Nicholas I, Noble David G, Robinson James A, Brown, Andrew F, Hughes, Julian, Proctor Deborah, Gibbons David W, and Galbraith Colin A. (2002) The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002-2007. British Birds Vol 95 No 9 Pages 410-448

⁵ NCC (1989), *Guidelines for the Selection of Biological ASSIs* p35 S5.7. Peterborough: Nature Conservancy Council.

VARIATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Central Ayres ASSI

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17	Reclamation of land from sea, estuary or marsh.
18	Bait digging on inter-tidal areas, (if it is shown to be damaging).
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of

	pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling,.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, amphibian, bird, fish or invertebrate.

Cronk e King Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmhtaght, Bee as Eiriny's

The Slieu Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS


www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin


227000



479500

479500

227000



Citation

Site name: Cronk e King

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Patrick

Planning Authority: Department of Infrastructure

Local Authority: Patrick Commissioners

National Grid Reference: SC 2697 7960

Area: 3.02ha (=7.46 acres)

Ordnance Survey Sheets: 1:50,000 OS Landranger Map 95 and 1:10,560 sheet SC 28 SE

Date notified: 21st October 2013

Date confirmed: 19th February 2014

Date of last revision: N/A

Purpose

This site is designated for its species-rich neutral grassland dominated by sweet vernal grass *Anthoxanthum odoratum*.

Description

Cronk e King is comprised of five hay meadows that have received traditional hay meadow management consistently for many years. There is a narrow strip of wet, sedge-rich land to the upper north-westerly boundary. The meadows are dominated by sweet vernal grass *Anthoxanthum odoratum* and a number of plants which are indicative of a species-rich meadow. The site supports northern marsh *Dactylorhiza purpurella*^{***}, heath spotted *D. maculata*^{***} and common spotted orchids *D. fuchsii*^{***}. Other floral interest includes yellow rattle *Rhinanthus minor*, greater bird's-foot trefoil *Lotus pedunculatus*, common bird's-foot trefoil *Lotus corniculatus*, common sorrel *Rumex acetosa*, heath bed straw *Galium saxatile* and meadow buttercup *Ranunculus acris*.

The marshy grassland supports sedges including oval sedge *Carex ovalis* with rushes such as toad rush *Juncus bufonius* and sharp-flowered rush *Juncus acutiflorus*. Marsh thistles *Cirsium palustre* provide nectar for visiting butterflies including dark green fritillaries *Argynnis aglaja*.

Brown hares *Lepus europaeus* are abundant in the meadows; common blue butterflies *Polyommatus icarus* are recorded throughout the warmer summer months. Curlew *Numenius arquata*^{*}, linnet *Carduelis cannabina*[#], goldfinch *Carduelis carduelis* and meadow pipit *Anthus pratensis* breed on the site. House sparrow *Passer domesticus*^{*#}, house martin *Delichon urbica*, starling *Sturnus vulgaris*^{*#} and swallow *Hirundo rustica* also breed in the vicinity and utilise these fields. Visiting (non-breeding) birds include lesser black backed gull *Larus fuscus*, short-eared owl *Asio flammeus*^{*}, wheatear *Oenanthe oenanthe* and snipe *Gallinago gallinago*.

* = species on Schedule 1 of the Wildlife Act 1990

*** = species on Schedule 7 of the Wildlife Act 1990

= Red List Bird of Conservation Concern (UK, IoM and Channel Islands)

List of operations requiring DEFA's consent

The operations listed below may damage the features of interest of **Cronk e King ASSI**. Before any of these operations are undertaken you must consult DEFA, and may require our consent.

It is usually possible to carry out some of these operations in certain ways, or at specific times of year, or on certain parts of the ASSI, without damaging the features of interest. If you wish to carry out any of these activities please contact a DEFA Biodiversity Officer who will give you advice and, where appropriate, issue consent. Please help us by using the form sent to you with the notification documents to ask us for consent to carry out these operations.

In certain circumstances it will not be possible to consent these operations, because they would damage the features of interest. Where possible a Biodiversity Officer will suggest alternative ways in which you may proceed, which would enable consent to be issued. To proceed without DEFA's consent may constitute an offence. If consent is refused, or conditions attached to it, which are not acceptable to you, you will be provided with details of how you may appeal to DEFA.

Standard reference number	Type of operation
1	Cultivation, including ploughing, rotovating, harrowing and re-seeding.
2	Grazing, the introduction of grazing and alterations to the grazing regime (including type of stock, intensity or seasonal pattern of grazing).
3	Stock feeding, the introduction of stock feeding and alterations to stock feeding practice.
4	Mowing or cutting of vegetation (where already damaging), the introduction of mowing and alterations to the mowing or cutting regime (such as from haymaking to silage).
5	Application of manure, slurry, silage liquor, fertilisers and lime.
6	Application of pesticides, including herbicides (weed killers) whether terrestrial or aquatic, and veterinary products.
7	Dumping, spreading or discharge of any materials.
8	Burning and alterations to the pattern or frequency of burning.
9	Release into the site of any wild, feral, captive-bred or domestic animal, plant, seed or micro-organism (including genetically modified organisms).
10	Killing, injuring, taking or removal of any wild animal (including dead animals or parts thereof), or their eggs and nests, including pest control and disturbing them in their places of shelter.
11	Destruction, displacement, removal or cutting of any plant or plant remains, including tree, shrub, herb, hedge, dead or decaying wood, moss, lichen, fungus, leaf-mould, and turf.
12	Tree and/or woodland management (where already damaging), the introduction of tree and/or woodland management (where applicable) and alterations to tree and/or woodland management (including planting, felling, pruning and tree surgery, thinning coppicing, changes in species composition, removal of fallen timber).
13 a)	Draining (including moor- gripping, the use of mole, tile, tunnel or other artificial drains).
13 b)	Modification to the structure of water courses (e.g. rivers, streams, springs, ditches and drains), including their banks and beds, as by re-alignment,

	regrading, damming, or dredging.
13 c)	Management of aquatic and bank vegetation for drainage purposes.
14	Alterations to water levels and tables and water utilisation (including irrigation, storage, abstraction from existing water bodies and through boreholes). Also the modification of current drainage practices.
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, hard rock, sand, gravel, topsoil, subsoil, lime, limestone pavement, shells and spoil.
21	Destruction, construction, removal, rerouting, or regrading of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, including soil and soft rock exposures or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures or the undertaking of engineering works, including drilling.
24 a)	Modification of natural or man-made features (including cave entrances) and clearance of boulders, large stones, loose rock or scree.
26	Use of vehicles or craft.
27	Recreational or other activities.
28 a)	Game and waterfowl management and hunting practices (where already damaging), introduction of game or waterfowl management (where applicable) and alterations to game and waterfowl management and hunting practice.

Notes:

1. This is a list of operations appearing to DEFA to be likely to damage the special features of this ASSI, as required under Section 27 of the Wildlife Act 1990.
2. Any reference to 'animal' in this list shall be taken to include any mammal, reptile, amphibian, bird, fish or invertebrate.

Date notified: 21st October 2013

Date confirmed: 19th February 2014

Date of last revision: N/A

Views about Management

This is not a legal document

A statement of DEFA's views about the management of Cronk e King Area of Special Scientific Interest (ASSI)

This statement represents DEFA's views about the management of the ASSI for nature conservation. This statement sets out in principle, our views on how the area's special conservation interest can be conserved and enhanced. DEFA does not have a duty to notify the owners and occupiers of ASSI of its views about the management of the land, but DEFA has determined that this would be good practice with ASSI designations.

Not all of the management principles will be equally appropriate to all parts of the ASSI. Also, there may be other activities, additional to our current views, which can be beneficial to the conservation and enhancement of the features of interest.

This statement does not constitute consent for any of the 'operations requiring DEFA's consent'. The written consent of DEFA is required before carrying out any of those operations. DEFA welcomes consultation with owners, occupiers and users of the ASSI to ensure that the management of this site conserves and enhances the features of interest, and to ensure that all necessary prior consents are obtained.

Management Principles

Continuation of the hay meadow management will protect the existing natural habitats of interest on this site. This involves cutting after 31st July each year in a wildlife-friendly manner and removing the hay. Winter sheep grazing (November to April) should not cause a problem for the designated interest. Such management, avoiding agricultural improvement, will maintain the diverse grassland flora and suitable soil conditions for the orchids. Farmyard manure could be applied with prior consent in order to maintain productivity, but artificial fertilisers may be damaging and should be avoided. The application of pesticides should be avoided generally but the Department will discuss issues arising and the most appropriate way of tackling them within the designated area.

Date notified: 21st October 2013

Date confirmed: 19th February 2014

Date of last revision: N/A

Cronk y Bing Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin



**Isle of Man
Government**
Reilthys Eilian Vannin

Department of Agriculture,
Fisheries and Forestry


**CRONK Y BING
AREA OF SPECIAL
SCIENTIFIC INTEREST**

Area: 17.71ha (= 43.76 acres) *
Designated: 14th March 2006
Confirmed: 9th August 2006

*Note: the seaward boundary of the site
is defined as the mean low water mark
between points A and B on this map.
The low water mark, and hence the
overall site area, may be subject to
variation with time.

The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Cronk y Bing

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Andreas

Planning Authority: Department of Local Government and the Environment

Local authority: Andreas Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets NX 30

National Grid Reference (centroid): NX381017

Area: 17.71ha (43.76 acres)

Date notified: 14th March 2006

Date confirmed: 9th August 2006

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its wildlife habitat – soft cliffs, dunes and coastal grassland.

Description and reasons for notification:

Cronk y Bing consists of a wide strip of sand dunes, with soft cliffs on the seaward side and a flat plateau of semi-fixed dune grassland on the landward side, where sand has been removed from the dunes in the past. The seaward dunes are “yellow dunes”, with a relatively mobile structure, dominated by marram grass *Ammophila arenaria*. The loose, sandy structure of the yellow dunes favours plants such as sea spurge *Euphorbia paralias*, sea bindweed *Calystegia soldanella*, sea holly *Eryngium maritimum*, and common cornsalad *Valerianella locusta*. These dunes rise to around 4m on the seaward edge, ending abruptly in soft cliffs in which the burrows of mining bees and other invertebrates can be seen in summer.

The plateau of grassland behind the high ridge of dunes has marram areas with short, coastal turf and almost-bare ground, some of which is dominated by lichens, notably “spaghetti lichen”, *Cladonia portentosa*. True dog-lichen, *Peltigera canina*, is also frequent in places, along with low-growing burnet rose *Rosa pimpinellifolia*. Coastal grassland plants in the more developed turf include sea storksbill *Erodium maritimum*, common storksbill *Erodium cicutaria* agg., rest harrow *Ononis repens*, wild thyme *Thymus polytrichus*, wild carrot *Daucus carota*, harebell *Campanula rotundifolia*, sheep’s-bit *Jasione montana*, and hundreds of spikes of pyramidal orchid *Anacamptis pyramidalis***. Several of the above species – notably pyramidal orchid and true dog-lichen – rely on calcareous grassland habitats, and are consequently of limited distribution on the Island. The grassland plateau is relatively sheltered by the dunes, and attracts invertebrates and songbirds such as nesting meadow pipit.

The seaward side of the dunes consists of sandy shingle, strandline and a wide intertidal zone from which material blows in on the prevailing wind to form and maintain the dunes. The beach zone attracts occasional ground-nesting birds including oystercatcher *Haematopus ostralegus* and ringed plover *Charadrius hiaticula**. It also offers suitable habitat for a range of other feeding and breeding birds including little tern *Sterna albifrons** and curlew *Numenius arquata*.*

Cronk y Bing has been managed as a nature reserve, and remains relatively natural and representative of dunes and cliffs along this stretch of coast, unlike adjacent dune areas which have received vehicle damage, extensive quarrying, tipping and other disturbance. It therefore represents a good example of dune habitat, complementing the Ayres NNR habitats to the north.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

As an additional point of botanical interest, the range of coastal habitats and vegetation communities present at Cronk y Bing offers an important refuge for those plant species which are ephemeral in nature, and which do not appear at the same site year after year. Such species require a certain amount of suitable habitat in order to maintain their presence as part of the native flora of the Island, and will become extinct if no habitats are available for them to colonise on an occasional basis. Examples which have been recorded here in the past include several rare and protected species, such as oysterplant *Mertensia maritima***, Isle of Man cabbage *Coinceya monensis subsp. monensis***, smooth cat's-ear *Hypochoeris glabra*** and viper's bugloss *Echium vulgare***.

The proximity to the Lhen Trench benefits birds and semi-aquatic invertebrates on the site, linking it with adjacent feeding wader habitats and other semi-natural habitats inland.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Cronk y Bing

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17	Reclamation of land from sea, estuary or marsh.
18	Bait digging on inter-tidal areas, (if it is shown to be damaging).
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of

	boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest (excluding walking, with or without dogs).

† Note: each type of operation has a standard reference number. For each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* “animal” includes any mammal, reptile, bird, fish or invertebrate.

Dalby Coast Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmhtaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Keillrys Ellan Vannin


Dalby Coast Area of Special Scientific Interest

Designated: 7th October 2009

Varied and confirmed: 1st March 2010

Area: 62.1 hectares (153.4 acres)

National Grid Reference (centroid): SC 212782



CONFIRMATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Dalby Coast

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Patrick

Local authority: Patrick Commissioners

Planning Authority: Department of Local Government and the Environment

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC27, SC28

National Grid Reference (centroid): SC 212782

Area: 62.1 hectares (153.4 acres)

Date notified: 7th October 2009

Date confirmed: 1st March 2010

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its geology, flora and wildlife habitats – species-rich meadow, hard coastal cliffs and slopes, coastal grassland, neutral flush, reedbed, vegetated shingle and intertidal habitats.

Description and reasons for notification:

The Dalby Coast ASSI boundary encompasses an important matrix of coastal habitats, grading from species-rich hay meadow to rocky shore. Of most botanical interest is the thin strip of land that forms at the base of the steep coastal slopes that forms the interface between the steep, sloping maritime grassland and the rocky shore. These areas are characterised by base-rich flushes, with orchids and sedges occurring in abundance. Within this narrow zone, a maritime influence quickly dominates and botanical interest is increased with elements of saltmarsh vegetation dominating the upper shore.

The steep rocky cliffs and coastal broughs that stretch from Glen Maye to an area south of Niarbyl are one of the largest continuous areas of unimproved coastal habitat in the Isle of Man. The coast is generally rocky with gently sloping grassy slopes becoming steep south of Niarbyl with low cliffs in places. A number of small streams and springs flow down to the sea and create botanically-interesting wet flushes. One of the largest stands of reedbed occurs to the south of the site.

The coast at Niarbyl is also an important geological site. Recently recognised sandstones of Silurian age (Dalby Group, Niarbyl Formation) meet Ordovician mudstones and siltstones of the Manx Group (Creggan Mooar Formation) on the beach to the north of the cottages. A major fault line across the beach marks the contact between these rocks from two ancient continents that were once on opposite sides of the world, and the closure of the Iapetus Ocean that once separated them.

This stretch of coast, particularly the area to the north of Niarbyl is little disturbed by man. In 'The Naturalist in Man', Garrad wrote '*Where the farmland runs down to the cliff top a substantial hedge marks the limit of cultivation and prevents the descent of grazing animals. In consequence a narrow band of vegetation remains almost unaffected by farming activities....A more varied wildlife survived (here) because of this unintentional protection and the favourable microclimate produced by the nearness of the sea*'.

In addition to providing valuable nesting habitat for a range of coastal birds such as fulmar *Fulmarus glacialis* raven *Corvus corax* and chough *Pyrhcorax pyrrhcorax**, the cliffs support patches of diverse coastal grassland, cliff crevice plants and scrub.

Detached cliff sections which now lie within the intertidal zone have similar characteristics to the main cliffs and slopes. Plants present include abundant thrift *Armeria maritima*, English stonecrop *Sedum anglicum*, sea plantain *Plantago maritima*, sea spleenwort *Asplenium marinum*, sea campion *Silene uniflora*, wild thyme *Thymus polytrichus*, sheep's-bit scabious *Jasione montana*, common bird's-foot trefoil *Lotus corniculatus*, red fescue *Festuca rubra*, silver hair-grass *Aira caryophyllaea*, honeysuckle *Lonicera periclymenum*, cat's ear *Hypochoeris radicata*, bramble *Rubus fruticosus* agg., dandelion *Taraxacum* sp., and a good range of plants typical of short, neutral grassland. Small freshwater flushes on the sides of the coastal slopes vary the flora, with species such as brookweed *Samolus valerandi*, yellow-sedge *Carex viridula* and lesser spearwort *Ranunculus flammula*.

The whole site is notable for its diversity of habitats. In particular the wet flushes at the base of the cliffs have a high species diversity. Orchid species include Heath-spotted orchid *Dactylorhiza maculata***, Common spotted orchid *Dactylorhiza fuchsia***, Northern marsh orchid *Dactylorhiza purpurella*** and there are historic records of O'Kelly's orchid *Dactylorhiza fuchsii subsp okellyi***.

Sedges are particularly well represented across all habitats and highlight the diversity of habitats and species-richness of the site with 15 species identified; Distant sedge *Carex distans*, tawny sedge *Carex hostiana*, yellow sedge *Carex viridula subsp. serotina*, long-bracted sedge *Carex extensa*, carnation sedge *Carex panicea*, common sedge *Carex nigra*, sand sedge *Carex arenaria*, false-fox sedge *Carex otrubae*, glaucous sedge *Carex flacca*, star sedge *Carex echinata*, oval sedge *Carex ovalis*, flea sedge *Carex pulicaris*, yellow sedge *Carex viridula subsp. oedocarpa*, yellow sedge *Carex viridula subsp. viridula* and hairy sedge *Carex hirta*.

Other interesting species recorded recently include black bog rush *Schoenus nigricans*, few-flowered spike rush *Eleocharis quinquefolia*** and saltmarsh flat-sedge *Blysmus rufus***.

The wide-range of semi-natural habitats grading from species-rich meadow to sea-cliff provide excellent breeding and feeding territory for birds, which are sufficient to merit site selection on the grounds of breeding bird diversity alone. Recorded breeding birds that contribute to the scoring of the site for species associated with hard coastline include Fulmar *Fulmarus glacialis*, Shelduck *Tadorna tadorna**. Eider *Somateria mollissima*, Oystercatcher *Haematopus ostralegus*, Ringed plover *Charadrius hiaticula**, Great-black backed gull *Larus marinus*, Rock pipit *Anthus petrosus*, Stonechat *Saxicola torquata*, Raven *Corvus corax* and Chough *Pyrhcorax pyrrhcorax**.

A total of 32 definite or probable breeding bird species has been recorded since 1998; there are also recent records of a further 44 species of non-confirmed breeding and/or passage or wintering birds, of which 16 are listed on Schedule 1 of the Wildlife Act 1990.

Viviparous lizards*** have been recorded on the grass banks on the beach road to Niarbyl. It is likely that they occur along the entire coastline of the site.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

***= Protected under Schedule 5 of the Wildlife Act 1990

Other information:

The designation boundary abuts that of Glen Maye ASSI.

CONFIRMATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Dalby Coast ASSI

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control.
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17	Reclamation of land from sea, estuary or marsh.
18	Bait digging on inter-tidal areas, (if it is shown to be damaging).
19	Erection of sea defences or coastal protection works, including cliff or land-slip

	drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest.
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora.
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, amphibian, bird, fish or invertebrate.

1st March 2010

Dhoon Glen

Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



Isle of Man
Government

Reiltys Ellan Vannin



**Isle of Man
Government**
Reillys Ellan Vannin

Department of Agriculture,
Fisheries and Forestry

**DHOON GLEN
AREA OF SPECIAL
SCIENTIFIC INTEREST**

ASSI boundary



Area: 20.92ha (= 51.69 acres) *
OS Grid reference: SC458865
(approximate centre of site)

Designated: 27th September 2006
Boundary varied: 21st March 2007

*Note: the seaward boundary of the site is
defined as the Lowest Astronomical Tide
between points A and B on this map.


The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

Base Map is

Crown Copyright

Department of Local Government
and the Environment, Isle of Man.

Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Dhoon Glen

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Maughold

Planning Authority: Department of Local Government and the Environment

Local authority: Maughold Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC48

National Grid Reference (centroid): SC458865

Area: 20.92ha (51.69 acres)

Date notified: 27th September 2006

Date varied: 21st March 2007

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its flora and wildlife habitats – broadleaved woodland, mixed plantation, open water, strandline and shingle.

Description and reasons for notification:

Dhoon Glen consists of a series of small, steep tributary valleys and one exceptionally deep glen, notable for a tall and powerful waterfall. The sides of the tributaries and glen have a very diverse assortment of woodland types, ranging from extensive areas of semi-natural broadleaved woodland through to mixed, broadleaved and conifer plantation. The most diverse areas have a canopy of ash *Fraxinus excelsior*, oak *Quercus petraea*, birch *Betula pendula* and rowan *Sorbus aucuparia*, although sycamore *Acer pseudoplatanus* and beech *Fagus sylvatica*-dominated areas also exhibit a diverse semi-natural groundflora. Herb species present include abundant bluebell *Hyacinthoides non-scripta*, with wood anemone *Anemone nemorosa*, yellow pimpernel *Lysimachia nemorum*, wood horsetail *Equisetum sylvaticum*, sanicle *Sanicula europaea*, wood sorrel *Oxalis acetosella*, opposite-leaved golden saxifrage *Chrysosplenium oppositifolium*, foxglove *Digitalis purpurea*, greater stitchwort *Stellaria holostea*, honeysuckle *Lonicera periclymenum* and primrose *Primula vulgaris*. The Glen is particularly rich in ferns, which include hay-scented buckler-fern *Dryopteris aemula***, hard fern *Blechnum spicant*, soft shield-fern *Polystichum setiferum*, lady-fern *Athyrium filix-femina*, male fern *Dryopteris filix-mas*, scaly male fern *Dryopteris affinis*, broad buckler-fern *Dryopteris dilatata*, hart's-tongue *Phyllitis scolopendrium*, black spleenwort *Asplenium adiantum-nigrum* and, at the coastal end of the Glen, sea spleenwort *Adiantum marinum*. Killarney filmy fern *Trichomanes speciosum*** (gametophyte stage only) has been recorded from the base of the waterfall. This species is internationally rare and threatened, and is listed under Annex 2 of the EU Habitats and Species Directive. It is also protected under the Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention).

The site boundary was chosen to include the full range of the Glen from fast-flowing tributaries to the sea, where the fresh water meets the tide in a series of small, inter-tidal rockpools noted for their diversity and consequent educational value. Other coastal features relating to the Glen include the presence of diverse, regenerating coastal woodland and scrub, with frequent coastal cliff and grassland plants also present. These notably include scattered clumps of very dwarf oak trees, kept to a height of a few feet only due to the very exposed conditions. These oaks are relatively safe from grazing livestock, and appear to represent a relict native oak population. They are consequently of research importance as well as providing a valuable habitat feature in their own right.

The large area of diverse, well-developed semi-natural habitats throughout the Glen provides excellent breeding and feeding territory for birds, which are almost diverse enough on the site to merit selection on the grounds of woodland or coastal breeding bird assemblage alone. Glenside breeding species include songthrush *Turdus philomelos** (an RSPB Red List species of concern), kestrel *Falco tinnunculus**, swallow *Hirundo rustica*, grey wagtail *Motacilla cinerea*, wood warbler *Phylloscopus sybillatrix*, willow warbler *Phylloscopus trochilus* and goldcrest *Regulus regulus*. Breeding birds along the coastal edge of the Glen include chough *Pyrrhocorax pyrrhocorax**, stonechat *Saxicola torquata*, meadow pipit *Anthus pratensis*, oystercatcher *Haematopus ostralegus*, Fulmar *Fulmarus glacialis* and herring gull *Larus argentatus* (all the latter are on the RSPB Amber List).

The diversity of habit types that supports such a wide range of bird species also benefits invertebrates, in particular moths. Macro-moth species of note include the UK Nationally Notable Black-banded *Polymixis xanthomista*, and the Grey *Hadena caesia* ssp *mananii****. The Isle of Man is the "type location" for these species; hence their Manx populations are of importance to the taxonomic study of macro-moths. In total, 80% of all the recently-recorded macro-moth species on the Island have been recorded from Dhoon Glen and the adjacent coast, making this an exceptionally diverse site.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

*** = Protected under Schedule 5 of the Wildlife Act 1990

Other information:

Dhoon Glen is centred on the Dhoon Granite, an igneous intrusion into the surrounding Manx slate. Igneous intrusions are an unusual feature on the Island and hence of geological as well as biological interest. The Glen also contains significant archaeological features, in particular the relics of mining and quarrying activity. The presence of spectacular waterfalls and dramatic geology and flora have led to a long history of use as a "pleasure glen", and recreational access is still of importance in the publicly-accessible parts of the site.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Dhoon Glen

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseeding
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	The introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning and lighting of fires.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.

23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles# or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora (excluding walking, with or without dogs).
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, amphibian, reptile, bird, fish or invertebrate.

"vehicles" includes bicycles.

Eary Vane Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmhtaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**


Reiltys Ellan Vannin



**Isle of Man
Government**
Reilbys Ellan Vannin


Department of Agriculture,
Fisheries and Forestry

**EARY VANE
AREA OF SPECIAL
SCIENTIFIC INTEREST**



Area: 3.96ha (= 9.78 acres)
O/S Grid reference: SC338818
(approximate centre of site)
Designated: 20th December 2006
Confirmed: 2nd May 2007

The grid lines on this map form
part of the National Grid,
and are spaced at 100m intervals
Base Map is
Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Eary Vane

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Marown

Planning Authority: Department of Local Government and the Environment

Local authority: Marown Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC38

National Grid Reference (centroid): SC338818

Area: 3.96ha (9.78 acres)

Date notified: 20th December 2006

Date confirmed: 2nd May 2007

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its habitat: species-rich rush pasture of the Sharp-flowered rush and Marsh bedstraw *Juncus acutiflorus* – *Galium palustre* type, with Purple moor-grass and Tormentil *Molinia caerulea* – *Potentilla erecta* mire.

Description and reasons for notification:

Eary Vane consists of a series of small fields on the lower south-eastern edge of Slieu Ruy, with the smaller hill of Cronk Breck to the south. The hillside location has resulted in a localised mire which ultimately drains eastwards into a tributary of the River Glass. The fields have not been subjected to intensive agricultural improvement and now support fen meadow habitat, dominated by rush pasture of the Sharp-flowered rush and Marsh bedstraw *Juncus acutiflorus* – *Galium palustre* type. This is accompanied by a range of associated habitats including Purple moor-grass *Molinia caerulea*-dominated mire, ditches and streams with aquatic vegetation, gorse and willow scrub, and dry acid grassland. The area has good diversity, with over half of all the higher plant species typical of the best Manx examples of this habitat recorded in recent years.

The structure of the vegetation is varied, ranging from tall tussocky rush pasture to short turf on the sod banks and drier parts of the area, and short boggy vegetation alongside streams and drains and in dips and hollows. Large numbers of Heath spotted-orchid *Dactylorhiza maculata*** are present throughout most of the site. Other species of the marshy grassland areas include frequent Sharp-flowered rush *Juncus acutiflorus*, Jointed rush *Juncus articulatus*, Heath rush *Juncus squarrosus*, Purple moor-grass *Molinia caerulea*, Velvet bent *Agrostis canina*, Star sedge *Carex echinata*, Dwarf yellow-sedge *Carex viridula* subspecies *viridula*, Carnation sedge *Carex panicea*, Marsh violet *Viola palustris*, Common lousewort *Pedicularis palustris*, Devil's-bit scabious *Succisa pratensis*, Cuckoo-flower *Cardamine pratensis* and Marsh pennywort *Hydrocotyle vulgaris*. Plants more typical of dry, acid grassland occur on the edges of sod banks and around the drier parts of the site, including Red fescue *Festuca rubra*, Mat-grass

Nardus stricta, Wavy hair-grass *Deschampsia flexuosa*, Sweet vernal grass *Anthoxanthum odoratum*, Pill sedge *Carex pilulifera*, Green-ribbed sedge *Carex binervis*, Harebell *Campanula rotundifolia*, Heath bedstraw *Galium saxatile*, Tormentil *Potentilla erecta*, Heath milkwort *Polygala serpyllifolia*, Bilberry *Vaccinium myrtillus* and Ling *Calluna vulgaris*.

The grassland habitat is varied by the presence of small boggy areas dominated by Sphagnum moss, with bog plants such as Common cotton-grass *Eriophorum angustifolium*, Round-leaved sundew *Drosera rotundifolia*, Bog asphodel *Narthecium ossifragum* and Bog stitchwort *Stellaria uliginosa*. Other wetland plants grow in and alongside the stream and ditches which cross the site, most of which have both submerged and emergent vegetation. In addition to bog plants this includes aquatic and emergent species such as Ivy-leaved crowfoot *Ranunculus hederaceus*, Lesser spearwort *Ranunculus flammula*, Tufted forget-me-not *Myosotis laxa*, Creeping forget-me-not *Myosotis secunda*, Marsh willowherb *Epilobium palustre* and the moss *Philonotis fontana*.

A total of 78 plant species has been recorded at Eary Vane within the last year, making this a very good example of the habitat type. The presence of a wide variety of both sward structure and species also makes this area suitable for birds and invertebrates which are typical of Manx upland margin habitats, including breeding Wren *Troglodytes troglodytes*, Meadow pipit *Anthus pratensis*, and Curlew *Numenius arquata**.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

This site is named after the old farmstead of Eary Vane, situated to the north of the ASSI fields.

Species-rich rush pastures are internationally threatened, having suffered very heavy losses due to agricultural improvement, drainage and development. This applies particularly to species-rich, lowland examples. Countries within the British Isles are considered to have a special responsibility for the protection of this habitat, which is in danger of extinction if the current European rate of loss continues. In view of this, "Purple moor-grass and Rush Pastures" are now a Priority Habitat in the UK Biodiversity Action Plan, and in recent years several of the most diverse UK examples have been designated SSSIs and/or Special Areas of Conservation.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST **Appendix II**

Site Name: Eary Vane

Operations likely to damage the special interest of the site

<u>Standard reference number</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.

23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
26	Use of vehicles or craft likely to damage or disturb features of interest.
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora (excluding walking, with or without dogs).
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, bird, fish or invertebrate.

Glen Maye Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture


Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin



**Isle of Man
Government**
Reihys Eilan Vannin

Department of Agriculture,
Fisheries and Forestry

**GLEN MAYE
AREA OF SPECIAL
SCIENTIFIC INTEREST**

 Designation boundary

Area: 15.92ha (39.34 acres)
O/S Grid reference: SC229798
(approximate centre of site)

Designated: 31/10/2007

Confirmed: 22/04/2008

*Note: the seaward boundary of the site is
defined as the Lowest Astronomical Tide
between points A and B on this map.

The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

Base Map is
Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Glen Maye

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Patrick

Local authority: Patrick Commissioners

Planning Authority: Department of Local Government and the Environment

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC27, SC28

National Grid Reference (centroid): SC229798

Area: 15.92ha (39.34 acres)

Date notified: 31st October 2007

Date confirmed: 22nd April 2008

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its flora and wildlife habitats – broadleaved woodland, mixed plantation, coastal scrub, open water, hard coastal cliffs and slopes, coastal grassland and intertidal habitats.

Description and reasons for notification:

Glen Maye is one of the largest National Glens, consisting of a steep, wide gorge with broadleaved and mixed plantation on the sides, leading up to very tall, south-facing cliffs and west-facing coastal slopes. An exceptional range of associated habitats is present on the site, with good natural gradation between woodland and scrub, scrub and grassland, coastal slopes and the shore. The type and aspect of the open water on the site is also varied, as several small streams and flushes, and one large tributary, feed into the Glen from both the north and south sides. The largest tributary forms a substantial glen in its own right, also with semi-natural broadleaved vegetation and scrub on either side of the watercourse. The Glen Maye ASSI boundary has been chosen to include the full range of these habitats.

The Glen woodland is varied both in the canopy and the groundflora. The canopy reflects Victorian tree-planting, but also contains semi-natural regrowth and native scrub species. Trees and shrubs include ash *Fraxinus excelsior*, oak *Quercus petraea*, sycamore *Acer pseudoplatanus*, wych elm *Ulmus glabra*, alder *Alnus glutinosa*, beech *Fagus sylvatica*, blackthorn *Prunus spinosa*, hazel *Corylus avellana*, grey willow *Salix cinerea*, eared willow *Salix aurita*, rowan *Sorbus aucuparia*, holly *Ilex aquifolium*, elder *Sambucus nigra*, Wild and naturalised roses *Rosa spp.*, and small patches of sweet chestnut *Castanea sativa* and conifers including Scots pine *Pinus sylvestris*. Introduced shrubs such as snowberry, *Symphoricarpos albus*, rhododendron, *Rhododendron ponticum agg.* and fuchsia, *Fuchsia magellanica*, occur but in small amounts.

The majority of the woodland, and also the scrub on the coastal slopes, has a groundflora which is semi-natural in character and supports a good range of typical native woodland herbs, including bluebell *Hyacinthoides non-scriptus*, wild garlic *Allium ursinum*, wood sorrel *Oxalis acetosella*, red campion *Silene dioica*, primrose *Primula vulgaris*, pignut *Conopodium majus*, greater stitchwort *Stellaria holostea*, yellow pimpernel *Lysimachia nemorum* and hairy woodrush *Luzula pilosa*. There are small populations of the rare, protected species, wood vetch *Vicia sylvatica*** and hairy brome *Bromopsis ramosa*. Wet areas of the glen banks support shade-tolerant marsh plants such as marsh marigold *Caltha palustris*, wild angelica *Angelica sylvestris*, lesser spearwort *Ranunculus flammula* and opposite-leaved golden saxifrage *Chrysosplenium oppositifolium*.

At the coastal end of the site, the woodland grades into coastal slopes and sheer cliffs. In addition to providing valuable nesting habitat for a range of coastal birds such as fulmar *Fulmarus glacialis*, the cliffs support patches of diverse coastal grassland, cliff crevice plants and scrub.

Detached cliff sections which now lie within the intertidal zone have similar characteristics to the main cliffs and slopes. Plants present include abundant thrift *Armeria maritima*, English stonecrop *Sedum anglicum*, sea plantain *Plantago maritima*, sea spleenwort *Asplenium marinum*, sea campion *Silene uniflora*, wild thyme *Thymus polytrichus*, sheep's-bit scabious *Jasione montana*, common bird's-foot trefoil

Lotus corniculatus, red fescue *Festuca rubra*, silver hair-grass *Aira caryophyllaea*, honeysuckle *Lonicera periclymenum*, cat's ear *Hypochoeris radicata*, bramble *Rubus fruticosus* agg., dandelion *Taraxacum* sp., and a good range of plants typical of short, neutral grassland. Small freshwater flushes on the sides of the coastal slopes vary the flora, with species such as brookweed *Samolus valerandi*, yellow-sedge *Carex viridula* and lesser spearwort *Ranunculus flammula*.

The whole site is notable for its abundance and diversity of ferns. At the coastal end there is frequent sea spleenwort *Asplenium marinum*, maidenhair spleenwort *Asplenium trichomanes*, black spleenwort *Asplenium adiantum-nigrum* and bracken *Pteridium aquilinum*. The mild and sheltered climate of the river banks and woodland supports abundant soft shield fern *Polytrichum setiferum*, lady fern *Athyrium filix-femina*, male fern *Dryopteris filix-mas*, scaly male fern *Dryopteris affinis*, broad buckler-fern *Dryopteris dilatata*, hart's-tongue *Phyllitis scolopendrium*, western polypody *Polypodium interjectum* and hard fern *Blechnum spicant*. The rarities narrow buckler-fern *Dryopteris carthusiana** and southern polypody *Polypodium cambricum* were last recorded from this area.

The Glen Maye river and its main tributary are too shaded and fast-flowing to support a wide range of submerged and emergent plants. Where these factors permit, small populations of tenacious species such as willow moss *Fontinalis antipyretica* and the freshwater alga *Lemanea* sp. are present. The river is rocky, with one substantial waterfall and several smaller torrents, occasional small pools and numerous small caves on the rocky banks. The conditions are ideal for breeding grey wagtail *Motacilla cinerea* which breed here. The Glen forms part of a migratory route for sea trout.

The well-developed semi-natural habitats throughout the Glen provide excellent breeding and feeding territory for birds, which are almost diverse enough to merit site selection on the grounds of breeding bird diversity alone. Recorded glenside breeding birds include song thrush *Turdus philomelos**#, mistle thrush *Turdus viscivorus*, blackbird *Turdus merula*, blackcap *Sylvia atricapilla*, chaffinch *Fringilla coelebs*, goldcrest *Regulus regulus*, kestrel *Falco tinnunculus**, willow warbler *Phylloscopus trochilus* and dunnoek *Prunella modularis*. Breeding birds on the cliffs, coastal grassland and scrub at the mouth of the Glen include fulmar, linnet *Carduelis cannabina*#, rock pipit *Anthus spinoletta*, stonechat *Saxicola torquata*, meadow pipit *Anthus pratensis*, whitethroat *Sylvia communis*, herring gull *Larus argentatus* and great black-backed gull *Larus marinus*. Eider *Somateria mollissima* have bred here and shelduck *Tadorna tadorna** have been recorded from the intertidal waters adjacent to the Glen mouth. Other birds recorded on the site include twite *Carduelis flavirostris**#, ringed plover *Charadrius hiaticula**, peregrine *Falco peregrinus** and grey heron *Ardea cinerea*. A total of 29 definite or probable breeding bird species has been recorded since 1998; there are also recent records of a further 36 species of non-confirmed breeding and/or passage or wintering birds, of which 13 are listed on Schedule 1 of the Wildlife Act 1990 and one of these is confirmed breeding. For birds this represents one of the most diverse National Glens.

* = Protected under Schedule 1 of the Wildlife Act 1990

= Birds of Conservation Concern Red List species ¹

** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

This is a very well-documented and popular Glen, long recognised as an important botanical site. Recreational access is of importance and there is a long history of use as a "pleasure glen" both for walking access and fishing. The Glen Maye River corridor reaches back inland as far as Glen Rushen ASSI; Glen Maye represents the culmination of this important habitat corridor.

¹ Gregory Richard D, Wilkinson Nicholas I, Noble David G, Robinson James A, Brown, Andrew F, Hughes, Julian, Proctor Deborah, Gibbons David W, and Galbraith Colin A. (2002) The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002-2007. **British Birds** Vol 95 No 9 Pages 410-448

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Glen Maye

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	The introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	The introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires (where already damaging) and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herbs, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
18	Bait digging on inter-tidal areas, (where already damaging).
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works,

	including drilling,.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest (excluding walking, with or without dogs).
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, bird, fish or invertebrate.

Glen Rushen Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmmltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin



**Isle of Man
Government**
Raijys Eilan Vannin

Department of Agriculture,
Fisheries and Forestry

**GLEN RUSHEN
AREA OF SPECIAL
SCIENTIFIC INTEREST**


□ ASSI boundary

Area: 12.27ha (= 30.32 acres)
O/S Grid reference: SC23667652
(approximate centre of site)

Designated: 20th December 2006
Confirmed: 2nd May 2007

The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

Base Map is
Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Glen Rushen

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Patrick

Planning Authority: Department of Local Government and the Environment

Local authority: Patrick Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC27

National Grid Reference (centroid): SC23667652

Area: 12.27ha (30.32 acres)

Date notified: 20th December 2006

Date confirmed: 2nd May 2007

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its habitat: wet unimproved acid grassland, valley mire and flush, developing native broadleaved woodland, river valley and streams.

Description and reasons for notification:

Glen Rushen forms part of an important river habitat corridor leading from Dalby Mountain moorland through Glen Rushen and Glen Moar to eventually reach the sea at Glen Maye. The Glen Rushen stretch is notable for its naturalness and diversity, which includes sloping, damp acid grassland with patches of mire vegetation and small, boggy flushes on the valley sides. These support abundant orchids *Dactylorhiza maculata subspecies ericetorum**** and a thriving colony of pale butterwort *Pinguicula lusitanica****, a carnivorous plant which requires undisturbed, nutrient-poor acidic habitats and which has a very localised distribution on the Island. A wide range of wet, acid grassland species is present including Purple moor-grass *Molinia caerulea*, sharp-flowered rush *Juncus acutiflorus*, heath rush *Juncus squarrosus*, heath woodrush *Luzula multiflora*, yellow sedge *Carex viridula subspecies oedocarpa*, common sedge *Carex nigra*, star sedge *Carex echinata*, common lousewort *Pedicularis sylvatica*, marsh lousewort *Pedicularis palustris*, marsh arrowgrass *Triglochin palustris*, devil's-bit scabious *Succisa pratensis*, marsh pennywort *Hydrocotyle vulgaris*, marsh thistle *Cirsium palustre* and bog stitchwort *Stellaria uliginosa*. Drier grassland areas such as the sides of old sod banks support plants such as sheep's fescue *Festuca ovina*, sweet vernal-grass *Anthoxanthum odoratum*, red fescue *Festuca rubra*, mat-grass *Nardus stricta*, pill sedge *Carex pilulifera*, spring sedge *Carex caryophyllaea*, cat's-ear *Hypochaeris radicata*, heath milkwort *Polygala serpyllifolia*, hard fern *Blechnum spicant*, eyebright *Euphrasia agg.*, common bird's-foot trefoil *Lotus corniculatus*, bilberry *Vaccinium myrtillus* and abundant tormentil *Potentilla erecta*.

The grassland habitat throughout the Glen Rushen area is characterised by small flushes, boggy areas and mire alongside the watercourses, which support a range of bog and marsh plants including common cotton-grass *Eriophorum angustifolium*, *Sphagnum* mosses, flea sedge *Carex pulicaria*, star sedge *Carex echinata*, round-leaved sundew *Drosera rotundifolium*, pale butterwort *Pinguicula lusitanica*, heath spotted-orchid *Dactylorhiza maculata subspecies*

*ericetorum****, bog asphodel *Narthecium ossifragum*, bog pondweed *Potamogeton polygonifolius*, bog pimpernel *Anagallis tenella*, lesser spearwort *Ranunculus flammula* and marsh violet *Viola palustris*.

The river itself is notable for its natural course and the presence of marshy, valley mire vegetation alongside, locally dominated by whorled caraway *Carum verticillatum*, a distinctive western British species with a limited distribution on the Island. Scattered broadleaved trees and willow scrub alongside the river and the larger tributary include ash *Fraxinus excelsior*, wych elm *Ulmus glabra*, common alder *Alnus glutinosa* and rowan *Sorbus aucuparia*. Some of these have been planted, and some have grown up naturally. The groundflora in both cases is diverse, with some woodland species including bluebell *Hyacinthoides non-scriptus*, primrose *Primula vulgaris*, pignut *Conopodium majus*, and red campion *Silene dioica*.

The Glen Rushen valley provides a diverse and structurally varied habitat for birds, invertebrates, small mammals and frogs *Rana temporaria***. Hen harrier *Circus cyanea**, chough *Pyrrhocorax pyrrhocorax** and peregrine falcon *Falco peregrinus** breed near to Glen Rushen, and its fields and river banks form a key feeding area. The marshy fields also support breeding curlew *Numenius arquata**.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 5 of the Wildlife Act 1990

*** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

This area links several important areas of semi-natural habitats, notably wet heath on Dalby Mountain, dry dwarf shrub heath on either side of the valley, and the Glen Moar/Glen Maye River.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST **Appendix II**

Site Name: Glen Rushen

Operations likely to damage the special interest of the site

<u>Standard reference number</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.

23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora.
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, bird, fish or invertebrate.

Greeba Mountain & Central Hills Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:


- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture
Rheynn Chymmltaght, Bee as Eiriny's

This Sleaf Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



Isle of Man
Government
Reillys Ellen Vannia



**Isle of Man
Government**



Ennagh, Ballyvaughan

Department of Agriculture,
Fisheries and Forestry

**GREEBA MOUNTAIN
AND CENTRAL HILLS
AREA OF SPECIAL
SCIENTIFIC INTEREST**

Area: 1080.95ha (2671 acres)

Greeba Mountain and Central Hills ASSI

 Greeba Mountain and Central Hills ASSI
 (excluded area)

O/S Grid reference: SC335840
(approximate centre of site)

Designated: 23rd July 2008

Confirmed: 19th January 2009



The grid lines on this map form
part of the National Grid,
and are spaced at 1km intervals

Base Map is
Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Greeba Mountain and Central Hills

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Region: Isle of Man

Parishes: Braddan, German, Marown and Michael

Local authorities: Braddan, German, Marown & Michael Commissioners

Planning Authority: Department of Local Government and the Environment

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC27, SC28
National Grid Reference (centroid): SC335840 **Area:** 1,080.95ha (2671 acres)

Date notified: 23rd July 2008

Date confirmed: 19th January 2009

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its assemblage of upland breeding birds, and its upland wildlife habitats – dwarf shrub heath (heather moorland), acid grassland, mire (including blanket bog and acid hillside flushes), and river catchment.

Description and reasons for notification:

Greeba Mountain and Central Hills ASSI is a large area of upland moorland and grassland comprising five hills in a roughly southwest to northeast line: Greeba Mountain, Slieau Ruy, Lhargee Ruy, Colden and Slieau Maggle. The majority of the area is covered with heather moorland, interspersed with blanket bog and flush, acid grassland and small streams. The streams form the head of a large catchment including the Blaber River (a tributary of Glen Helen and the River Neb), the River Glass and associated Baldwin trout fisheries, and the Greeba River (a tributary of the River Dhoo which runs eventually to Douglas). The Central Hills area thus has a strong ecological influence on wetland habitats throughout the central region of the Island.

The majority of the habitat consists of dwarf shrub heath, i.e. heather-dominated moorland. Some parts of the area are wetter than others, with a full range of wet and dry heath present. Species include ling *Calluna vulgaris* (which dominates the majority of the site), bell heather *Erica cinerea*, cross-leaved heath *Erica tetralix* (on the wetter ground), bilberry *Vaccinium myrtillus*, common cottongrass *Eriophorum angustifolium*, heath grass *Danthonia decumbens*, heath rush *Juncus squarrosus*, deergrass *Trichophorum cespitosum*, green-ribbed sedge *Carex binervis*, and common lousewort *Pedicularis sylvatica*. In total, over 900 hectares of dry dwarf shrub heath has been recorded on the Central Hills ASSI, amounting to 15% of the total area for this habitat on the Island. Whilst broadly categorised as “dry” heath, there is a diverse range of water levels, with frequent wet areas throughout the ASSI on both the plateaus and the slopes.

Acid grassland occurs throughout the ASSI, interspersed with bog and heath. Many heath and bog plants grow amongst the grasses, which are dominated by mat grass *Nardus stricta*, brown bent *Agrostis vinealis*, common bent *Agrostis capillaris*, red fescue *Festuca rubra* and purple moor-grass *Molinia caerulea*. On the lower-lying parts of the area sweet vernal-grass *Anthoxanthum odoratum* is abundant. Other grassland species include tormentil *Potentilla erecta*, heath milkwort *Polygala serpyllifolia*, heath bedstraw *Galium saxatile*, common bird’s-foot trefoil *Lotus corniculatus*, common dog’s-violet *Viola riviniana*, sorrel *Rumex acetosa*, wild thyme *Thymus polytrichus*, common cat’s-ear *Hypochoeris radicata*, heath woodrush *Luzula multiflora* and heath spotted-orchid *Dactylorhiza maculata****.

The Central Hills contain 20% of the total area recorded as acid/neutral flush habitat on the Island, interspersed with blanket bog. These mire habitats occur in a complex mosaic with acid grassland and heath. Species present include round-leaved sundew *Drosera rotundifolia*, ling *Calluna vulgaris*, lousewort *Pedicularis sylvatica*, crowberry *Empetrum nigrum*, common cotton-grass *Eriophorum angustifolium*, hare’s-

tail cottongrass *Eriophorum vaginatum*, bog asphodel *Narthecium ossifragum*, velvet bent *Agrostis canina*, heath spotted-orchid *Dactylorhiza maculata****, bulbous rush *Juncus bulbosus*, purple moor-grass *Molinia caerulea*, many-stalked spike-rush *Eleocharis multicaulis* and star sedge *Carex echinata*. *Sphagnum* moss is frequent in the wettest areas, blanketing some parts and forming large patches or cushions of moss on the drier, grassier areas. Mosses present include *Sphagnum palustris* agg., *Sphagnum auriculatum*, *Sphagnum capillifolium*, *Sphagnum cuspidatum* and *Polytrichum commune*. In the more low-lying parts of the site, particularly at the north end, the diversity of mire species is higher, with plants characteristic of valley mire and fen meadows as well as bog species. Additional species found in the lower areas include sharp-flowered rush *Juncus actutiflorus*, lesser spearwort *Ranunculus flammula*, devil’s-bit scabious *Succisa pratensis*, carnation sedge *Carex panicea*, yellow-sedge *Carex viridula* subsp. *oedocarpa*, flea sedge *Carex pulicaris*, marsh lousewort *Pedicularis palustris*, jointed rush *Juncus articulatus* and greater bird’s-foot trefoil *Lotus pedunculatus*. Frogs *Rana temporaria*** breed in considerable numbers in the wetter areas.

Greeba Mountain and Central Hills ASSI supports a wide range of characteristic upland breeding birds. These include Hen harrier *Circus cyaneus**, Red grouse *Lagopus lagopus*, Raven *Corvus corax*, Curlew *Numenius arquata**, Stonechat *Saxicola torquata* and Snipe *Gallinago gallinago*. Whinchat *Saxicola rubetra** and Wheatear *Oenanthe oenanthe* are known to have bred on the ASSI in the recent past, and nearby breeding Peregrine *Falco peregrinus* and Long-eared owl *Asio otus** rely on the ASSI as hunting territory. In addition to the important breeding assemblage of upland birds, the area supports many other species, including breeding Dunnock *Prunella modularis*, Grasshopper warbler *Locustella naevia**#, Grey wagtail *Motacilla cinerea*, Skylark *Alauda arvensis**#, Songthrush *Turdus philomelos**#, Linnet *Carduelis cannabina*, Kestrel *Falco tinnunculus**, Lesser redpoll *Carduelis cabaret*, Reed bunting *Emberiza schoeniclus**#. Non-breeding birds which benefit from the area as a feeding ground include Fieldfare *Turdus pilaris*, Golden plover *Pluvialis apricaria**, Jack snipe *Limnocryptes minimus**, Redwing *Turdus iliacus*, Snow bunting *Plectrophenax nivalis*, Sparrowhawk *Accipiter nisus**, Swift *Apus apus** and Woodcock *Scolopax rusticola**.

* = Protected under Schedule 1 of the Wildlife Act 1990

= Birds of Conservation Concern Red List species ¹

** = Protected under Schedule 5 of the Wildlife Act 1990

*** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

Blanket bog is a Priority Annex 1 habitat under the EU Habitats Directive; Northern Atlantic wet heaths with *Erica tetralix* and European dry heaths are both Annex 1 habitats. These habitats are all internationally rare and/or threatened.

¹ Gregory Richard D, Wilkinson Nicholas I, Noble David G, Robinson James A, Brown, Andrew F, Hughes, Julian, Proctor Deborah, Gibbons David W, and Galbraith Colin A. (2002) The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002-2007. **British Birds** Vol 95 No 9 Pages 410-448

Site Name: Greeba Mountain and Central Hills

Operations likely to damage the special interest of the site

Standard reference number†	Type of operation
1	Cultivation including ploughing, rotovating, harrowing and reseeding
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling,.
24	Modification of natural or man-made features (including cave entrances),

	clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora.
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, bird, amphibian, fish or invertebrate.

Jurby Airfield Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**

Reillys Ellan Vannin



Department of Agriculture,
Fisheries and Forestry

JURBY AIRFIELD AREA OF SPECIAL SCIENTIFIC INTEREST

Area: 63.04 ha


Site Boundary

Designated: July 20th 2004
Varied and confirmed:
18th January 2005

The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

100 0 100 Meters

Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Area: Isle of Man

Site name: JURBY AIRFIELD

Parish: Jurby

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

National Grid Reference: SC 362983

Area: approximately 63.04 ha. (155.7 acres)

Ordnance Survey Sheets: 1:50,000 : Map 95

1:10,560 : SC 39 NE

1:25,000 Isle of Man Public Rights of Way and Outdoor Leisure Map (North)

Date notified: 20th July 2004, varied and confirmed 18th January 2005

Purpose

This site is designated for its semi-improved neutral grassland of the type crested dog's-tail and knapweed *Cynosurus cristatus* - *Centaurea nigra* lowland hay meadow.

Description:

A large, open and flat area of species-rich, neutral grassland on sandy soil, lying in between the airfield runways. The grassland is managed as a hay meadow. The grassland is variously dominated by red fescue *Festuca rubra*, Yorkshire fog *Holcus lanatus* and sweet vernal-grass *Anthoxanthum odoratum*, with frequent crested dog's-tail *Cynosurus cristatus*, common mouse-ear *Cerastium fontanum*, common bird's-foot-trefoil *Lotus corniculatus*, field wood-rush *Luzula campestris*, bulbous buttercup *Ranunculus bulbosus*, yarrow *Achillea millefolium*, yellow rattle *Rhinanthus minor*, ribwort plantain *Plantago lanceolata*, heath wood-rush *Luzula multiflora*, common bent-grass *Agrostis capillaris* and common sorrel *Rumex acetosa*.

Floral interest includes a number of species restricted to relatively unimproved sites on free-draining neutral or calcareous soils, which, in addition to bulbous buttercup and yellow rattle, includes lady's bedstraw *Galium verum*, spring sedge *Carex caryophyllaea*, crested hair-grass *Koeleria macrantha*, eyebright *Euphrasia nemorosa*, kidney vetch *Anthyllis vulneraria* and ox-eye daisy *Leucanthemum vulgare*.

Other notable plants are the squirrel-tail fescue *Vulpia bromoides*, and common centaury *Centaureum erythraea*.

The site is a stronghold for breeding skylarks *Alauda arvensis*^{***}. Other breeding bird species include the linnet *Carduelis cannabina*^{**}, grey partridge *Perdix perdix*^{**}, corncrake *Crex crex*^{*/**} and curlew *Numenius arquata*^{*}. In the winter flocks of golden plover *Pluvialis apricaria*^{*} and lapwing *Vanellus vanellus*^{*} have been recorded on the wider airfield. Visiting non-breeders include hen harrier *Circus cyaneus*^{*/**} and, peregrine *Falco peregrinus*^{*}.

Other information:

The location is known to have had significant prehistoric and Viking occupation. RAF Jurby survives as significant groups of buildings, perimeter defences and the runways.

Full species lists are available from the Biological Records Centre, Manx Museum, Douglas.

^{*} species on Schedule 1 of the Wildlife Act 1990

^{**} Red List Bird of Conservation Concern

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Jurby Airfield

Operations likely to damage the special interest of the site

Type of operation

1. Cultivation including ploughing, rotovating, harrowing and reseedling.
2. Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3. Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4. Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime, including conversion from hay making to silage or cessation of mowing, (except within 2 m of the edge of the runways or race tracks, access track for ambulances by route agreed with DAFF).
5. Application of manure, fertilisers and lime.
6. Application of pesticides, including herbicides (weedkillers).
7. Dumping, spreading or discharge of any materials.
8. Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9. The release into the site of any wild, feral or domestic animals **, plant or seed.
10. The killing or removal of any wild animal **, including pest control.
11. The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12. The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13. Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14. The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15. Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20. Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21. Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22. Storage of materials.
23. Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24. Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, infilling of pits, and quarries.
25. Removal of geological specimens, including rock samples, minerals and fossils.
26. Use of vehicles or craft likely to damage or disturb features of interest, (except on the runways and race tracks or by emergency vehicles whilst attending emergencies)
27. Recreational or other activities likely to damage features of interest, (except on runways and race tracks).

28. Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

**** “animal” includes any mammal, reptile, bird, fish or invertebrate.**

Operations 16-19 excluded as not relevant to this site.

January 18th 2005

Langness, Derbyhaven & Sandwick Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmltaght, Bee as Eirinys

This Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



Isle of Man
Government
Reillys. Ellan Vannin



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Area: Isle of Man

Parishes: Malew

Site name: Langness, Sandwick and Derbyhaven

Status : Area of Special Scientific Interest (ASSI). Notified under Section 27 of the Wildlife Act 1990.

Planning Authority: Department of Local Government and the Environment

Local authority: Malew Commissioners

National Grid Reference: SC 291681 - SC276652 **Area:** 310 ha. (765 acres)

Ordnance Survey Sheet: 1:50,000 OS Landranger 95 1:10,560 SC 26 NE
1:25,000 Isle of Man Public Rights of Way and Outdoor Leisure Map

Date notified: October 19th 2000 **Varied and confirmed:** 27th April 2001

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its fauna, flora, geology, geomorphology and landscape.

Description and reasons for notification:

Fauna

Langness is of outstanding scientific importance as the only place in the British Isles where the lesser mottled grasshopper, *Stenobothrus stigmaticus*** has been recorded. The grasshopper is found in the short grass around rock outcrops.

Langness is the most important Manx locality for wintering and migrant waders, and wildfowl. They feed mainly in the intertidal zone but also on the upper shore and coastal grassland. The sheltered coast is important for ducks (particularly teal *Anas crecca*, widgeon *Anas penelope* and shelduck *Tadorna tadorna*) and waders (lapwing *Vanellus vanellus**, oystercatcher *Haematopus ostralegus* and ringed plover *Charadrius hiaticula*) as well as visiting rarities. Significant numbers of waders roost here and large feeding flocks of curlew *Numenius arquata* and golden plover *Pluvialis apricaria* have been recorded in the area. Raven *Corvus corax* and chough *Pyrrhocorax pyrrhocorax** nest on the rocky coast and lapwing* have bred on the saltmarsh. In winter there are heaps of seaweed on the strandline in which choughs dig and feed. Its location as a southern promontory makes Langness a key site for migrant birds and a great variety of species have been recorded, comparable with the range of species on the Calf, which is a British Isles Bird Observatory. These include wryneck *Jynx torquilla**, little auk *Alle alle*, goosander *Mergus merganser* and little egret *Egretta garzetta*.

The birds are the main vertebrates but frogs *Rana temporaria*** and common lizards *Lacerta vivipara* ** have been recorded. Around the shores grey seals *Halichoerus grypus*** haul out on the rocks.

The peninsula also attracts migrant butterflies (red admiral *Vanessa atlanta*, painted lady *Vanessa cardui* and clouded yellow *Colias croceus*) and moths (silver-Y *Autographa gamma*). The heath and grassland are habitats for a wide variety of invertebrates, including Grey *Hadena caesia mananii*** and pod lover moths *Hadena perplexa capsophila***. The native butterflies include dark green fritillary *Argynnis*

aglaja, grayling *Hipparchia semele*, common blue *Polyommatus icarus* and small copper *Lycaena phlaeas*. A total of 15 species of butterfly have been recorded. Other invertebrates recorded include green tiger beetle *Cicindela campestris*, great yellow bumble bee *Bombus distinguendus*, the saltmarsh centipede *Schendyla peyerimhoffi* and a rare moth *Pyrausta sanguinalis**** which also occurs on the Ayres.

Flora

The dominant vegetation is believed to have been gallic heath (Manx gorse *Ulex gallii* and heathers, *Calluna vulgaris* and *Erica cinerea*) and dune vegetation (marram grass- *Ammophila arenaria* and sand sedge *Carex arenaria*). Now relatively little heath remains in pockets on the shallower soils. Where the ground is not intensively managed as golf greens, the vegetation is un-improved grassland, heath and scrub on limestone, Manx slate and conglomerate and on sand.

One of the largest areas of saltmarsh on the Island is on the Castletown side of the peninsula. Such species as sea club-rush *Bolboschoenus maritimus*, grass-leaved orache *Atriplex littoralis*, maritime rush *Juncus maritimus* and sea aster *Aster tripolium* occur there. Interesting coastal grassland plants include spring squill *Scilla verna* and Danish milk vetch *Astragalus danicus*. Around the margin of Langness freshwater seepage creates flushes of interesting plants; black bog rush *Schoenus nigricans*, ragged robin *Lychnis flos-cuculi*, celery leaved buttercup *Ranunculus sceleratus* and bog pimpernel *Anagallis tenella*. A water tank between the cliffs and the golf course, with planted white water lily *Nymphaea alba*, attracts damselflies (*Enallagma cyathigerum*, *Ischnura elegans* and *Pyrrhosoma nymphula*) and adds to the diversity of habitats. The strandline vegetation includes the dramatic sea kale *Crambe maritima*, sea radish *Raphanus raphanistrum ssp maritimus* and a variety of more common flowering plants.

Geology, geomorphology and landscape

The Langness peninsula is an example of a "tombolo", a rocky island joined by a sandy spit to the main island. The area has geological formations of limestone, Manx slate and conglomerate overlain with sand and shingle at the land-ward end. This combination occurs nowhere else on the Island. Dramatic rock arches of conglomerate or "pudding stone" occur in the south east. This conglomerate overlies Manx slates illustrating an "unconformity" or a gap in the geological sequence. Within the conglomerate there is a dyke of basalt: a volcanic intrusion. The variety of rocks leads to soils which are diverse in structure and acidity.

Other information:

The land has a history of copper mining, lime burning and farming. The importance of human history in the area has been reflected in the designation of several listed monuments under the terms of the Manx Museum and National Trust Act 1959.

This site forms a cohesive ecological unit which is important for its terrestrial ecology and together with the marine and inter-tidal area of Derbyhaven and Sandwick bays, meets the criteria for designation as a wetland of international importance under the RAMSAR Convention. Much of the peninsula and its adjoining foreshores have been a designated Bird Sanctuary since 1936.

Full species lists are available from the Biological Records Centre, Manx Museum, Douglas.

* listed in Schedule 1 of Wildlife Act 1990

** listed in Schedule 5 of Wildlife Act 1990

*** Proposed Red Data Book species

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Langness, Sandwick and Derbyhaven

Operations likely to damage the special interest of the site

Type of operation


1. Cultivation including ploughing, rotovating, harrowing and reseeding (excluding that required to manage the tees, greens, fairways and practice area*).
2. Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3. Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4. Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing), (excluding that required to manage the tees, greens, fairways and practice area).
5. Application of manure, fertilisers and lime (excluding that required to manage the tees, greens, fairways and practice area).
6. Application of pesticides, including herbicides (weedkillers) (excluding herbicides required to manage the tees, greens, fairways and practice area).
7. Dumping, spreading or discharge of any materials, (excluding that required to manage the tees, greens, fairways and practice area).
8. Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9. The release into the site of any wild, feral or domestic animals **, plant or seed (excluding domestic dogs, and excluding grass seed sown on tees, greens, fairways and practice areas).
10. The killing or removal of any wild animal **, including pest control (excluding that required to manage the tees, greens, fairways and practice area).
11. The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf, (excluding that required to manage the tees, greens, fairways and practice area).
12. The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13. Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains), (excluding that required to manage the tees, greens, fairways and practice area).
14. The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes) (excluding irrigation required to manage the tees, greens, fairways and practice area).
15. Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16. Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17. Reclamation of land from sea, estuary or marsh.
18. Bait digging on inter-tidal areas, (if it is shown to be damaging).
19. Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
20. Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil, (excluding sand removal for bunkers).

21. Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22. Storage of materials (excluding that required to manage the tees, greens, fairways and practice area where locations have been agreed with DAFF).
23. Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling, (excluding any locations agreed with DAFF).
24. Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, infilling of pits, and quarries.
25. Removal of geological specimens, including rock samples, minerals and fossils.
26. Use of vehicles or craft likely to damage or disturb features of interest,
(excluding vehicles required by those taking part in a game of golf or to manage the tees, greens, fairways and practice area, or in other locations agreed with DAFF, or for overnight parking of camper vans, when in locations agreed with DAFF, outside bird breeding season - ie excluding 1st May to 16th July).
27. Recreational or other activities likely to damage features of interest (excluding walking, with or without dogs).
28. Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

* tees, greens, fairways and practice area are areas indicated provisionally on accompanying prints of aerial photographs.

** "animal" includes any mammal, reptile, bird, fish or invertebrate.

April 2nd 2001



Isle of Man

Ellan Vannin

AT 4 of 1959

**MANX MUSEUM AND NATIONAL TRUST
ACT 1959**



Isle of Man

Ellan Vannin

MANX MUSEUM AND NATIONAL TRUST ACT 1959

Index

Section	Page
1 Short title	5
2 Interpretation of terms	5
<i>Constitution</i>	7
3 Constitution of the Trust	7
3A Legal status	8
3B Who comprises the Trust	8
3C Application of particular provisions of Statutory Boards Act 1987	8
<i>Appointment and term of office of trustees</i>	8
3D Minister's appointee	8
3E Appointment of chairperson and other trustees	8
3F Requirements for making appointment recommendation	9
3G Term of office of trustees	9
3H General powers concerning appointments not affected	10
3I Vice-chairperson	10
<i>Manx Museum</i>	10
4 [Repealed]	10
5 Manx Museum	10
6 Trust may obtain specimens of birds, fishes, animals, or eggs	11
<i>Ancient Monuments</i>	11
7 Trust to declare ancient monument to which Act shall apply	11
8 Gift, etc, of ancient monument to Trust	11
9 Sale or gift by limited owner	11
10 Guardianship of ancient monuments	11
11 Protection of ancient monuments	12
12 Effect of preservation order	13
13 Notice to owners of such monuments	13
14 Voluntary contributions	14
15 Injury to ancient or historic monument	14
16 Schemes for preserving amenities of ancient monuments	14
17 Power of Trust to enter upon lands	16
18 Service of documents	16

19	Placing of tablets on ancient monuments of historical interest	16
20	Reports of finding of archaeological objects.....	17
21	Restriction of export of archaeological objects.....	17
21A	Restrictions on use of metal detectors	18
22	Prohibition of injury, etc, to archaeological objects.....	19
23	Restriction of excavation for archaeological purposes.....	19
	<i>Manx National Trust</i>	20
24	Powers and duties of Manx Museum and National Trust	20
	<i>General</i>	20
25	Power to acquire real estate, etc	20
25A	Certain property of Trust to be inalienable	21
25B	Power to enter into agreements restricting use of land	22
26	Powers exercisable over Trust	22
26A	Trustees' duty to consult on heritage matters.....	23
27	Power to make, etc, bye-laws	23
28	Trust to be exempt from Income Tax.....	23
29	Ministerial meetings with stakeholders	23
30	Crown rights	24
31	Commencement of Act	24
	<i>Transitional provisions for Manx Museum and National Trust (Amendment) Act 2011</i>	24
32	Definitions for ss 32 to 35	24
33	Appointment of new chairperson and other trustees	24
34	Existing trustees continue until appointments take effect	25
35	Expiry of transitional provisions.....	25
	FIRST SCHEDULE	27
	SECOND SCHEDULE	27
	THIRD SCHEDULE	29
	ENDNOTES	31
	TABLE OF LEGISLATION HISTORY	31
	TABLE OF RENUMBERED PROVISIONS	31
	TABLE OF ENDNOTE REFERENCES	31

**Isle of Man***Ellan Vannin*

MANX MUSEUM AND NATIONAL TRUST ACT 1959

Received Royal Assent: 20 November 1959
Passed: 19 January 1960
Commenced: 19 January 1960

[AN ACT to consolidate and revise the Manx Museum and National Trust Acts, 1886 to 1951.]¹

GENERAL NOTES

1. The maximum fines in this Act are as increased by the *Manx Museum Act 1984* Sch and by the *Criminal Justice (Penalties, Etc.) Act 1993* s 1.
2. References to hard labour are to be construed in accordance with the *Custody Act 1995* Sch 3 para 1.

1 Short title

This Act may be cited for all purposes as the Manx Museum and National Trust Act, 1959.

1A. Main purposes of Act

The main purposes of this Act are to —

- (a) provide for the Manx Museum and National Trust (*Thie-tashtee Vannin as y Barrantys Ashoonagh*); and
- (b) promote public confidence in the Trust and its relationship with the Government.²

2 Interpretation of terms

- (1) In the construction of this Act the following expressions shall have, unless inconsistent with the context, the meanings hereby assigned to them (that is to say): —

“**ancient monument**” shall include any monument or group of monuments, and any part or remains of a monument, or group of monuments, the preservation of which is in the opinion of the trust a matter of public interest by reason of the historic, architectural, traditional, artistic,

scientific or archaeological interest attaching thereto, and the site of any such monument or of any remains thereof, and any part of the adjoining land, which may in the opinion of the Trust be required for the purpose of fencing, covering in, or otherwise preserving the monument from injury, or for the purpose of preserving the amenities of a monument or for providing or facilitating access thereto or for the exercise of proper control or management with respect thereto. And shall include any dwelling house, building or erection, and the fittings, furniture and contents thereof illustrative of the mode of life, occupation, habits and customs of the people of this Island in bygone days;

“appointment recommendation” has the meaning given under section 3E(1) and (3);³

“archaeological object” means any chattel including ancient human and animal remains, whether in a manufactured or partly manufactured or unmanufactured state which by reason of the archaeological interest attaching thereto, or of its association with any Manx historical event or person, has a value substantially greater than its intrinsic (including artistic) value, and notwithstanding the provisions of section 30 of this Act the said expression includes all treasure trove;⁴

“chairperson” means the Trust’s chairperson, appointed under section 3E(1);⁵

“Council of Ministers appointee” has the meaning given under section 3E(1);⁶

“Department” means the Department of Community, Culture and Leisure;⁷

“entitled” means beneficially entitled and the expression **“land”** means land which is the site of an ancient monument whether the land is or is not subject to encumbrance;

“inspector” means an inspector of ancient monuments appointed by the Trust;

“maintenance” and **“maintain”** shall include the fencing, repairing and covering in of a monument and the doing of any other act or thing which may be required for the purpose of repairing the monument or protecting it from decay or injury or the restoration of the same to its known or assumed original condition when erected, and without prejudice to the generality of the foregoing provisions power to make examination of the monument or open up the monument and make excavations therein for the purpose of examination or otherwise;

“member”, of the Department, means a person who, under section 2 of the Government Departments Act 1987, is a member of the Department;⁸

“Minister” means the Minister for the Department;⁹

“Minister’s appointee” has the meaning under section 3D(1);¹⁰

“monument” shall include any building, structure or other work, whether above or below the surface of the land, other than an ecclesiastical

building for the time being used for ecclesiastical purposes, and any cave or excavation;

“**Museum**” means the land and buildings comprising the Public Museum, Library and Art Gallery comprised in a Conveyance made 25th October 1921 between (1) the Trustees of Henry Bloom Noble deceased and (2) the Government Property Trustees;¹¹

“**owner**” shall include —

- (a) any person entitled for an estate in fee to the possession or receipt of the rents and profits of any land;
- (b) any person entitled under any existing or future will or other instrument for the term of his own life or the life of any other person to the possession or receipt of the rents and profits of any land and tenure;
- (c) the guardians of any minor or the receiver of a mental patient;¹²
- (d) any body corporate or corporation sole or any trustees for charitable public or private persons or purposes in whom any property may be vested;
- (e) the Crown Estate Commissioners, any Department or Statutory Board and any other Government or Tynwald body, Board or committee in whom any property may be vested by Act, order, gift, conveyance or otherwise;¹³

“**the Trust**” means the Manx Museum and National Trust;

“**trustee**” means the Minister’s appointee or a Council of Ministers appointee (including the chairperson).¹⁴

- (2) References in any enactment or other document to the Manx Museum and Ancient Monuments Trustees shall be construed as reference to the Manx Museum and National Trust.
- (3) In this Act —
 - (a) a reference to a function includes a power; and
 - (b) a reference to performing a function includes exercising a power.¹⁵

Constitution

3 Constitution of the Trust

- (1) The Manx Museum and National Trust (*Thie-tashtee Vannin as y Barrantys Ashoonagh*) is continued in existence.
- (2) The Trust may continue to perform its functions under this Act under the name Manx National Heritage (*Eiraght Ashoonagh Vannin*).¹⁶

3A Legal status

The Trust continues to be a body corporate with perpetual succession and to have a common seal.

3B Who comprises the Trust

The Trust is comprised of the following persons as its trustees —

- (a) a chairperson;
- (b) at least 8, but no more than 14, other trustees.

3C Application of particular provisions of Statutory Boards Act 1987

- (1) Schedule 2 (exercise of functions of Statutory Boards) of the *Statutory Boards Act 1987*, other than paragraphs 1, 3(2)(a), 6 and 7, applies to the Trust and the trustees.
- (2) The provisions applied under subsection (1) apply as if the Trust were a Statutory Board, and with other necessary changes.

Appointment and term of office of trustees

3D Minister's appointee

- (1) The Minister is to appoint one person as a trustee (the “**Minister's appointee**”).
- (2) Only a member of the Department (but not the Minister) can be the Minister's appointee.
- (3) The Minister must tell the Council of Ministers about the appointment, or any change in the appointment, of the Minister's appointee.

3E Appointment of chairperson and other trustees

- (1) The persons who are to be the chairperson and the trustees, other than the Minister's appointee, (each a “**Council of Ministers appointee**”) are to be appointed by the Council of Ministers on the recommendation (an “**appointment recommendation**”) of an interview panel.
- (2) The interview panel is comprised of the following persons as its members —
 - (a) the Minister, as its chairperson;
 - (b) the Minister's appointee;
 - (c) two other persons appointed by the Council of Ministers
- (3) If appointments have been made under subsection (1) and they are still in force, the Trust may recommend (also an “appointment

recommendation”) to the Council of Ministers the appointment of further trustees.

- (4) However the Trust can not make an appointment recommendation —
 - (a) for the chairperson; or
 - (b) for a person if, were the person to be appointed, a majority of the trustees would be persons recommended under subsection (3).
- (5) Neither the Minister nor a member of Tynwald can be appointed to the interview panel under subsection (2)(c) or be a Council of Ministers appointee.
- (6) Despite subsections (1) and (3), the appointment of a Council of Ministers appointee does not take effect until it is approved by Tynwald.

3F Requirements for making appointment recommendation

The following apply for making an appointment recommendation —

- (a) the areas of interest of the Trust and the Department must be considered;
- (b) appropriate professional advice must be obtained.

3G Term of office of trustees

- (1) The Minister’s appointee —
 - (a) holds office as a trustee at the Minister’s will, but goes out of office on ceasing to be a member of the Department or on resignation by written notice to the Minister; and
 - (b) is eligible for reappointment.
- (2) Council of Ministers appointees may be appointed at different times and for different terms.
- (3) A Council of Ministers appointee —
 - (a) may be appointed for any term up to 5 years, but goes out of office on becoming a member of Tynwald or on resignation by written notice to the Council of Ministers; and
 - (b) is eligible for reappointment.
- (4) However, a Council of Ministers appointee can not be appointed as the chairperson or as a trustee for more than 2 consecutive terms.
- (5) Subsection (4) does not prevent a Council of Ministers appointee from being appointed to the following after the 2 consecutive terms end —
 - (a) for the chairperson - as a trustee other than the chairperson;
 - (b) for a trustee other than the chairperson - as the chairperson.

3H General powers concerning appointments not affected

To avoid any doubt, sections 20 and 22 of the Interpretation Act 1976 apply for an appointment under section 3D or 3E.

Vice-chairperson

3I Vice-chairperson

- (1) The trustees must, by ballot, elect a trustee other than the chairperson or the Minister's appointee (an "eligible trustee") to be the Trust's vice-chairperson.
- (2) Elections for vice-chairperson must be held annually or on any earlier vacancy in that office.
- (3) The office becomes vacant if the person holding it ceases to be an eligible trustee or resigns by written notice to the chairperson.

Manx Museum

4 [Repealed]¹⁷

5 Manx Museum

- (1) The Museum shall be held by the Trust as a Public Museum, Library and Art Gallery and also for any purpose of a historical, scientific, archaeological, ecclesiastical, physical, industrial or educational character with or incidental to a Public Museum, Library or Art Gallery.
- (2) The Trust shall keep the Museum and any additions thereto in good order and repair.
- (3) The Museum shall be used for storing, classifying, and exhibiting objects of art, or of antiquarian, scientific, literary, historical, or educational interest, whether the property of such Trust or of any other person or body.
- (4) All persons or public bodies owning or having the custody or control of articles of any description suitable to be kept, stored, or exhibited in the Museum, are hereby authorised and empowered to transfer the same, either absolutely or on such conditions as they may deem proper, to the Trust, or may, with the consent of the Trust, deposit them on loan.
- (4A) Public records, within the meaning of the *Public Records Act 1999*, may not be transferred to the Trust under subsection (4) above otherwise than on loan and in accordance with section 3 of that Act.¹⁸

Museum to be open to the public

- (5) The Museum shall be open to the public on such days and for such hours and subject to such conditions as the Trust may determine, subject to the approval of Tynwald.
- (6) The Trust shall have power to make such reasonable charges as the Trust may determine, for the inspection of any document which has been removed from the General Registry to the Museum under section 6 of the *Public Records Act 1999* or for the supply of a copy of any such document.¹⁹
- (7) [Repealed]²⁰

6 Trust may obtain specimens of birds, fishes, animals, or eggs

Notwithstanding any enactment to the contrary, it shall be lawful for any person, authorised in writing by the Trust, and acting on their behalf, to take and kill any wild bird, fish, or other animal, or to take the eggs of any wild birds, which may be required by the Trust for the purposes of the Museum, without being liable to any proceedings or penalty for so doing.

*Ancient Monuments***7 Trust to declare ancient monument to which Act shall apply**

The Trust may, from time to time, by Order, with the consent of the owner, declare that any monument shall be deemed to be an ancient monument to which this Act applies.²¹

8 Gift, etc, of ancient monument to Trust

The owner of any ancient monument may by deed or will give, devise or bequeath to the Trust such ancient monument and it shall be lawful for the Trust to accept such gift, devise or bequest if they think it expedient to do so.

9 Sale or gift by limited owner

Any sale, gift, devise or bequest to the Trust of an ancient monument by an owner who is not the owner in fee simple in possession for his own benefit of such property shall include and convey to the Trust the absolute ownership of such ancient monument.

10 Guardianship of ancient monuments

- (1) The owner of any ancient monument may by deed under his hand constitute the Trust guardians of such monument and the Trust may accept such guardianship if they think it expedient to do so.

- (2) Every person deriving title to any ancient monument from, through or under any owner who has constituted the Trust guardians of the monuments under this section shall be bound by the deed executed by the owner for that purpose, notwithstanding that such owner was not the absolute holder in fee simple in possession of such ancient monument, and the execution of any such deed by any person who was not such absolute owner shall not render him subject to any liability on account of any depreciation of property attributable thereto.
- (3) The owner of an ancient monument of which the Trust are guardians shall, except save as in this Act expressly provided, have the same estate right, title and interest in and to such monument in all respects as if the Trust had not been constituted guardians thereof, but the Trust may maintain such monument and shall for the purposes of such maintenance at all reasonable times have access to such monument by themselves or their workmen for the purpose of inspecting and maintaining it as in this Act defined, and may bring to such monument such materials and do such acts and things as may be required for the maintenance thereof.

11 Protection of ancient monuments

- (1) If the Trust is of opinion that any ancient monument is in danger of destruction or removal or damage from neglect or injudicious treatment, and that the preservation of the monument is of national importance, the Trust may make an order (in this Act referred to as 'a preservation order') placing the monument under the protection of the Trust.
- (2) Where the Trust has reason to believe that any ancient monument is in danger as aforesaid they may themselves, or by any person authorised in writing by them, enter at any reasonable time upon any premises for the purpose of enabling them to determine by inspection of the monument whether it is proper for them to make an order under this section. Provided that unless the Trust consider the inspection of the monument is a matter of immediate urgency, not less than seven clear days' notice to the occupier of the premises shall be given of their intention so to enter upon them.
- (3) On the making of a preservation order the Trust shall cause a copy of such order to be fixed on some conspicuous part of the monument and shall also cause a copy of the order together with a notice stating the effect thereof to be served upon the owner and upon the occupier of the monument.
- (4) The preservation order shall have effect for a period of eighteen months after the date on which it is made. On the expiration of that period and for a period of one month thereafter the owner or occupier of the premises affected by such order may apply by petition to Tynwald to have such order set aside. If —

- (a) at the first sitting of Tynwald held after the presentation of such petition or any sitting to which it may be continued or adjourned Tynwald shall not pass a resolution setting the same aside, or
 - (b) no such petition is presented within either of the said periods,²²
- the monument to which such order refers shall thenceforth vest in the Trust in all respects as if the Trust had been nominated guardians of such monument by the owner thereof.²³
- (5) Not less than fourteen days' notice in writing shall be given by the owner and occupier of any ancient monument to the Trust of the sitting of Tynwald at which it is proposed to move for a resolution setting aside a preservation order. And the Trust may appear before Tynwald and be heard in opposition to the resolution to set aside the preservation order. The costs in connection with any such petition shall be in the discretion of Tynwald.

12 Effect of preservation order

While a preservation order is in force the monument to which the order relates shall not be demolished or removed, nor shall any additions or alterations be made thereto, or any work carried out in connection therewith, except with the written consent of the Trust.

13 Notice to owners of such monuments

- (1) The Trust shall from time to time cause to be prepared and published a list containing such monuments as are considered by the Trust as being monuments the preservation of which is of national importance, or should otherwise be included in such list, and shall inform the owners and occupiers of such monuments of their intention and of the penalties which may be incurred by a person guilty of an offence under the next succeeding subsection.
- (2) Where an owner of an ancient monument which is included in any such list of monuments proposes to demolish or remove in whole or in part, structurally alter or make additions to the monument, he shall forthwith give notice of his intention to the Trust and shall not commence any work of demolition, removal, alteration or addition for a period of one month after having given such notice, and any person guilty of a contravention or non-compliance with this provision shall be liable on summary conviction at the suit of the Trust to a fine not exceeding £2,500 or to imprisonment for a term not exceeding one month or to both.
- (3) This section shall not apply to any structure which is occupied as a dwelling house by any person other than a person employed as a caretaker thereof or his family.

14 Voluntary contributions

The Trust shall have power to receive voluntary contributions towards the cost of the maintenance and preservation of any ancient monument and the amenities of such monument, and to enter into any agreement with the owner of any such monument, or with any other person, as to the maintenance and preservation of the monument and the cost thereof, notwithstanding that the Trust are not the owners or guardians of the monument, and may make regulations relating to any monument as to access or otherwise including the power to make charges for such access in cases where the monument is vested in the Trust.

15 Injury to ancient or historic monument

If any person injures or defaces any monument of which the Trust are the owners or guardians, or which is the subject of a preservation order, that person shall on summary conviction at the suit of the Trust be liable to a fine not exceeding £5,000 and in addition to the fine to pay such sums as the court by which he is tried may think just for the purpose of repairing any damages caused by him, or to imprisonment with or without hard labour for a term not exceeding one month, and for the purpose of this section the owner of an ancient monument of which the Trust are guardians or which is subject to a preservation order shall be liable in all respects as if he were not the owner thereof.

16 Schemes for preserving amenities of ancient monuments

- (1) For the purpose of preserving the amenities of any ancient monument, the Trust may, subject to the provisions of this section, prepare and confirm a scheme (hereafter in this Act referred to as ‘a preservation scheme’) for any area comprising or adjacent to the site of the monument, being an area to which, in the opinion of the Trust, it is necessary or expedient for that purpose that the scheme should apply.
- (2) Every preservation scheme shall define by reference to a map annexed thereto the area to which the scheme is applicable (hereafter in this Act referred to as ‘the controlled area’) and may provide for all or any of the following matters (that is to say) —
 - (a) for prohibiting or restricting the construction, erection or execution of buildings, structures and other works above ground within the controlled area, or the alteration or extension of such buildings, structures or works in such manner as materially to affect their external appearance;
 - (b) for prescribing the position, height, size, design, materials, colour and screening, and otherwise regulating the external appearance of buildings, structures and other works above ground within the controlled area;

- (c) for prohibiting or restricting the felling of trees, quarrying and excavations within the controlled area;
 - (d) for otherwise restricting the user of land within the controlled area to such extent as may appear to the Trust to be expedient for the purpose of preserving the amenities of the monument;
 - (e) for such other matters as appear to the Trust to be incidental to or consequential on the foregoing provisions of this section or to be necessary for giving effect to these provisions.
- (3) The provisions of Part I of the Second Schedule to this Act shall have effect with respect to the confirmation, variation and revocation of preservation schemes.
- (4) Nothing in any preservation scheme shall affect any building, structure or other work above ground or any alteration or extension thereof, if it was constructed, erected or executed before the date when notice of intention to confirm the scheme was published in two newspapers printed and circulating in this Isle under Part I of the Second Schedule to this Act, and for the purpose of this provision a building, structure or other work, and any alteration or extension thereof shall be deemed to have been constructed, erected or executed before that date —
 - (a) if its construction, erection or execution was begun before that date; or
 - (b) if and so far as its construction, erection or execution was necessary for the purpose of performing a contract made before that date.
- (5) Any person whose property is injuriously affected by the coming into force of a preservation scheme shall be entitled to obtain compensation in respect thereof from the Trust, subject to the provisions of Part II of the Second Schedule to this Act.
- (6) If any person contravenes any provision of a preservation scheme for the time being in force he shall be liable on summary conviction at the suit of the Trust to a fine not exceeding £100 for every day on which the contravention occurs or continues.²⁴
- (7) If, after any person has been convicted of a contravention of a preservation scheme by reason that any building, structure or other work is not in conformity with the scheme, the contravention continues after the expiration of such period as the court before whom he was convicted may determine, the Trust shall have power to do all such acts as, in their opinion, are necessary for removing so much of the building, structure or work as is not in conformity with the scheme, or for making it conform with the scheme, and any expenses incurred by the Trust in so doing shall be recoverable summarily as a civil debt from the person convicted.
- (8) [Repealed]²⁵

17 Power of Trust to enter upon lands

- (1) Subject as hereinafter provided, an inspector or any person specially authorised in writing in that behalf by the Trust, after giving not less than fourteen days' notice in writing to the occupier of his intention so to do and on production of his authority if so required by or on behalf of the occupier, may enter for the purposes of investigation at all reasonable times upon any land which the Trust have reason to believe contains an ancient monument, and may make excavations in the land for the purposes of examination or may enter and remain for a reasonable time on any land for the purpose of making observations and collecting data and specimens of or relating to natural history, geology or any other matter within the purview of the Trust:

Provided that —

- (a) no person shall, under the powers conferred by this section, enter any dwelling house or any building, park, garden, pleasure ground or other land used for the amenity or convenience of a dwelling house except with the consent of the occupier; and
 - (b) no excavation shall be made under the said powers except with the consent of every person whose consent to the making of the excavation would, apart from this section, be required.
- (2) If any person wilfully obstructs or hinders any person duly authorised by the Trust in the exercise of the powers conferred by this section, he shall be liable on summary conviction at the suit of the Trust to a fine not exceeding £200.

18 Service of documents

- (1) Every document, required or authorised to be served under this Act upon the owner or occupier of an ancient monument, may be served either by delivering it or leaving it at the usual or last known place of abode of the person on whom it is to be served, or by sending it by post as a registered letter addressed to him at his last known place of abode, or, if that cannot be found, by fixing it on some conspicuous part of the monument.
- (2) Any such document may, as the case requires, be addressed to the 'owner' or 'occupier' of the monument (describing it) without further name or description.

19 Placing of tablets on ancient monuments of historical interest

The Trust may, with the consent of the owner of an ancient monument which is of historical interest, place or cause to be placed on such ancient monument a tablet, plate, inscription, or other notice, stating the facts in relation to such ancient monument which give rise to the historical interest thereof.

20 Reports of finding of archaeological objects

- (1) Every person who finds any archaeological object shall, within fourteen days after he has found such object, make a report of such finding to a member of the police on duty in the district in which such object was found or to an officer of the Trust, and shall when making such report state his own name and address, the nature or character of the said object and the time and place at which and the circumstances in which it was found by him, and shall also, and whether he has or has not made such report as aforesaid, and irrespective of the person to whom he has made such report (if any) give to any member of the police or officer of the Trust on request any information within his knowledge in relation to such object or the finding thereof and shall permit any member of the police or officer of the Trust to inspect, examine or photograph such object.
- (2) Every person who finds an archaeological object and —
 - (a) fails without reasonable excuse to make a report of such finding in accordance with this section, or
 - (b) makes under this section a report of such finding which is to his knowledge false or misleading in any material respect, or
 - (c) in contravention of this section fails or refuses to give to a member of the police or the officer of the Trust information in relation to such archaeological object or the finding thereof, or
 - (d) gives to a member of the police or an officer of the Trust information in relation to such archaeological object or the finding thereof which is to his knowledge false or misleading in a material respect,

shall be guilty of an offence under this section and shall be liable on summary conviction at the suit of the Trust to a fine not exceeding £1,000.

21 Restriction of export of archaeological objects

- (1) Any person who —
 - (a) removes an archaeological object from the Island to the United Kingdom; or
 - (b) exports an archaeological object from the Island to a country outside the United Kingdom; or
 - (c) sells an archaeological object for such removal or export,

without or otherwise than in accordance with a licence issued by the Trust under subsection (2) below, shall be guilty of an offence and shall be liable on summary conviction to imprisonment for a term not exceeding 6 months, or to a fine not exceeding £5,000 or a sum equal to two times the value of the goods, whichever is the greater, or to both.

- (2) The Trust may at their discretion issue to any person a licence to remove from the Island to the United Kingdom or to export from the Island to a country outside the United Kingdom any archaeological object and may make any such licence subject to such conditions and restrictions as they shall think proper.²⁶

21A Restrictions on use of metal detectors

[P1979/46/42]

- (1) If a person uses a metal detector in a protected area without the written consent of the Trust he shall be guilty of an offence and liable on summary conviction to a fine not exceeding £1,000.
- (2) In this section —
‘metal detector’ means any device designed or adapted for detecting or locating any metal or mineral in the ground; and
‘protected area’ means any place which is either —
 - (a) the site of an ancient monument or of any monument under the ownership or the guardianship of the Trust by virtue of this Act; or
 - (b) situated in an area which the Trust has by order designated as an area of archaeological importance for the purposes of this section.
- (3) An order made by the Trust under subsection (2) above, shall be laid before Tynwald as soon as may be after it is made, and if Tynwald at the sitting before which such order is so laid resolves that the order shall be annulled, the order shall thereupon cease to have effect.
- (4) The Trust may by resolution depute the ²⁷Chairperson or Vice-Chairperson of the Trust to exercise the power under subsection (2) above, to make an order designating areas of archaeological importance, subject to such conditions, exceptions and qualifications as may be specified in such resolution; and from the date specified in such resolution, the Chairman or Vice-Chairman may exercise such power, subject as aforesaid.
- (5) If a person without the written consent of the Trust removes any object of archaeological or historical interest which he has discovered by the use of a metal detector in a protected area he shall be guilty of an offence and liable on summary conviction to a fine not exceeding £5,000 or two times the value of the object, whichever is the greater.
- (6) A consent granted by the Trust for the purposes of this section may be granted either unconditionally or subject to conditions.
- (7) If any person —

- (a) in using a metal detector in a protected area in accordance with any consent granted by the Trust for the purposes of this section; or
- (b) in removing or otherwise dealing with any object which he has discovered by the use of a metal detector in a protected area in accordance with any such consent;

fails to comply with any condition attached to the consent, he shall be guilty of an offence and liable, in a case falling within paragraph (a) above, to the penalty provided by subsection (1) of this section, and in a case falling within paragraph (b) above, to the penalty provided by subsection (5) of this section.

- (8) In any proceedings for an offence under subsection (1) above, it shall be a defence for the accused to prove that he used the metal detector for a purpose other than detecting or locating objects of archaeological or historical interest.
- (9) In any proceedings for an offence under subsection (1) or (5) of this section, it shall be a defence for the accused to prove that he had taken all reasonable precautions to find out whether the place where he used the metal detector was a protected area and did not believe that it was.²⁸

22 Prohibition of injury, etc, to archaeological objects

- (1) No person shall injure, deface or destroy any archaeological object, nor shall any person alter any archaeological object otherwise than under and in accordance with a licence in that behalf granted under this section.
- (2) The Trust may, if they think fit, issue to any person a licence to alter a specified archaeological object in such manner, to such extent and subject to such conditions as are specified in such licence.
- (3) Every person who injures, defaces, destroys or alters an archaeological object in contravention of this section shall be guilty of an offence under this section, and shall be liable on summary conviction at the suit of the Trust to a fine not exceeding £5,000 or, at the discretion of the court, to imprisonment for any term not exceeding six months or to both such fine and such imprisonment.

23 Restriction of excavation for archaeological purposes

- (1) No person shall without or otherwise than in accordance with a licence issued by the Trust under this section, dig, or excavate, in or under any land (whether with or without removing the surface of the land) for the purpose of searching generally for archaeological objects or of searching for, exposing or examining any particular structure or thing of archaeological interest known to be or believed to be in or under such land or for any other archaeological purpose.

- (2) The Trust may at their discretion issue to any person a licence to dig or excavate in or under any specified land for any specified archaeological purposes and may insert in any such licence such conditions and restrictions as they shall think proper.
- (3) Any person who digs or excavates in or under any land in contravention of this section shall be guilty of an offence under this section and shall be liable on summary conviction at the suit of the Trust to a fine not exceeding £2,500.
- (4) Nothing in this section shall apply to or render unlawful digging or excavation in or under any land for or in the course of any agricultural or industrial operation nor shall a licence under this section operate to render lawful the doing of anything which would be unlawful if this section had not been passed.

Manx National Trust

24 Powers and duties of Manx Museum and National Trust

The Trust shall —

- (a) promote the permanent preservation for the benefit of the people of the Isle of Man of lands and tenements (including buildings) of beauty or historic interest and, as regards lands, the preservation (so far as practicable) of their natural aspect, features and animal and plant life; the preservation of buildings of national interest or architectural, historic or artistic interest, and places of national interest or beauty, and the protection and augmentation of the amenities of such buildings and places and their surroundings;
- (b) preserve all furniture and pictures and chattels of any description having national or historic or artistic interest;
- (c) provide access to, and enjoyment of, such buildings, places and chattels by the public, whenever desirable.

General

25 Power to acquire real estate, etc

Subject to section 25A, the Trust shall have power —

- (a) to acquire land and personal estate,
- (b) to hold land and personal estate;
- (c) to take land on lease or letting;
- (d) to let or lease land not immediately required for the purposes of the Trust;
- (e) with the consent of Tynwald, to sell or exchange land;

- (f) to sell or exchange personal estate;
- (g) with the approval of Tynwald, to raise money by borrowing on the security of the trust property or any part thereof, by way of specific mortgage thereof and by charging or appropriating as security for money borrowed the rents, profits or income derivable from the trust property or any part thereof;
- (h) for the purpose of promoting the interests, powers and duties of the Trust, to act as trustees of any charity or endowment whether such charity or endowment was or is established before or after the passing of this Act and to accept real or personal property given to them as a charity or endowment or upon trust for any purposes in connection with the charity or endowment, and may act as sole trustees of any charity or endowment notwithstanding more than one trustee was either originally appointed or substituted and whether such Trustees were appointed by the court or otherwise;
- (i) the Trustees of any charity may transfer or convey to the Trust any real or personal property comprising the assets or endowment thereof for the purpose of promoting the interests, powers and duties of the Trust and the Trust may accept such transfer or conveyance and shall thereafter be the trustees thereof.²⁹

25A Certain property of Trust to be inalienable

[P1907/CXXXVI/21; P1939/CXXXVI/12]

- (1) If, after the commencement of this section, the Calf of Man shall become vested in the Trust, it shall be held by the Trust for preservation for the benefit of the people of the Isle of Man in accordance with the objects of the Trust, and shall not be chargeable with any debts or liabilities of the Trust and shall be inalienable.
- (2) The Trust may, in respect of any other lands or tenements (including buildings) which are vested in it (whether so vested before or after the commencement of this section), by order determine that such lands or tenements are proper to be held for the benefit of the people of the Isle of Man, and such lands or tenements shall thereupon be so held by the Trust and shall, while the order is in force, be inalienable.
- (3) An order under subsection (2) above shall not come into operation until it has been approved by Tynwald.
- (4) Subsections (1) and (2) above shall not prevent the exercise of the powers contained in section 25(d) of this Act, in relation to any inalienable lands and tenements if Tynwald is satisfied that such exercise is consistent with the purposes of the Trust and has approved the exercise of those powers.

- (5) Notwithstanding subsections (1) and (2) above, the Trust may grant any easement or right (not including a right to the exclusive possession of the surface) over or in respect of any inalienable lands and tenements, and if such grant be by way of lease the same shall be valid and effectual without the approval of Tynwald.
- (6) For the avoidance of doubt section 28 of the *Interpretation Act 1976* (power to amend or revoke public documents) shall apply to subsection (2) of this section.
- (7) This section shall not prejudice the operation of any enactment relating to the compulsory acquisition of land.³⁰

25B Power to enter into agreements restricting use of land

[P1937/LVII/8]

- (1) Where any person is willing to agree with the Trust that any land or any part thereof shall, so far as his interest in the land enables him to bind it, be made subject either permanently or for a specified period to conditions restricting the planning development or use thereof in any manner, the Trust may, if it thinks fit, enter into an agreement with him or accept a covenant from him to that effect.
- (2) The Trust shall have power to enforce such an agreement or covenant against persons deriving title under him in the like manner and to the like extent as if the Trust were possessed of, or entitled to, or interested in, adjacent land and as if the agreement or covenant had been and had been expressed to be entered into for the benefit of that adjacent land.³¹

26 Powers exercisable over Trust

The Trust with respect to the Trust property —

- (a) may plant, drain, level, and otherwise improve and alter, any part or parts of such property, so far as they deem necessary or desirable, and they may make enclosures for these purposes and for the purpose of protecting or renovating turf and for protecting trees and plantations;
- (b) may make and maintain roads, footpaths, and ways over such property, and may make and maintain ornamental ponds and waters on such property;
- (c) may, on such property, erect any necessary buildings, and may maintain and repair such buildings;
- (d) shall, by all lawful means, prevent, resist and abate all enclosures and encroachments upon, and all attempts to enclose or encroach upon such property or any part thereof, or to appropriate or use the same, or the soil, timber, or roads thereof, or any part thereof for any purpose inconsistent with this Act;

- (e) may set apart, from time to time, parts of such property upon which persons may play games or hold meetings or gatherings for athletic sports or may land from boats;
- (f) may make such reasonable charges for the admission of the public to such property, or any part thereof, or for the use by the public of any such property or any part thereof, as they may from time to time determine.

26A Trustees' duty to consult on heritage matters

In performing the Trust's functions relating to the Island's heritage, the trustees must consult with the persons or organisations they consider appropriate.³²

27 Power to make, etc, bye-laws

- (1) The Trust may from time to time make, revoke and alter bye-laws for the regulation and protection of the trust property, for the prevention of nuisances upon the trust property, and for the preservation of order upon the trust property.
- (2) Any such bye-law may impose penalties for any breach thereof.
- (3) Any such penalties may be recovered, on summary conviction, at the suit of the Trust, or any officer of Police, and shall be added to the General Revenue of the Trust.
- (4) Any bye-laws made, revoked or altered by the Trust under this section shall be subject to the approval of Tynwald.

28 Trust to be exempt from Income Tax

The income derived from any property of the Trust shall be exempt from the provisions of the Income Tax Acts, 1946 and 1958, and any Act amending the same.

29 Ministerial meetings with stakeholders

- (1) The Minister may convene and chair meetings ("stakeholder meetings") with any person whom the Minister considers to be a stakeholder of the Trust in order to discuss any matter relating to the Island's heritage.
- (2) The Minister may participate in the discussion.
- (3) Before convening a stakeholder meeting the Minister may consult the trustees as to —
 - (a) the conduct and form of the meeting; or
 - (b) the persons he or she proposes to invite to attend.
- (4) The Minister may require all or any of the trustees and the director to attend a stakeholder meeting.

- (5) In this section, “director” means the Trust’s director or, if the Trust has no office of that title, its most senior officer from time to time.”.

30 Crown rights

Nothing in this Act affects prejudicially any estate, right, power, privilege or exemption of the Crown, and in particular nothing herein contained authorises the Trust to take, use, or in any manner interfere with, any land, or hereditaments or any rights of whatsoever description belonging to Her Majesty in right of Her Crown, or vested in the Government Property Trustees for the public service of this Isle, without the consent in writing of those Commissioners on behalf of Her Majesty, or the Government Property Trustees, as the case may be, first had and obtained for that purpose.

31 Commencement of Act

This Act shall come into operation when the Royal Assent thereto has been by the Governor announced to Tynwald and a Certificate thereof has been signed by the Governor and the Speaker of the House of Keys.

*Transitional provisions for Manx Museum and
National Trust (Amendment) Act 2011³³*

32 Definitions for ss 32 to 35

In sections 32 to 35 —

“amending Act” means the Manx Museum and National Trust (Amendment) Act 2011;

“commencement” means the day on which Royal Assent to the amending Act is announced as mentioned in section 10(1) of the Interpretation Act 1976;

“existing office” means the offices, under old section 3, of a member of the Trust or of the Chairman or Vice-Chairman of the Trust;

“new appointment provisions” means sections 3B and 3D to 3G;

“old section 3” means section 3 as in operation before the commencement.

33 Appointment of new chairperson and other trustees

- (1) During the period mentioned in subsection (2) —

- (a) the Minister must appoint the Minister’s appointee; and
- (b) the Council of Ministers must appoint at least 9 Council of Ministers appointees, one of whom must be appointed as chairperson.

- (2) For subsection (1), the period is —

- (a) if the Keys is dissolved after the passage, by both the Keys and the Council, of the Bill for the amending Act but before the commencement the period starting on the commencement and ending on the first anniversary of the commencement; or
 - (b) if the Keys is not dissolved during the period mentioned in paragraph (a) the period starting on the first dissolution of the Keys after the commencement and ending on the first anniversary of the dissolution.
- (3) The new appointment provisions apply to the appointments, despite there being existing offices that are still occupied.
- (4) However, the appointments must all take effect on the same day.

34 Existing trustees continue until appointments take effect

- (1) Old section 3 continues to apply to the appointment of a person to, and to the holding of, an existing office until the start of the day (the “changeover time”) on which the appointments under section 33 take effect.
- (2) Each holder of an existing office continues in that office until the changeover time.
- (3) Subsections (1) and (2) apply despite the new appointment provisions and section 3 (tenure of office of members of Statutory Boards) of the Statutory Boards Act 1987, as applied under old section 3(2)(e).
- (4) At the changeover time, each existing office holder goes out of that office.
- (5) Subsection (4) does not prevent an existing office holder from holding an appointment mentioned in section 33(1).

35 Expiry of transitional provisions

Sections 32, 33 and 34, this section and the crossheading immediately before section 32 expire on the second anniversary of the commencement.

FIRST SCHEDULE³⁴**SECOND SCHEDULE**

Section 16

PART I*Provisions as to confirmation, variation and revocation of preservation schemes*

1. (1) Before confirming a preservation scheme (hereafter in this Schedule referred to as a 'scheme') the Trust shall cause to be published in two newspapers printed and circulating in this Isle, and in such other manner as they think best for informing persons affected, notice of their intention to confirm the scheme, of the place where copies thereof may be inspected, and of the time (which shall not be less than three months) within which and the manner in which representations with respect to the scheme may be made, and shall cause such notice as aforesaid to be given to every local authority whose area comprises any part of the controlled area.
2. Any person affected by the scheme may, within the time appointed under the last foregoing paragraph for making representations, send to the Trust written objection to the scheme stating the specific grounds of objection and the specific modifications required.
3. The Trust after considering any representations and objections duly made with respect to a scheme, may by order confirm the scheme either with or without modifications:
Provided that —
 - (a) where an objection has been duly made to the scheme by any person appearing to the Trust to be affected thereby and has not been withdrawn, the Trust unless they consider the objection to be frivolous or have modified the scheme as required by the objection, shall, before confirming the scheme, direct a public inquiry to be held as hereinafter provided and consider the report of the person who held the inquiry; and
 - (b) a scheme as so confirmed shall not apply to any area to which it would not have applied if it had been confirmed without modification.
4. Any inquiry under this schedule shall be held in accordance with rules made by the Trust for the purpose, and such rules may contain provisions as to the costs of the inquiry.
5. A scheme when so confirmed shall come into force on such date as may be specified in the order confirming it.

6. A scheme may be varied or revoked by a subsequent scheme and the Trust may by order revoke a scheme if they think in the circumstances that the scheme ought to be revoked.

7. As soon as practicable after the making of an order under this schedule confirming or revoking a scheme the order shall be published in two newspapers printed and circulating in this Isle and in such other manner as the Trust think best for informing persons affected, and a copy of the order shall be sent to every local authority whose area comprises any part of the controlled area.

PART II

Provisions as to compensation

1. No person shall be entitled to compensation in respect of a preservation scheme unless within three months from the date on which the scheme comes into force, or within such further time as the Trust may in special circumstances allow, he makes a claim for the purpose in such manner as the Trust may by regulations prescribe.

2. A person shall not be entitled to compensation by reason of the fact that any act or thing done or caused to be done by him has been rendered abortive by a preservation scheme, if or so far as the act or thing was done after the date on which the Trust published in two newspapers printed and circulating in this Isle notice of their intention to confirm the scheme, or by reason of the fact that the performance of any contract made by him after that date is prohibited by the scheme.

3. Where any provision of a preservation scheme was, immediately before the scheme came into force, already in force by virtue of this or any other Act, no compensation shall be payable by reason of any property being injuriously affected by that provision of the preservation scheme if compensation has been paid, or could have been claimed, or was not payable, by reason of that property having been injuriously affected by the provision already in force.

4. Where any provision of a preservation scheme could, immediately before the scheme came into force, have been validly included in a scheme, order, regulation or bye-law by virtue of any other Act, then —

- (a) if no compensation would have been payable by reason of the inclusion of that provision in the scheme, order, regulation or bye-law, no compensation shall be payable in respect of that provision of the preservation scheme: and
- (b) if compensation would have been so payable, the compensation payable in respect of that provision of the preservation scheme shall not be greater than the compensation which would have been so payable.

5. Any dispute as to whether any property is injuriously affected by a preservation scheme, or as to the amount of the sum which is to be paid as compensation in respect of such a scheme, shall be determined by arbitration under and in accordance with the Arbitration Acts, 1910 to 1935.

THIRD SCHEDULE³⁵

ENDNOTES

Table of Legislation History

Legislation	Year and No	Commencement

Table of Renumbered Provisions

Original	Current

Table of Endnote References

¹ Long title repealed by Manx Museum and National Trust (Amendment) Act 2011 s.3.

² Section 1A inserted by Manx Museum and National Trust (Amendment) Act 2011 s.4

³ Definition of “appointment recommendation” inserted by Manx Museum and National Trust (Amendment) Act 2011 section 5(1).

⁴ Definition of ‘archeological object’ substituted by Manx Museum Act 1984 s 1.

⁵ Definition of “chairperson” inserted by Manx Museum and National Trust (Amendment) Act 2011 s. 5(1).

⁶ Definition of “Council of Ministers appointee” inserted by Manx Museum and National Trust (Amendment) Act 2011 s.5(1).

⁷ Definition of “Department” inserted by inserted by Manx Museum and National Trust (Amendment) Act 2011 s.5(1).

⁸ Definition of “member” inserted by inserted by Manx Museum and National Trust (Amendment) Act 2011 s.5(1).

⁹ Definition of “Minister” inserted by inserted by Manx Museum and National Trust (Amendment) Act 2011 s.5(1).

¹⁰ Definition of “Minister’s appointee” inserted by inserted by Manx Museum and National Trust (Amendment) Act 2011 s.5(1).

¹¹ Definition of ‘Museum’ amended by Statute Law Revision Act 1997 Sch 1.

¹² Para (c) amended by Statute Law Revision Act 1997 Sch1.

¹³ Para (e) amended by Statute Law Revision Act 1997 Sch 1.

¹⁴ Definition of “trustee” inserted by Manx Museum and National Trust (Amendment) Act 2011 s.5(1).

¹⁵ Subsection (3) added by by Manx Museum and National Trust (Amendment) Act 2011 s.5(2).

¹⁶ Sections 3 to 3E substituted by Manx Museum and National Trust (Amendment) Act 2011 s.6.

¹⁷ S 4 repealed by Statute Law Revision Act 1989 Sch 2.

¹⁸ Subs (4A) inserted by Public Records Act 1999 Sch 3.

¹⁹ Subs (6) added by Manx Museum Act 1984 s 3 and amended by Public Records Act 1999 Sch 3.

²⁰ Subs (7) repealed by Public Records Act 1999 Sch 4.

²¹ S 7 amended by Governor's General Functions (Transfer) Act 1980 Sch 1.

²² Para (b) inserted by Statute Law Revision Act 1992 Sch 1.

²³ Subs (4) amended by Statute Law Revision Act 1992 Sch 1.

²⁴ Subs (6) amended by Manx Museum Act 1984 Sch.

²⁵ Subs (8) repealed by Statute Law Revision Act 1983 Sch 2.

²⁶ S 21 substituted by Manx Museum Act 1984 s 1.

²⁷ References to Chairman and Vice-chairman superseded by Manx Museum and National Trust (Amendment) Act 2011 s. 7.

²⁸ S 21 inserted by Manx Museum Act 1984 s 2.

²⁹ S 25 amended by Manx Museum and National Trust (Amendment) Act 1986 s 3.

³⁰ S 25A inserted by Manx Museum and National Trust (Amendment) Act 1986 s 3.

³¹ S 25B inserted by Manx Museum and National Trust (Amendment) Act 1986 s 2.

³² Section 26A inserted by Manx Museum and National Trust (Amendment) Act 2011 s.8.

³³ Sections 32 to 35 and the cross-heading preceding them inserted by Manx Museum and National Trust (Amendment) Act 2011 s 10 until . 18/10/2013. The version of section 3 referred to in these sections read as follows—

3 Constitution of Trust

- (1) The Trust shall continue to consist of the following members-
 - (a) [Para (a) repealed by Manx Museum Act 1984 s 5.]
 - (b) one member of the Legislative Council, to be elected by the Legislative Council; [Para (b) substituted by Manx Museum and National Trust (Amendment) Act 1986 s 1.]
 - (c) the Speaker of the House of Keys;
 - (d) one member of the House of Keys, to be elected by the House of Keys; [Para (d) substituted by Manx Museum and National Trust (Amendment) Act 1986 s 1.]
 - (e) one person to be appointed by the Council of the National Trust constituted by the National Trust Acts of the Imperial Parliament;
 - (f) two persons to be from time to time appointed by the Mayor, Aldermen and Burgesses of the Borough of Douglas;
 - (g) one person to be from time to time appointed by each of the following authorities, that is to say: Ramsey Town

-
- Commissioners, Castletown Town Commissioners, Peel Town Commissioners;
- (h) one person to be from time to time appointed by the University of Liverpool;
 - (i) one person to be from time to time appointed by the Isle of Man Education Authority;
 - (j) [Para (j) repealed by Statute Law Revision Act 1997 Sch 2.]
 - (k) one person to be appointed by each of the five societies or institutions selected from time to time by the Trust as representatives of societies or institutions engaged in scientific, artistic, historical or kindred pursuits within this Isle;
 - (l) seven persons to be co-opted by the Trust.[Para (l) amended by Manx Museum and National Trust (Amendment) Act 1986 s 1.]
- (2) With respect to the tenure of office of the members of the Trust, the following provisions shall take effect-
- (a) to (c) [Paras (a) to (c) repealed by Members of Boards (Miscellaneous Provisions) Act 1976 Sch 2.]
 - (d) if any member of the Trust appointed or elected by any of the various bodies referred to in subsection (1) of this section ceases to be a member of that body or (except in the case of representatives of the University of Liverpool and the Council of the National Trust) resides out of the Isle of Man for a continuous period of twelve months, he shall, ipso facto, cease to be a member of the Trust;
 - (e) section 3 of the Statutory Boards Act 1987 applies to the Trust as it applies to a Statutory Board. [Para (e) substituted by Council of Ministers Act 1990 Sch 1.]
 - (f) [Para (f) repealed by Statutory Boards Act 1987 Sch 4.]
- (3) With respect to the Trust the following provisions shall take effect —
- (i) the Trust shall continue to be a body corporate by the name of the Manx Museum and National Trust with perpetual succession and a common seal;
 - (ii) the Trust shall elect annually by ballot two of their number to be Chairman and Vice-Chairman respectively and shall also so proceed on any casual vacancy occurring in the office of Chairman or Vice-Chairman by resignation of such office or from any other cause;
 - (iii) Schedule 2 to the Statutory Boards Act 1987 (except paragraphs 1, 6 and 7) applies to the Trust as it applies to a Statutory Board.

³⁴ Sch 1 repealed by Statute Law Revision Act 1983 Sch 2.

³⁵ Sch 3 repealed by Statute Law Revision Act 1989 Sch 2.

Maughold Cliffs & Brooghs Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im




**Isle of Man
Government**

Reiltys Ellan Vannin

Maughold Cliffs & Brooghs ASSI

2009 aerial




Note: The seaward boundary follows the lowest astronomical tide between points A and B on the map.

Area: 53.63 hectares (132.52 acres)

National Grid Reference (centroid) : SC 487923

Designated: 30th March 2011
 Confirmed: 5th September 2011

Maughold Cliffs & Brooghs ASSI




Legend
 Maughold Cliffs & Brooghs ASSI

Note: The seaward boundary follows the lowest astronomical tide between points A and B on the map.

Area: 53.63 hectares (132.52 acres)

National Grid Reference (centroid) : SC 487923

Designated: 30th March 2011
 Confirmed: 5th September 2011



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Maughold Cliffs & Brooghs

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Maughold

Local authority: Maughold Commissioners

Planning Authority: Department of Infrastructure

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheet SC49SE

National Grid Reference (centroid): SC 487 923 **Area:** 53.63 hectares (132.52 acres)

Date notified: 30th March 2011

Date confirmed: 5th September 2011

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its breeding avifauna and semi-natural coastal habitats; hard cliffs and slopes, coastal grassland, acid grassland and scrub.

Description and reasons for notification:

Maughold cliffs and brooghs ASSI boundary encompasses a wide-range of semi-natural habitats grading from species-rich coastal grassland to hard cliff and intertidal rocky shore. It provides excellent breeding and feeding territory for seabirds and other coastal species, which are sufficient to merit site selection on the grounds of having a good range of representative breeding birds.

A total of 35 definite or probable breeding bird species has been recorded since 1999; there are also recent records of a further 47 species of possible breeders, passage or wintering birds, of which 5 are listed on Schedule 1 of the Wildlife Act 1990.

Of most interest is the colony of cliff-breeding sea birds which includes a colony of shags *Phalacrocorax aristotelis** and the largest colony of cormorants *Phalacrocorax carbo* on the Isle of Man. Other breeding seabirds include puffin *Fratercula arctica*, fulmar *Fulmarus glacialis*, kittiwake *Rissa tridactyla*, black guillemot *Cepphus grylle*, guillemot *Uria aalge* and razorbill *Alca torda*. Other important cliff breeders include raven *Corvus corax*, peregrine falcon *Falco peregrinus** and chough *Pyrhocorax pyrrhocorax**. In addition to providing valuable nesting habitat the cliffs support patches of diverse coastal grassland, cliff crevice plants and scrub.

The whole site is notable for its diversity and extent of typical coastal habitats. The steep rocky cliffs and coastal brooghs that stretch from Port-e-Vullen to an area south of the lighthouse, stopping short of Port Mooar are one of the largest continuous stretches of tall cliff and unimproved coastal habitat in the Isle of Man. The coast is generally composed of steep cliffs with *Fucus* covered rocks below and gently sloping grassy slopes above. A number of small flushes and streams occur in the grassland and add to the botanical-interest. One of these forms St Maughold's Well which is of cultural and historical interest. Bracken clothes the slopes in summer and small patches of scrub add to the habitat diversity. Heath-spotted orchid *Dactylorhiza maculata*** occur in large numbers (c.200) in a number of isolated patches. Grey seals*** haul out and breed on the beaches and rocks at the base of the cliffs.

Viviparous lizards *Lacerta vivipara**** have been recorded on the coastal broughs in the vicinity of the footpath at the southern end of the ASSI. It is likely that they occur on other south-facing areas of the site.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

***= Protected under Schedule 5 of the Wildlife Act 1990

Other information

Geology and archaeology

The coastal cliffs expose Manx Group rocks of Ordovician age (about 482 million years old). Within this group, the Maughold Formation extends from Port e Vullen eastwards to Stack ny Skey, where it meets the Creg Agneash Formation. The Maughold Formation consists of typically "pin-stripe" lamination of thin bands of alternating mudstone and sandstone with some siltstone bedded between layers, whereas the Creg Agneash Formation is paler and more dominated by layers of sandstone. As with other rocks of this age, the exposures show the effects of tectonic forces in the earth's crust by much folding, distortion and the presence of faults.

Minerals were mined here in the past, particularly hematite or iron ore. Evidence of 6th – 7th century working of Maughold iron ore has been found by analysis of a smelting site near Andreas. Iron ore was produced in some quantity during the 19th century, the hey-day of mining in the Isle of Man, when annual production could reach 2,000 tons. The most visible remains of this now defunct industry is several adit entrances along the coast near the foot of the cliffs, and there is an old mine shaft at Maughold Head.

At the summit of Maughold Head near the site of the iron ore exposure is an earthwork enclosure or fortification, probably dating to the Iron Age. The summit of the Head later also served as a gathering point for medieval militia undertaking coastal lookout ('Watch and Ward') duties. A stone burial cist just south of the Head is probably of Bronze Age date and a cairn on the summit may be of similar age. On the north slope of the Head lies a spring which feeds a well, named after St Maughold and associated with an early medieval monastic settlement which stood nearby under the present churchyard. A shelter in the churchyard displays cross-slabs from the early Christian period which have been gathered from the churchyard and wider afield within the surrounding parish.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Maughold Cliffs & Brooghs

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseeded
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weed killers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control.
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17	Reclamation of land from sea, estuary or marsh.
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.

20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest.
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora.
28	Game and waterfowl management and hunting practices, where already damaging, introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, reptile, amphibian, bird, fish or invertebrate

Poyll Vaaish Coast Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Re: llys Ellan Vannin



**Isle of Man
Government**
Reiltys Eilan Varrin

Department of Agriculture,
Fisheries and Forestry

**POOYL VAAISH COAST
AREA OF SPECIAL
SCIENTIFIC INTEREST**

Designation boundary



Area: 44.76ha (= 110.6 acres)
O/S Grid reference: SC242678
(approximate centre of site)

Designated: 18th February 2007
Varied and confirmed: 9th August 2007


*Note: the seaward boundary of the site is
defined as the Lowest Astronomical Tide
between points A and B on this map.

The grid lines on this map form
part of the National Grid,
and are spaced at 500m intervals

Base Map is

Crown Copyright

Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Poyll Vaaish Coast

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parishes: Arbory and Malew

Planning Authority: Department of Local Government and the Environment

Local authority: Arbory Commissioners, Malew Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC 26NW

National Grid Reference (centroid): SC242678 **Area:** 44.76ha (110.6 acres) –
based on position of Lowest
Astronomical Tide boundary – see Map

Date notified: 19th February 2007

Date confirmed: 9th August 2007

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its breeding and visiting birds and its intertidal, saltmarsh, vegetated strandline, vegetated shingle and coastal grassland habitats.

Description and reasons for notification:

Poyll Vaaish Coast ASSI consists of low, rocky, carboniferous limestone coastline with coarse shingle, saltmarsh and a series of small bays and pools. The aspect is predominantly south-facing, and this factor, coupled with the varied nature of the terrain, has led to ideal conditions for a mix of sun- and shelter-loving plants and invertebrates alongside species which can tolerate very exposed coastal conditions. The combination of limestone geology and varied terrain make this an important and irreplaceable location for Manx coastal biodiversity.

The area has a very varied patchwork of vegetation communities including vegetated shingle, vegetated strandline, saltmarsh and coastal grassland. Overall the site is rich in plant biodiversity, as it not only contains many coastal terrestrial plants but also a good range of seaweeds in its intertidal zone. The saltmarsh is ungrazed, and varies from lower saltmarsh with muddy ground washed by the tide, to middle and upper saltmarsh which merges with the adjacent habitats of vegetated shingle and coastal grassland. Upper saltmarsh is usually limited to scattered patches, whereas the lower, wetter vegetation forms one of the largest continuous patches of saltmarsh on the Island. Species present include frequent glasswort *Salicornia* spp., with red fescue *Festuca rubra* and common saltmarsh grass *Puccinellia maritima*. Notable plants include sea purslane *Atriplex portulacoides***, a rarity on the Island which here forms patches of shrubby saltmarsh vegetation with other saltmarsh plants such as sea aster *Aster tripolium*, sea plantain *Plantago maritimum*, annual sea-blite *Suaeda maritima*, greater sea-spurrey *Spergularia media*, lesser sea-spurrey *Spergularia marina*, and spear-leaved orache *Atriplex prostrata*.

The soft ground and tidal pools of the saltmarsh attract an important assemblage of breeding and wintering coastal birds. These include chough *Pyrrhocorax pyrrhocorax**, breeding eider *Somateria mollissima**, breeding Ringed plover *Charadrius hiaticula**, and breeding Shelduck *Tadorna tadorna**. Birds attracted to the site are not limited to typically coastal species; in total, 95 species have been recorded feeding and/or breeding on the saltmarsh, shingle, rocks and grassland in the last five years, including songbirds, waders, wildfowl and birds of prey. Of these, 35 are protected under Schedule 1 of the Wildlife Act 1990, making this an important site for rare and vulnerable birds on the Island.

Other notable species known to breed on and/or visit Poyll Vaaish include Skylark *Alauda arvensis**, Black redstart *Phoenicurus ochruros**, Reed bunting *Emberiza schoeniclus**, Song thrush *Turdus philomelos**, Twite *Carduelis flavirostris**, Curlew *Numenius arquata**, Lapwing *Vanellus vanellus**, Golden plover *Pluvialis apricaria**, Black-tailed godwit *Limosa limosa*, Teal *Anas crecca*, Shoveler *Anas clypeata*, Whooper swan *Cygnus cygnus**, Hen harrier *Circus cyaneus**, Kestrel *Falco tinnunculus**

Long-eared owl *Asio otus** and Barn owl *Tyto alba**. Eight of the species recorded on the area are on the RSPB Red List of Birds of Conservation Concern, and 47 of the remaining species are on the Amber List, hence this is an important site for birds which are threatened in the wider geographical context.

Whilst the seaward side of Poyll Vaaish Coast grades into a species-rich inter-tidal zone, the upper edge grades from vegetated shingle into coastal grassland. The vegetated shingle is exceptionally diverse, with a colourful array of ruderal and coastal plants such as sea mayweed *Tripleurospermum maritimum*, sea kale *Crambe maritima*, smooth sowthistle *Sonchus oleraceus*, sea spurge *Euphorbia paralias*, common poppy *Papaver rhoeas*, rock samphire *Crithmum maritimum*, sea sandwort *Honckenya peploides*, common scurvy-grass *Cochlearia officinalis*, sea beet *Beta vulgaris* ssp. *maritima*, sea radish *Raphanus raphanistrum* ssp. *maritimus*, low-growing blackthorn *Prunus spinosa*, pellitory-of-the-wall *Parietaria judaica*, creeping cinquefoil *Potentilla reptans*, curled dock *Rumex crispus*, common orache *Atriplex patula* and Babington's orache *Atriplex glabriuscula*. Many of these plants produce abundant seed and attract insects, both of which provide valuable food for birds, including flocks of linnet *Carduelis cannabina*, starlings *Sternus vulgaris** and house sparrows *Passer domesticus**.

The upper edge of the site abuts the coastal path, and has a mixture of weedy plants on shingle, and established coastal grassland. Species which are typical of unimproved meadow grassland occur here, such as lady's bedstraw *Galium verum*, lesser hawkbit *Leontodon saxatilis*, common cat's-ear *Hypochoeris radicata*, common bird's-foot trefoil *Lotus corniculatus*, wild carrot *Daucus carota*, dandelion *Taraxacum agg.*, red fescue *Festuca rubra*, common milkwort *Polygala vulgaris*, cock's-foot *Dactylis glomerata*, common ragwort (cushag) *Senecio jacobaea*, common knapweed *Centaurea nigra*, field scabious *Knautia arvensis*, wild thyme *Thymus polytrichus*, and yarrow *Achillea millefolium*, in conjunction with salt-tolerant plants including buckshorn plantain *Plantago coronopus*, thrift *Armeria maritima*, rock sea-spurrey *Spergularia rupicola*, spring squill *Scilla verna*, sea mouseear *Cerastium diffusum*, and sea campion *Silene uniflora*. Unimproved grassland of this level of diversity is a rare and vulnerable habitat on the Isle of Man, hence this site is of importance for the conservation of these species, which form a distinctive element of the Manx landscape and support a wide range of associated birds and invertebrates.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

Geology

Poyll Vaaish Coast is part of the well-known and significant Carboniferous sequence between Langness and Port St Mary. The rocks here are rich in well-preserved fossils.

Rocks here form part of the important sequence of Carboniferous age features which are exposed along the length of the coast between Langness and Port St Mary, and include the disruptive marine volcanic structures of Scarlett and the basal conglomerate within Langness ASSI. Within the Poyllvaaish Coast ASSI boundary the strata are mostly fine-grained limestones of the Balladoole Formation to the north west of Poyllvaaish Farm and, around Poyll Ritchie, fine-grained, irregularly bedded carbonate mudstones of the Knockrushen Formation. Both formations are rich in fossils, such as crinoids, brachiopods, corals and ammonoids and, because fossiliferous rocks are so restricted in the Isle of Man, have considerable educational value.

Boundary

The boundary of the site has been set at Lowest Astronomical Tide.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Poyll Vaaish Coast

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17	Reclamation of land from sea, estuary or marsh.
18	Bait digging on inter-tidal areas.
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.

23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora (excluding walking, with or without dogs).
28	Introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* “animal” includes any mammal, reptile, bird, fish or invertebrate.

Port St Mary Ledges & Kallow Point Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**

Reiltys Ellan Vannin


Port St Mary Ledges & Kallow Point ASSI

Designated: 1st November 2010

Varied and confirmed: 28th April 2011

Area: 14.79 hectares (= 36.55 acres)

National Grid Reference (centroid): SC 211 669




Port St Mary Ledges & Kallow Point ASSI

Designated: 1st November 2010

Varied and confirmed: 28th April 2011

Area: 14.79 hectares (= 36.55 acres)

National Grid Reference (centroid): SC 211 669



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Port St Mary Ledges & Kallow Point

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Rushen

Planning Authority: Department of Infrastructure

Local authority: Port St Mary Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC 26NW

National Grid Reference (centroid): SC211669 **Area:** 14.79ha (36.55 acres)

– based on position of Lowest Astronomical Tide boundary – see Map

Date notified: 1st November 2010

Date confirmed: 28th April 2011

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its matrix of maritime habitats, principally its intertidal limestone ledges featuring distinctive zonation from the eulittoral zone to the sublittoral zone, vegetated shingle, hard cliff and coastal grassland habitats. The site also hosts a number of important breeding and visiting bird species.

Description and reasons for notification:

Port St Mary Ledges & Kallow Point ASSI consists predominantly of low, rocky, carboniferous limestone coastline with an associated patchwork of vegetation communities comprising vegetated shingle, hard cliff and coastal grassland. The aspect is predominantly south-facing, and this factor, coupled with the varied nature of the terrain, has led to ideal conditions for a mix of sun and shelter-loving plants and invertebrates alongside species which can tolerate very exposed coastal conditions. The combination of limestone geology and varied terrain make this an important and irreplaceable location for Manx coastal biodiversity, a fact that has been acknowledged by generations of marine biologists from the Port Erin Marine Laboratory who have conducted research on the site for over 70 years. The site is internationally recognised by the scientific community and research on the site has been published in nearly 100 peer-reviewed scientific publications.

Bird species known to breed on Port St Mary ledges and Kallow Point include herring gull *Larus argentatus*, oystercatcher *Haematopus ostralegus*, rock pipit *Anthus petrosus* and ringed plover *Charadrius hiaticula**. In addition there are many birds that use the site for feeding and/or roosting, they include black guillemot *Ceppus grylle*, chough *Pyrrhocorax pyrrhocorax**, cormorant *Phalacrocorax carbo*, dunlin *Calidris alpina*, eider *Somateria mollissima*, fulmar *Fulmarus glacialis*, house sparrow *Passer domesticus**, knot *Calidris canuta*, merlin *Falco columbarius*, purple sandpiper *Calidris maritima*, redshank *Tringa tetanus*, shag *Phalacrocorax aristotelis**, shelduck *Tadorna tadorna*, snipe *Gallinago gallinago*, song thrush *Turdus philomelos**, starling *Sturnus vulgaris**, stonechat *Saxicola torquata*, turnstone *Arenaria interpres*, wheatear *Oenanthe oenanthe* and whimbrel *Numenius phaeopus*.

Five of the species recorded on the area are on the RSPB Red List of Birds of Conservation Concern and 34 of the remaining species are on the Amber List, hence this is an important site for birds which are threatened in the wider geographical context.

The Port St Mary ledges are categorised as Moderately Exposed Rock with a wide variety of associated biotope categories. To the West of the site, there is an area of shingle and a promenade wall which has unvegetated sections and other areas densely covered by the lichen *Verrucaria maura*.

Drainage holes in the wall input freshwater into this section and in those areas the green algae *Enteromorpha intestinalis* is present. There is an extensive area of shingle approaching the landward end of the outer harbour and further areas of *Enteromorpha* corresponding with other freshwater inputs. The habitat on the seaward side of the outer harbour is dominated by dense serrated wrack *Fucus serratus*.

The ledge formation provides a wide range of intertidal habitats including open expanses of freely draining rock, pock-marked areas retaining water, boulder fields and variety of rock pools and deep varied crevices.

On the mid-shore to the west of the site the dominant species is bladder wrack *Fucus vesiculosus* with limpets *Patella vulgata*, barnacles *Semibalanus balanoides* and abundant grazing snails (littorinids). There are also a number of pools characterised by the coralline algae *Corallina officinalis*, with a number of other species of green, brown and red seaweeds, littorinids and the top shell *Gibbula cineraria*. There are also some small patches of the red seaweed *Osmundea pinnatifida* and *Gelidium pusillum*, a scarce biotope in the Isle of Man and also in the UK. Towards the East of the site the shore is dominated by a series of ledges of around 0.5m high, vegetated mainly by serrated wrack *Fucus serratus* with other algae species and barnacles.

The lower shore ledges are dominated by the brown algae thongweed *Himanthalia elongata* and various red algae species with a carpet of the coralline algae *Corallina officinalis* and encrusting coralline algae *Lithothamnion*. To the west of the site there is a small patch of dulse (*Palmaria palmata*) on very exposed rock, an uncommon biotope in the Isle of Man and the UK. At the bottom of the shore and moving into the sublittoral zone a wide zone of kelp *Laminaria digitata* dominates with diverse fauna including the painted top shell *Calliostoma zizyphinum*. The crevices in the limestone ledges are highly diverse and are home to a wide variety of sponges species.

The site is recognised as one of the best rocky shores in the British Isles for marine ecological studies. It has been extensively studied by scientists over the past century and it has also been selected as a monitoring site for a British Isles-wide project assessing the impact of climate change on coastal habitats. One of the reasons this site was chosen for this important study was the good historical baseline of the site. Internationally renowned scientists are using the site to detect changes in species composition as a result of climate change, assisting the Isle of Man in understanding and adapting to the effects of climate change. For example, in 2004 Kallow Point was the first Manx site where the southern barnacle *Chthamalus stellatus* was recorded. Kallow Point is also a good site to monitor the loss of cooler water species, for example the tortoiseshell limpet, *Tectura testudinalis*.

Port St Mary ledges are one of only two intertidal limestone ledges still remaining intact on the Isle of Man, following the loss of a significant proportion of the Island's most diverse intertidal site as a result of the construction of the Ronaldsway runway extension. Limestone ledges form a very small proportion of the Island's total intertidal area and are thus of particular interest. The intertidal communities associated with limestone ledges are also significantly different from those associated with the predominant Manx slate shores. The other remaining limestone ledges site is at Scarlett. This site is much more diverse than Port St Mary ledges but is subject to less wave action and therefore significantly different in terms of intertidal communities.

The upper (northerly) edge of the site comprises terrestrial habitats that grade from vegetated shingle to hard cliff and coastal grassland, predominantly on the western end of the site. The vegetated shingle is not particularly diverse, with a colourful array of ruderal and coastal plants such as sea mayweed *Tripleurospermum maritimum*, smooth sowthistle *Sonchus oleraceus*, cleavers *Galium aparine*, curled dock *Rumex crispus*, common scurvy-grass *Cochlearia officinalis*. Low-growing blackthorn *Prunus spinosa* and elder *Sambucus nigra* grow on the most stabilised and landward shingle. Many of these plants produce abundant seed and attract insects, both of which provide valuable food for birds.

The hard cliffs are sparsely vegetated with common scurvy-grass *Cochlearia officinalis*, rock sea-spurrey *Spergularia rupicola* and sea spleenwort *Asplenium marinum*. The coastal grassland is

particularly species-rich, being dominated by red fescue *Festuca rubra* and containing vernal species such as spring squill *Scilla verna*, lesser celandine *Ranunculus ficaria* and bluebell *Hyacinthoides non-scripta*. Many meadow species are common, such as lady's bedstraw *Galium verum*, hawkbit *Leontodon spp*, common cat's-ear *Hypochaeris radicata*, common bird's-foot trefoil *Lotus corniculatus*, wild carrot *Daucus carota*, dandelion *Taraxacum agg.*, glaucous sedge *Carex flacca*, cock's-foot *Dactylis glomerata*, common ragwort (cushag) *Senecio jacobaea*, common knapweed *Centaurea nigra*, bulbous buttercup *Ranunculus bulbosus*, wild thyme *Thymus polytrichus* and yarrow *Achillea millefolium*, in conjunction with salt-tolerant plants including buck's-horn plantain *Plantago coronopus*, thrift *Armeria maritima* and sea campion *Silene uniflora*. Unimproved grassland of this level of diversity is a rare and vulnerable habitat on the Isle of Man, hence this site is of importance for the conservation of these species, which form a distinctive element of the Manx landscape and support a wide range of associated birds and invertebrates.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

Geology

The rocks at Kallow Point present useful exposures of the Carboniferous Limestone which in the Isle of Man is limited to the area around Castletown and along the coast between Langness and Port St Mary.

At Kallow Point itself, near horizontal beds of the Knock Rushen Formation consist of thick, fine-grained carbonate mud, containing fossils of corals, brachiopods, bryozoa and crinoids. The fossilised remains of animal burrows contribute to evidence that this was once the floor of a tropical sea at a time (350 to 300 million years ago) when the deposits that now make up the Isle of Man were situated near the equator. Ancient ripple marks and overturned brachiopods indicate deposition in a lagoon.

The Knock Rushen Formation is faulted against Manx Group rocks (Ordovician; Manx Group) on both sides of the Kallow Point peninsula and is thus an outlier of Carboniferous rock on the western side of Bay Ny Carrickey. The Formation re-appears on the other side of the Bay slightly to the east of Kentraugh.

An additional feature of interest is the hummocky surface of the limestone below the sea wall, indicating the beginnings of nodule formation. The limestone was extensively quarried for building stone and for supplying limekilns on the site.

The Derbyhaven area has been substantially covered by the recent runway extension. The Scarlett area is subjected to considerably less wave action than Port St Mary Ledges, and so the two areas are complementary as special intertidal sites worthy of protection.

Boundary

The boundary of the site has been set at Lowest Astronomical Tide.

Site Name: Port St Mary Ledges and Kallow PointOperations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseedling
2	Grazing where already damaging, the introduction of grazing and changes in grazing regime (including type of stock or intensity or seasonal pattern of grazing and cessation of grazing).
3	Stock feeding, where already damaging, the introduction of stock feeding and changes in stock feeding practice.
4	Mowing or other methods of cutting where they are already damaging, changes in mowing or cutting regime (including conversion from hay making to silage or cessation of mowing).
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16	Coastal fishing or fisheries management and seafood or marine life collection, where already damaging, the introduction of coastal fishing and changes in coastal fishing practices or fisheries management and seafood or marine life collection, including the use of traps and fish cages.
17	Reclamation of land from sea, estuary or marsh.
18	Bait digging on inter-tidal areas.
19	Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.

20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest,
27	Recreational or other activities likely to damage features of interest or soil, fauna and flora (excluding walking, with or without dogs).
28	Introduction of game or waterfowl management and changes in game or waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* "animal" includes any mammal, amphibian, reptile, bird, fish or invertebrate.



WILDLIFE ACT 1990

RAMSEY BAY (MARINE NATURE RESERVE) (No. 2) BYELAWS 2011

Approved by Tynwald: 20th January 2012

Coming into operation: 1st January 2012

The Department of Environment, Food and Agriculture, on application of the Wildlife Committee, makes these Byelaws under section 33 of the Wildlife Act 1990¹.

Part 1

Title, commencement, duration and interpretation

1 Title

These Byelaws are the Ramsey Bay (Marine Nature Reserve) Byelaws 2011.

2 Commencement and duration

These Byelaws shall come into operation on 1st January 2012.

3 Interpretation

In these Byelaws –

“the Act” means the Wildlife Act 1990;

“authorised person” means a person authorised in writing by the Department to exercise powers under these Byelaws;

“the Conservation Zone” is that area numbered as Zone 1 on the map in the Schedule and is that area to the West of the line running North from the point at 54° 18'.66N, 04° 21'.10W and known as Gob ny Rona up to, and including, the mean high water mark on the shoreline but not including any point inland from the piers at the entrance to Ramsey harbour and the harbour itself.

¹ 1990 c.2

“the Department” means the Department of Environment, Food and Agriculture;

“the Horse Mussel Conservation Zone” consists of that area of sea, up to and including the mean high water mark, running East of a line passing through the Point of Ayre Lighthouse being $04^{\circ} 22'.10W$ and to the North of a line which transects that line running North, being $54^{\circ} 24'.60N$. The area extending out to the nautical three-mile limit and marked as Zone 2 on the map contained in the Schedule;

“the Eelgrass Conservation Zone” is that area numbered as Zone 3 on the map in the Schedule and consists of the area South of the line running East from the point at $54^{\circ} 18'.90N$ and known as the Ballure Arches to a point where it meets the line running North at $4^{\circ} 21'.10W$ up to, and including the mean high water mark;

“the Fisheries Management Zone” is that area numbered as Zone 4 on the map in the Schedule and consists of the area to the East of the line running North at $4^{\circ} 21'.10W$ to the point where it meets the line running East at $54^{\circ} 24'.06N$ and extends to the nautical three-mile limit;

“gate gear” includes any appliance with a rigid frame mouth, but which does not have teeth, which is towed through the waters and is manufactured, adapted, used or intended for use for the purpose of fishing for scallops and queen scallops;

“gill and other specified nets” means any of the following, that is to say gill nets, beach seines, trammel nets, tangle nets, stake nets, drift nets, ring nets, T nets, J nets, hoop nets, any trawl, Danish seine or similarly towed net, and any similar nets used for the enmeshing or entrapment of fish, but does not include a landing net used in the course of normal angling by rod and line;

“longline” means fishing gear which comprises a main line carrying one or more hooks on branch lines (snoods) set either at or near the bottom or drifting in midwater or near the surface intended to catch fish;

“scallop dredge ” includes any appliance with a rigid frame mouth and teeth, which is towed through the waters and is manufactured, adapted, used or intended for use for the purpose of fishing for scallops or queen scallops;

“the Ramsey Bay Marine Nature Reserve” means the area indicated on the map in the Schedule and designated by the Department under the Ramsey Bay (Marine Nature Reserve) (Designation) Order 2011 and referred to in these Byelaws as “the Reserve”;

Part 2

General Restrictions within the Reserve

4 Prohibition of aggregate extraction

No person shall extract or remove any mineral, including sand, gravel, rock, oil, gas or coal from any part of the Reserve.

5 Prohibition of dumping of dredged material

No person shall cause to be dumped in any part of the Reserve any material that has been extracted by dredging.

6 Prohibition of netting

(1) No person shall use any gill or specified nets intended to capture any species of fish within the Reserve.

(2) But paragraph (1) does not prohibit the use of a landing net when being used as an auxiliary to angling with rod and line.

(3) Nor does paragraph (1) apply to a person in possession of a licence granted by the Department under bye-law 4(3) of the Sea-Fisheries (Protection of Migratory Fish Species) Bye-Laws 2005² or section 2AA of the Sea Fisheries Act 1971.

7 Prohibition of long-lining

No person shall use a long-line intended to capture any species of fish within the Reserve, including setting a long-line from the shore or any vessel.

8 Prohibition of dredging

(1) No person shall use any gate gear or scallop dredge within any part of the Reserve.

(2) Paragraph (1) does not apply to trawling, scallop-dredging or diving for scallops within the Fisheries Zone authorised by a licence issued by the Department.

² SD 689/05

9 Prohibition of littering

No person shall deposit, or cause, or suffer to be deposited, any paper, bottles, glass, rubbish or refuse within the Reserve.

10 Prohibition of construction

No person shall construct any pipeline or other structure within the Reserve except under the authority of a licence issued by the Department.

Part 3

Additional Restrictions in specified Zones

11 Restrictions within the Conservation Zone

No person may trawl, dredge or gather scallops or queen scallops by any means within the Conservation Zone.

12 Restrictions within the Horse Mussel Zone

No person may work a pot, trawl, dredge or extract scallops or queen scallops by any means within the Horse Mussel Zone.

13 Restrictions within the Eelgrass Zone

(1) This byelaw applies to the Eelgrass Zone.

(2) No person may –

(a) trawl;

(b) dredge;

(c) anchor;

(d) undertake recreational rod and line angling; or

(e) collect any living animal or plant on the foreshore or in the sea by any means, including hand gathering.

(3) But the taking of lugworm and razor clams from the shore (between Spring High Water Mark and Spring Low Water Mark) is only prohibited between 1 April and 30 September each year.

(4) Paragraph (3) above ceases to have effect on 31st March 2014.

Part 3
Research and investigations

14 Permit to undertake scientific research

Despite the prohibitions contained in these Bye-Laws, the Department may, if it considers it necessary, issue a permit to any person to undertake research or investigations within the Reserve and attach conditions and limitations to that permit.

Part 4
Obstruction and offences

15 Obstruction of an officer

No person shall wilfully obstruct –

(a) any officer of the Department in the execution of any research or scientific work or any work connected with the laying out, maintenance or management of the Reserve;

(b) any authorised person on the Reserve.

16 Production of authority

An authorised person shall, if required to do so, produce evidence of his authority under these Byelaws.

17 Offences

Any person contravening these Byelaws shall be guilty of an offence under section 33(5) of the Act.

18 Revocation

The Ramsey Bay (Marine Nature Reserve) Byelaws 2011³ are revoked.

MADE




8.12.11

Minister for Environment, Food and Agriculture

³ SD 0764/11

SCHEDULE RAMSEY BAY MARINE NATURE RESERVE



EXPLANATORY NOTE

(This note is not part of the Byelaws)

These Byelaws are consequent on the designation of the Ramsey Bay Marine Nature Reserve and specify restrictions on activities within the Reserve.

The Reserve consists of four 'Zones' –

- The Conservation Zone;
- The Horse Mussel Zone;
- The Fisheries Zone; and
- The Eelgrass Zone,

all of which have specific habitats which are protected under these Bye-Laws.

Contravention of any provision of these Byelaws is subject to a fine set down in section 33(5) of the Wildlife Act 1990.

Byelaw 4 prohibits the extraction of aggregate from any area of the Reserve.

Byelaw 5 prohibits the dumping of any dredged material in any part of the Reserve.

Byelaw 6 prohibits certain types of netting within the Reserve. These are specified in the definitions in byelaw 3.

Byelaw 7 prohibits longlining within the Reserve.

Byelaw 8 prohibits the use of any scallop dredge within the Reserve other than within the Fisheries Management Zone.

Byelaw 9 prohibits littering within the whole Reserve.

Byelaw 10 prohibits the construction of any pipeline or structure within the Reserve unless licensed by the Department.

Byelaw 11 prohibits the taking of scallops, by any means, within the Conservation Zone.

Byelaw 12 prohibits potting, trawling and dredging in the Horse Mussel Zone.

Byelaw 13 prohibits the extraction of living resources from the Eelgrass Zone, including the practice of catch and release angling. The collection of lugworm and razor clams will be permitted between 1 October and 31 March in the next year for a period of two years. It is intended to undertake a survey of the impact which this type of bait collection has in the Eelgrass Zone and provision is made for the practice to cease if necessary.

Byelaw 14 enables the Department to issue a permit for activities, which are otherwise prohibited within the Reserve, to be undertaken subject to conditions and limitations set down in that permit.

Byelaw 15 provides for an offence of obstructing any officer of the Department who is carrying out research, scientific work or any work connected with laying out, managing or maintaining the Reserve, or any authorised person.

Byelaw 16 provides that authorised persons shall, if required to do so, produce evidence of their authority under the Byelaws.

Byelaw 17 provides that any person contravening the Byelaws, including by failing to do something required by an authorised person, shall be guilty of an offence under section 33(5) of the Act; and

Under section 33(5) of the Act, a person found guilty of an offence at summary jurisdiction is liable to a fine of up to £2,500.

Byelaw 18 revokes previous Byelaws.

Ramsey Mooragh Shore Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

Thie Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



**Isle of Man
Government**

Keitrys Ellan Vannin



**Isle of Man
Government**
Raltys Eilan Varrin

Department of Agriculture,
Fisheries and Forestry

**RAMSEY MOORAGH SHORE
AREA OF SPECIAL
SCIENTIFIC INTEREST**

Area: 2.65ha (= 6.54 acres) *


Designated: 27th February 2006

Confirmed: 9th August 2006

*Note: the seaward boundary of the site is defined as the mean high water mark between points A and B on this map. The high water mark, and hence the overall site area, may be subject to variation with time.

The grid lines on this map form part of the National Grid, and are spaced at 500m intervals

Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Ramsey Mooragh Shore

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Ramsey Town

Planning Authority: Department of Local Government and the Environment

Local authority: Ramsey Town Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC 49SE, SC 49NE

National Grid Reference (centroid): SC452951

Area: 2.65ha (6.54 acres)

Date notified: 27th February 2006

Date confirmed: 9th August 2006

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its flora and habitat – vegetated strandline, vegetated shingle, semi-fixed dune and coastal grassland.

Description and reasons for notification:

Ramsey North Shore is a stretch of beach located between the Ramsey Mooragh Promenade and the mean high tide level. This length of shore corresponds to the small area in the centre of Ramsey Bay which has seabed sediments of sand, as opposed to gravel. This has resulted in a low sand dune forming against the promenade wall, with a zone of fine shingle on the seaward side, and associated vegetation and invertebrate life. Good examples of sandy strandline, vegetated shingle and dune grassland communities are all present on the site, which supports several rare or uncommon plants.

The sandy gravel on the seaward edge of the site has a continuous, stable strandline which has typical vegetated strandline species including sea rocket *Cakile maritima*, curled dock *Rumex crispus*, sea sandwort *Honckenya peploides*, and sand sedge *Carex arenaria*. The accumulated seaweed, driftwood and other organic flotsam supports abundant sandhoppers, which are particularly evident at high tide, when they in turn attract feeding birds including black redstart* and starlings*.

The vegetated strandline zone grades into foredune and semi-fixed “yellow” dune, often with very little foredune zone. The semi-fixed dune is pure sand at the north end of the site, and grades into fine shingle towards the south end. A wide range of coastal plant species is present, including prickly saltwort *Salsola kali*, marram grass *Ammophila arenaria*, sand sedge *Carex arenaria*, sand couch *Elytrigia juncea* ssp *boreoatlanticus*, sea mayweed *Matricaria maritima*, sea beet *Beta vulgaris* ssp *maritima*, sea bindweed *Calystegia soldanella*, sea plantain *Plantago maritima*, hare’s-foot clover *Trifolium arvense***, Isle of Man cabbage *Coincya monensis* ssp *monensis***, and abundant sea holly *Eryngium maritimum*, which is parasitised by conspicuous spikes of lesser broomrape *Orobanche minor*.

The most well-established zone of the dune is adjacent to the sea wall, where a range of coastal grassland species thrive, many of which are also characteristic of unimproved, neutral grassland (also a rare habitat on the Island). Species include lady’s bedstraw *Galium verum*, red fescue *Festuca rubra*, sheep’s fescue *Festuca ovina*, common cat’s-ear *Hypochoeris radicata*, wild carrot *Daucus carota*, ribwort plantain *Plantago lanceolata*, yarrow *Achillea millefolium*, rest harrow *Ononis repens*, kidney vetch *Anthyllis vulneraria* and bird’s-foot trefoil *Lotus corniculatus*.

Ramsey North Shore is significant in research terms, because it is the remaining Manx stronghold for Isle of Man cabbage, *Coincya monensis ssp monensis***. This species was the subject of the earliest known specific botanical record on the Island, as recorded in 1660 "...from the landing-place at Ramsey to the Town"[†]. Isle of Man cabbage is known to be in decline in all but its central range (northwest England), and has declined likewise on the Island. Ramsey North Shore is therefore a critical site for the monitoring and conservation of this species, both because of the continuity of records and because it is so suitable for the plant, and can provide seed for re-introduction programmes to suitable sites elsewhere.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 7 of the Wildlife Act 1990

[†] Rev J Ray, *Catalogus plantarum* 103 (per D.E. Allen, Flora of the Isle of Man, 1984 p67)

Other information:

As an additional point of botanical interest, the range of coastal habitats and vegetation communities present at Ramsey North Shore offers an important refuge for those plant species which are ephemeral in nature, and which do not appear at the same site year after year. Such species require a certain amount of suitable habitat in order to maintain their presence as part of the native flora of the Island, and will become extinct if no habitats are available for them to colonise on an occasional basis. Examples which have been recorded here in the past include several rare and protected species, such as oysterplant *Mertensia maritima***, dune fescue *Vulpia fasciculata*** and viper's bugloss *Echium vulgare***.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Ramsey Mooragh Shore

Operations likely to damage the special interest of the site

Type of operation

1. Cultivation including ploughing, rotovating, harrowing and reseeded.
2. The introduction of grazing.
3. The introduction of stock feeding and changes in stock feeding practice.
4. Mowing or other methods of cutting.
5. Application of manure, fertilisers and lime.
6. Application of pesticides, including herbicides (weedkillers).
7. Dumping, spreading or discharge of any materials.
8. Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9. The release into the site of any wild, feral or domestic animals *, plant or seed.
10. The killing or removal of any wild animal *, including pest control.
11. The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12. The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13. Reclamation of land from sea, estuary or marsh.
14. Erection of sea defences or coastal protection works, including cliff or land-slip drainage or stabilisation measures.
15. Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
16. Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
17. Storage of materials.
18. Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
19. Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
20. Removal of geological specimens, including rock samples, minerals and fossils.
21. Use of vehicles or craft likely to damage or disturb features of interest.
22. Recreational or other activities likely to damage features of interest (excluding walking, with or without dogs).

* "animal" includes any mammal, reptile, bird, fish or invertebrate.

Rosehill Quarry, Billown Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmiltaght, Bee as Eirinys

This Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS

www.gov.im



**Isle of Man
Government**

Reillys Ellan Vannin



**Isle of Man
Government**
Reilys Eilan Vannin

Department of Agriculture,
Fisheries and Forestry

**ROSEHILL QUARRY, BILLOWN
AREA OF SPECIAL
SCIENTIFIC INTEREST**

 Designation boundary


Area: 1.37ha (= 3.39 acres)

Designated: 2nd December 2005

Confirmed: 24th March 2006

The grid lines on this map form
part of the National Grid,
and are spaced at 50m intervals

Crown Copyright
Department of Local Government
and the Environment, Isle of Man.
Unauthorised production
infringes copyright.



NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site name: Rosehill Quarry, Billown

Status : Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Area: Isle of Man

Parish: Malew

Planning Authority: Department of Local Government and the Environment

Local authority: Malew Commissioners

Ordnance Survey Sheet: 1:50,000 OS Landranger Map No.95 and 1:10,560 sheets SC 27SE

National Grid Reference (centroid): SC 26856982

Area: 1.37 ha (3.39 acres)

Date notified: 2nd December 2005

Date confirmed: 24th March 2006

Date of last revision: N/A

Purpose

The purpose of this designation is to formalise the protection of an area which is important for its flora and habitat – open water, swamp vegetation, limestone grassland and scrub.

Description and reasons for notification:

Billown Quarry consists of a deeply-excavated, disused limestone quarry which has been allowed to flood and vegetate naturally. It now has a large, deep lake in the south part of the site and a small, shallow pool and marsh to the north. Calcareous grassland surrounds the water and also occurs on rock ledges around the quarry. The edges of the shallower lake have extensive fringes of swamp vegetation, grading into sedges and marshy grassland. The periphery of the site is sheltered all round by mixed hawthorn and gorse scrub, and trees of several stages of development.

The calcareous grassland in the centre of the site is species-rich, with limited grass cover and a high proportion of wild flowers including fairy flax *Linum catharticum*, wild carrot *Daucus carota*, selfheal *Prunella vulgaris*, common centaury *Centaureum erythraea*, common bird's-foot trefoil *Lotus corniculatus*, ox-eye daisy *Leucanthemum vulgare*, mouse-ear hawkweed *Pilosella officinalis*, black knapweed *Centaurea nigra* and eyebright *Euphrasia agg.* The sparse, fine grasses include crested dog's-tail *Cynosurum cristatum*, flattened meadow-grass *Poa compressa*, wood false-brome *Bromus sylvaticus* and red fescue *Festuca rubra ssp rubra*. The grassland is characterised by frequent patches of glaucous sedge *Carex flacca*, a species common to calcareous sites and consequently of rather local distribution on the Isle of Man. The site is notable for being one of only two sites on the Isle of Man for bee orchid *Ophrys apifera****. Overall, the vegetation is similar to that of the sheep's fescue *Festuca ovina* - meadow oat-grass *Avenula pratensis* grasslands – a habitat typical of the warmer, sunnier parts of the British Isles, found here at the northernmost edge of its range.

Both the two lakes in the flooded base of the quarry contain aquatic vegetation, notably stonewort *Chara* species, spiked water-milfoil *Myriophyllum spicatum* and curled pondweed *Potamogeton crispus*. All three species have a local distribution on the Island, due to the limited number of suitable habitats. The smaller and shallower of the two lakes supports a dense, varied fringe of emergent plants as well as submerged vegetation. Emergent species include bulrush *Typha latifolia*, common spike-rush *Eleocharis palustris*, mare's-tail *Hippuris vulgaris*, water horsetail *Equisetum fluviatile*, carnation sedge *Carex panicea*, lesser spearwort *Ranunculus flammula*, and water mint *Mentha aquatica*.

Birds, bats and invertebrates benefit from the sheltered environment of the quarry, which is provided partly by its depth and partly by the margins of developing scrub and woodland. At least fourteen bird species are known or thought to breed on the site include song thrush *Turdus philomelos**, willow warbler *Phylloscopus trochylus*, blackcap *Sylvia atricapilla*, blackbird *Turdus merula*, and mistle thrush *Turdus viscivorus*. The quarry is also known to attract feeding swallows *Hirundo rustica* and sand martins *Riparia riparia**, particularly in spring when suitable sheltered feeding areas are scarce. Invertebrates known to rely on the lakes and surrounding cover include common blue and blue-tailed damselflies *Enallagma cyathigerum* and *Ischnura elegans*, and the common darter dragonfly *Sympetrum striolatum*. In addition to birds and invertebrates, the site is known to attract feeding pipistrelle bats *Pipistrellus sp***.

The quarry is of interest to geologists, providing clear, accessible exposures of jointed, folded Carboniferous limestone, interspersed with impermeable shales and overlain with glacial boulder clay deposits.

Although a small site, Billown Quarry qualifies for ASSI selection because it fulfils the Priority Site Criteria for size, location, diversity, naturalness, representativeness and rarity, and research value, and for the presence of the features described above.

* = Protected under Schedule 1 of the Wildlife Act 1990

** = Protected under Schedule 5 of the Wildlife Act 1990

*** = Protected under Schedule 7 of the Wildlife Act 1990

Other information:

Other species thought likely to inhabit the site include the common frog *Rana temporaria* and common lizard *Lacerta vivipara*, both of which are protected under Schedule 5 of the Wildlife Act 1990. Bat species other than pipistrelles may also use the quarry for feeding or roosting. Further research on these species and on the possible presence of rare or local invertebrates and lower plants is warranted, due to the very limited Manx distribution of calcareous habitats.

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Site Name: Rosehill Quarry, Billown

Operations likely to damage the special interest of the site

<u>Standard reference number†</u>	<u>Type of operation</u>
1	Cultivation including ploughing, rotovating, harrowing and reseeded.
2	The introduction of livestock.
4	Introduction of mowing regime or other methods of cutting.
5	Application of manure, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers).
7	Dumping, spreading or discharge of any materials.
8	Burning, lighting of fires and changes in frequency or pattern of burning, where applicable.
9	The release into the site of any wild, feral or domestic animals *, plant or seed.
10	The killing or removal of any wild animal *, including pest control
11	The destruction, displacement, removal or cutting of any plant or plant remains, including herb, dead or decaying wood, moss, lichen, fungus, leaf mould or turf.
12	The introduction of tree and /or woodland management and changes in tree and/ or woodland management.
13	Drainage (including moor-gripping, the use of mole, tile, tunnel or other artificial drains).
14	The changing of water levels and tables and water utilisation (including irrigation, storage and abstraction from existing water bodies and through boreholes).
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
20	Extraction of minerals including peat, shingle, sand and gravel, topsoil, subsoil, chalk, lime, limestone pavement, shells and spoil.
21	Construction, removal or destruction of roads, tracks, walls, fences, hardstands, banks, ditches or other earthworks, or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures, or the undertaking of engineering works, including drilling.
24	Modification of natural or man-made features (including cave entrances), clearance of boulders, large stones, loose rock or scree and bettering, buttressing or grading rock faces and cuttings, in-filling of pits, and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft likely to damage or disturb features of interest.
27	Recreational or other activities likely to damage features of interest.
28	Game and waterfowl management and hunting practices.

† Note: each type of operation has a standard reference number; for each site, only those operations which are relevant to the site will be listed, hence there may be gaps in the numbering for some sites.

* “animal” includes any mammal, reptile, bird, fish or invertebrate.

Santon Gorge & Port Soldrick Area of Special Scientific Interest (ASSI)

Designation Documents

Please note:

- Notification as an ASSI confers no public right of entry to any land without the permission of the landowner.
- The citations and lists of operations requiring the Department of Environment, Food and Agriculture's consent may be subject to minor editorial changes and should not be assumed to be an exact facsimile of the original legal document.

Department of Environment, Food and Agriculture

Rheynn Chymmhtaght, Bee as Eirinys

This Slieau Whallian, Foxdale Road, St John's, Isle of Man IM4 3AS
www.gov.im



Isle of Man
Government

Reilly: Eilan Vannin

NOTIFICATION OF AN AREA OF SPECIAL SCIENTIFIC INTEREST Appendix I

Site Name: Santon Gorge and Port Soldrick **Parish(es):** Malew and Santon

Status: Area of Special Scientific Interest (ASSI) notified under Section 27 of the Wildlife Act 1990.

Planning Authority: Department of Infrastructure

Local Authorities: Malew Commissioners and Santon Commissioners

National Grid Reference (centroid): SC 2984 6924 **Area:** 24.35 hectares
(60.17 acres)

Ordnance Survey Sheet: **1:50,000:** OS Landranger Map No.95
1:10,000: OS Sheets SC 26 NE and 37 SW

Notification Date: 27th July 2012

Confirmation Date: 20th November 2012

Reasons for notification:

Santon Gorge is situated on the boundary between carboniferous limestone and Manx slate and is an important site primarily for the semi-natural woodland that exists on the gorge cliffs on either side of the Santon Burn. The woodland exists in association with many other habitats including scrub, coastal grassland, species-rich neutral marsh, coastal grassland and flushes and saltmarsh as well as an extensive area of inter-tidal habitat. These habitats support a large number of breeding birds.

General description:

The woodland at Santon Gorge forms the lower part of extensive semi-natural riparian woodland that extends several kilometres upstream along the banks of the Santon Burn. This lower section is the most botanically important and contains approximately 0.7 hectares of relic oak woodland within a larger area of 1 hectare of species-rich secondary semi-natural woodland.

The oak trees *Quercus x rosecea* would appear to be of coppice origin and are some of the Island's few relic ancient trees with stems over 200 years old. Other tree species include alder and ash. Many ancient woodland indicator species occur here, such as bitter vetchling *Lathyrus linifolius*, hazel *Corylus avellana* and aspen *Populus tremula*.

It is the only Island location for the hawkweed *Hieracium scabrecetum* which is a British Isles endemic and spindle *Euonymus europaeus*. Santon Gorge was a former site for hairy brome *Bromopsis ramosus* but this plant has not been recorded on the site since 2005, though it may still be re-found. Southern polypody *Polypodium cambricum* grows at the woodland fringe not far from the river and this location is one of only two sites on the Island for the species. Just south of the bridge at the northern end of the ASSI are a group of

veteran sycamores. Despite being non-native they are an impressive feature and may provide roost habitat for bats.

Blackthorn *Prunus spinosa*, hawthorn *Crataegus monogyna*, burnet rose *Rosa pimpinellifolia* and European gorse *Ulex europaeus* scrub is intimately mixed with coastal grassland and bracken.

The coastal scrub is contiguous with the riparian woodland. At the boundary between the woodland and scrub the woodland is naturally regenerating with ash *Fraxinus excelsior*, oak and sycamore *Acer pseudoplatanus* encroaching.

The most significant species amongst the scrub is the wood small-reed *Calamagrostis epigejos* that grows between and sometimes through the more patchy areas of scrub at Port Soldrick. The scrub at Port Soldrick is composed of blackthorn and there are occasional trees such as ash and notably one oak (*Quercus x rosacea*), indicating that the most sheltered parts of the bay will become woodland over time.

The promontory fort Cass ny Hawin (Grid Ref SC 297 693) contains a very species-rich calcareous grassland, a rare habitat on the Island. The area is sufficiently high above the open sea to only have minimal direct maritime influence. Due to the presence of the Fort the grassland has been afforded protection from agricultural improvement such as ploughing and fertilising. It is well grazed and at the peak of the monument the habitat is very dry and rich in plants such as knotted clover *Trifolium striatum*, kidney vetch *Anthyllis vulneraria* and spring sedge *Carex caryophyllea*. Spring sedge is a particularly notable species in the area.

Unimproved and very species-rich marsh occurs on the fringe of the woodland areas often intimately mixed with grazed woodland. The marshes are fed by permanent fresh water springs and grade from fresh spring habitat, with ivy-leaved crowfoot and watercress, to more stagnant and species-rich areas dominated by rush with ragged robin *Lychnis flos-cuculi*, water forget-me-not *Myosotis scorpioides* and water pepper *Persicaria hydropiper*.

A small and unmodified area of saltmarsh occurs in the lower reaches of the Santon Burn just before the gorge opens out on to the beach. The saltmarsh is dominated by red fescue *Festuca rubra* and saltmarsh rush *Juncus gerardii* with sea milkwort and sea club rush *Bolboschoenus maritimus* also present.

The Santon Burn is regarded as one of the most natural river systems on the Island. Salmon *Salmo salar*, sea trout *S. trutta morpha lacustris* and brown trout *S. trutta morpha fario* are present. Salmon and sea trout both migrate to and from the sea through Santon Gorge.

A line of freshwater springs from Santon Gorge to Port Soldrick create an extensive area of flush habitats along the coast. These flushes are species-rich and grade into brackish marsh towards the sea. Notable species include many-stalked spike rush *Eleocharis multicaulis*, slender clubrush *Isolepis cernua* and bog pimpinell *Anagallis tenella*.

The grazed slopes are species-rich and occur alongside coastal flushes. Typical species include thrift *Armeria maritima* and spring squill *Scilla verna*.

The mouth of the Santon Burn and Port Soldrick contain significant areas of un-vegetated and sparsely-vegetated shingle. Accumulations of seaweed make these areas attractive to feeding chough which nest on the coastal cliffs.

The shore to the north of the Runway extension includes limestone ledges that are more exposed than those to the south of the runway. They are diverse, with knotted wrack *Ascophyllum nodosum* and acorn barnacles *Semibalanus balanoides* and a number of pools containing beadlet anemones *Actinia equina* and other species. The "Runway Lights" area which was mainly lost when the Runway Extension Safety Area (RESA) was constructed was found to be the most species-rich site of all those surveyed on the Isle of Man, and the North Ronaldsway site also showed relatively high species richness. From an intertidal point of view, there is a natural boundary from where the RESA finishes up to Port Soldrick, including a diverse mosaic of rocky shore and shingle beaches, with freshwater input and a range of exposure levels.


The shores of the coves at Santon Gorge and Port Soldrick area are barren shingle and gravel, a common habitat in the Isle of Man but one considered uncommon in the UK. The upper shore along the coast between the two coves is rocky with a covering of lichens, including black tar lichen *Verrucaria maura*.


The lower shore throughout the site is dominated by fucoid (brown seaweeds) on rock. There are large areas of dense knotted wrack *Ascophyllum nodosum* on the mid shore to the south of Santon Gorge and to the south of Port Soldrick. To the north and south of this, the mid shore is dominated by bladder wrack *Fucus vesiculosus*, barnacle and limpet *Patella* sp. mosaic whilst the lower shore boulders are covered in serrated wrack *Fucus serratus*. The extreme lower shore from Ronaldsway to Santon Gorge is dominated by oarweed *Laminaria digitata* on moderately exposed sublittoral fringe rock. The input of freshwater at Santon Gorge creates an unusual freshwater-influenced intertidal environment, dominated by *Enteromorpha* spp on rock.

Additional information

Geological importance

At the mouth of the Santon Burn the Ordovician Manx Group rocks meets Carboniferous Limestone of the Derbyhaven Formation. The limestone beds tilt against the Ordovician strata along the boundary fault. In the inter-tidal zone of the west side of the gorge runs an Ordovician age igneous dyke which probably pushed through surrounding seabed sediment before it became rock and is irregularly shaped as a result. The Santon Gorge area is a candidate Manx Regionally Important Geological Site (RIGS).





NOTIFICATION OF SANTON GORGE AND PORT SOLDRICK AREA OF SPECIAL SCIENTIFIC INTEREST Appendix II

Standard reference number	Type of operation
1	Cultivation, including ploughing, rotovating, harrowing and re-seeding.
2	Grazing, the introduction of grazing and alterations to the grazing regime (including type of stock, intensity or seasonal pattern of grazing).
3	Stock feeding, the introduction of stock feeding and alterations to stock feeding practice.
4	Mowing or cutting of vegetation (where already damaging), the introduction of mowing and alterations to the mowing or cutting regime (such as from haymaking to silage).
5	Application of manure, slurry, silage liquor, fertilisers and lime.
6	Application of pesticides, including herbicides (weedkillers) whether terrestrial or aquatic, and veterinary products.
7	Dumping, spreading or discharge of any materials.
8	Burning and alterations to the pattern or frequency of burning.
9	Release into the site of any wild, feral, captive-bred or domestic animal, plant, seed or micro-organism (including genetically modified organisms).
10	Killing, injuring, taking or removal of any wild animal (including dead animals or parts thereof), or their eggs and nests, including pest control and disturbing them in their places of shelter.
11	Destruction, displacement, removal or cutting of any plant or plant remains, including tree, shrub, herb, hedge, dead or decaying wood, moss, lichen, fungus, leaf-mould, and turf.
12	Tree and/or woodland management (where already damaging), the introduction of tree and/or woodland management (where applicable) and alterations to tree and/or woodland management (including planting, felling, pruning and tree surgery, thinning coppicing, changes in species composition, removal of fallen timber).
13 a)	Draining (including moor- gripping, the use of mole, tile, tunnel or other artificial drains).
13 b)	Modification to the structure of water courses (e.g. rivers, streams, springs, ditches and drains), including their banks and beds, as by re-alignment, regarding, damming, or dredging.
13 c)	Management of aquatic and bank vegetation for drainage purposes.
14	Alterations to water levels and tables and water utilisation (including irrigation, storage, abstraction from existing water bodies and through boreholes). Also the modification of current drainage practices.
15	Infilling of ditches, dykes, drains, ponds, pools, marshes or pits.
16 a)	Freshwater fishery production and/or management, including sporting fishing and angling (where already damaging), the introduction of freshwater production and/or management (where applicable) and alterations to freshwater fishery production and/or management.
16 b)	Coastal fishing, fisheries management and seafood or marine life collection, including the use of traps or fish cages (where already damaging), the introduction of coastal fishing (where applicable), alterations to coastal fishing practice or fisheries management and seafood or marine life collection.
17	Reclamation of land from sea, estuary or marsh.
18	Bait digging in inter-tidal areas.
19	Erection and repair of sea defences or coastal protection works, including cliff or landslip drainage or stabilisation measures.
20	Extraction of minerals including peat, shingle, hard rock, sand, gravel, topsoil, subsoil, lime, limestone pavement, shells and spoil.
21	Destruction, construction, removal, rerouting, or regarding of roads, tracks, walls,


	fences, hardstands, banks, ditches or other earthworks, including soil and soft rock exposures or the laying, maintenance or removal of pipelines and cables, above or below ground.
22	Storage of materials.
23	Erection of permanent or temporary structures or the undertaking of engineering works, including drilling.
24 a)	Modification of natural or man-made features (including cave entrances) and clearance of boulders, large stones, loose rock or scree.
24 b)	Battering, buttressing or grading of geological exposures and cuttings (rock and soil) and infilling of pits and quarries.
25	Removal of geological specimens, including rock samples, minerals and fossils.
26	Use of vehicles or craft.
27	Recreational or other activities.
28 a)	Game and waterfowl management and hunting practices (where already damaging), introduction of game or waterfowl management (where applicable) and alterations to game and waterfowl management and hunting practice.
28 b)	Use of lead shot.

Notes:

1. This is a list of operations appearing to DEFA to be likely to damage the special features of this ASSI, as required under Section 27 of the Wildlife Act 1990.
2. Any reference to 'animal' in this list shall be taken to include any mammal, reptile, amphibian, bird, fish or invertebrate.

Date Notified: 27th July 2012

Date Confirmed: 20th November 2012



Isle of Man

Ellan Vannin

AT 2 of 1990

WILDLIFE ACT 1990



Isle of Man

Ellan Vannin

WILDLIFE ACT 1990

Index

Section	Page
PART I – WILDLIFE	7
<i>Protection of birds</i>	7
1 Protection of wild birds, their nests and eggs	7
2 Exceptions to s 1	9
3 Areas of special protection: birds	10
4 Exceptions to ss 1 and 3	11
5 Prohibition of certain methods of killing or taking wild birds	12
6 Sale etc of live or dead wild birds, eggs etc	14
7 Registration etc of persons in possession of certain captive birds.....	15
8 Protection of captive birds.....	17
<i>Protection of other animals</i>	18
9 Protection of certain wild animals.....	18
10 Exceptions to s 9.....	19
11 Prohibition of certain methods of killing or taking wild animals.....	20
<i>Protection of plants</i>	22
12 Protection of wild plants.....	22
<i>Miscellaneous</i>	23
13 Areas of special protection: animals and plants.....	23
14 Introduction of new species etc	24
15 [Repealed]	25
<i>Supplemental</i>	25
16 Power to grant licences	25
17 False statements made for obtaining registration or licence etc	27
18 Prohibition of hunting with dogs	27
19 Possession of implements etc.....	28
20 Enforcement.....	28
21 Summary prosecutions	29
22 Penalties, forfeitures etc.....	29
23 Power to vary Schedules.....	30
24 Advisory bodies and their functions	31

25	Regulations, orders, notices etc	31
26	Interpretation of Part I	32
26A	Exemption for certain flood risk management functions	33

PART II – NATURE CONSERVATION **34**

27	Areas of special scientific interest	34
28	Restoration where s 27 contravened	35
29	Compensation	36
30	Management agreements with owners etc of land	37
31	National nature reserves	38
32	Marine nature reserves	40
33	Byelaws for protection of marine nature reserves	40

PART III – MISCELLANEOUS AND GENERAL **42**

34	Offences by bodies corporate etc	42
35	Publication of information	42
36	Duty to have regard to environment etc	42
37	Financial provisions	43
38	General interpretation	43
39	Consequential amendments	43
40	Repeals	44
41	Short title and commencement	44

SCHEDULE 1 **45**

BIRDS WHICH ARE PROTECTED BY SPECIAL PENALTIES	45
--	----

SCHEDULE 2 **47**

BIRDS WHICH MAY BE KILLED OR TAKEN	47
------------------------------------	----

SCHEDULE 3 **48**

BIRDS WHICH MAY BE SOLD	48
-------------------------	----

SCHEDULE 4 **49**

CAPTIVE BIRDS WHICH MAY BE POSSESSED BY REGISTERED PERSONS ONLY	49
---	----

SCHEDULE 5 **51**

ANIMALS WHICH ARE PROTECTED	51
-----------------------------	----

SCHEDULE 6 **51**

ANIMALS WHICH MAY NOT BE KILLED OR TAKEN BY CERTAIN METHODS	51
---	----

SCHEDULE 7	52
PLANTS WHICH ARE PROTECTED	52
SCHEDULE 8	54
ANIMALS AND PLANTS TO WHICH SECTION 14 APPLIES	54
SCHEDULE 9	55
SCHEDULE 10	56
REPEAL OF ENACTMENTS	56
ENDNOTES	57
TABLE OF LEGISLATION HISTORY	57
TABLE OF RENUMBERED PROVISIONS	57
TABLE OF ENDNOTE REFERENCES	57

**Isle of Man***Ellan Vannin*

WILDLIFE ACT 1990

Received Royal Assent: 20 February 1990
Passed: 20 February 1990
Commenced: 1 January 1991

AN ACT to repeal and re-enact with amendments legislation for the protection of birds; to make new provision for the conservation of wild creatures and wild plants; to prohibit certain methods of killing or taking wild animals; to restrict the introduction of certain animals and plants; to amend certain related enactments; to make new provision relating to nature conservation; and for connected purposes.

GENERAL NOTE

1. The maximum fines in this Act are as increased by the *Criminal Justice (Penalties, Etc.) Act 1993* s 1.
2. Sections 1 to 8 not operative for bird sanctuaries in existence immediately prior to 1/1/1991.

PART I – WILDLIFE

Protection of birds

1 Protection of wild birds, their nests and eggs

[P1981/69/1]

- (1) Subject to the provisions of this Part, if any person intentionally or recklessly —
- (a) kills, injures or takes any wild bird;
 - (b) takes, damages or destroys the nest of any wild bird while that nest is in use or being built; or
 - (c) takes or destroys an egg of any wild bird,
- he shall be guilty of an offence.¹

- (2) Subject to the provisions of this Part, if any person has in his possession or control —
- (a) any live or dead wild bird or any part of, or anything derived from, such a bird; or
 - (b) an egg of a wild bird or any part of such an egg,
- he shall be guilty of an offence.
- (3) A person shall not be guilty of an offence under subsection (2) if he shows that —
- (a) the bird or egg had not been killed or taken, or had been killed or taken otherwise than in contravention of the relevant provisions; or
 - (b) the bird, egg or other thing in his possession or control had been sold (whether to him or any other person) otherwise than in contravention of those provisions;
- and in this subsection “the relevant provisions” means the provisions of this Part and of orders made under it and, in the case of a bird, egg or other thing falling within subsection (2), the provisions of the Protection of Birds Acts 1932 to 1975 and of orders made under those Acts.
- (4) Any person convicted of an offence under subsection (1) or (2) in respect of —
- (a) a bird included in Schedule 1 or any part of, or anything derived from, such a bird;
 - (b) the nest of such a bird; or
 - (c) an egg of such a bird or any part of such an egg,
- shall be liable to a special penalty.
- (5) Subject to the provisions of this Part, if any person intentionally or recklessly —
- (a) disturbs any wild bird included in Schedule 1 while it is building a nest or is in, on or near a nest containing eggs or young; or
 - (b) disturbs any nest or egg of such a bird; or
 - (c) disturbs dependent young of such a bird,
- he shall be guilty of an offence and liable to a special penalty.²
- (6) In this section “wild bird” does not include any bird which is shown to have been bred in captivity.
- (7) Any reference in this Part to any bird included in Schedule 1 is a reference to any bird included in Part I and, during the close season for the bird in question, any bird for the time being included in Part II of that Schedule.

2 Exceptions to s 1

[P1981/69/2]

- (1) Subject to the provisions of this section, a person shall not be guilty of an offence under section 1 by reason of the killing or taking of a bird included in Part I of Schedule 2 outside the close season for that bird, or the injuring of such a bird outside that season in the course of an attempt to kill it.
- (2) Subject to the provisions of this section, an authorised person shall not be guilty of an offence under section 1 by reason of —
 - (a) the killing or taking of a bird included in Part II of Schedule 2, or the injuring of such a bird in the course of an attempt to kill it;
 - (b) the taking, damaging or destruction of a nest of such a bird; or
 - (c) the taking or destruction of an egg of such a bird.
- (3) Subsections (1) and (2) shall not apply on Sundays in any area which the Department of Environment, Food and Agriculture (in this Act referred to as “the Department”) may by order prescribe for the purposes of those subsections.³
- (4) In this section and section 1 “**close season**” means —
 - (a) in the case of woodcock, the period in any year commencing with 1st February and ending with 30th September;
 - (b) in any other case, subject to the provisions of this Part, the period in any year commencing with 1st February and ending with 31st August.
- (5) The Department may by order made with respect to the whole or any specified part of the Island vary the close season for any wild bird specified in the order.
- (6) If it appears to the Department expedient that any wild birds included in Part II of Schedule 1 or Part I of Schedule 2 should be protected during any period outside the close season for those birds, it may by order made with respect to the whole or any specified part of the Island declare any period (which shall not in the case of any order exceed fourteen days) as a period of special protection for those birds; and this section and section 1 shall have effect as if any period of special protection declared under this subsection for any birds formed part of the close season for those birds.
- (7) Before making an order under subsection (6) the Department shall consult such bodies and organisations as appear to it to be appropriate.

3 Areas of special protection: birds

[P1981/69/3]

- (1) The Department may by order make provision with respect to any area specified in the order providing for all or any of the following matters, that is to say —
 - (a) that any person who, within that area or any part of it specified in the order, at any time or during any period so specified, intentionally or recklessly —
 - (i) kills, injures or takes any wild bird, game bird or poultry;⁴
 - (ii) takes, damages or destroys the nest of such a bird while that nest is in use or being built;
 - (iii) takes or destroys an egg of such a bird;
 - (iv) disturbs such a bird while it is building a nest or is in, on or near a nest containing eggs or young; or
 - (v) disturbs any nest or egg of such a bird; or
 - (vi) disturbs dependent young of such a bird,shall be guilty of an offence under this section;⁵
 - (b) that any person who, except as may be provided in the order, enters into that area or any part of it specified in the order at any time or during any period so specified shall be guilty of an offence under this section;
 - (c) that where any offence under this Part, or any such offence under this Part as may be specified in the order, is committed within that area, the offender shall be liable to a special penalty.
- (2) A person authorised in writing by the Department or, where it is necessary for the protection of the water supply, the Manx Utilities Authority, shall not by virtue of any such order be guilty of an offence by reason of —
 - (a) the killing or taking of a bird included in Part II of Schedule 2, or the injuring of such a bird in the course of an attempt to kill it;
 - (b) the taking, damaging or destruction of the nest of such a bird;
 - (c) the taking or destruction of an egg of such a bird; or
 - (d) the disturbance of such a bird or the nest, egg or dependent young of such a bird.⁶
- (3) The making of any order under this section with respect to any area shall not affect the exercise by any person of any right vested in him, whether as owner, lessee or occupier of any land in that area or by virtue of a licence or agreement.
- (4) Before making any order under this section the Department shall give particulars of the intended order either by notice in writing to every owner and every occupier of any land included in the area with respect

to which the order is to be made or, where the giving of such a notice is in its opinion impracticable, by advertisement in a newspaper published and circulating in the Island.

- (5) The Department shall not make an order under this section unless —
- (a) all the owners and occupiers aforesaid have consented thereto; or
 - (b) no objections thereto have been made by any of those owners or occupiers before the expiration of a period of 3 months from the date of the giving of the notice or the publication of the advertisement; or
 - (c) any such objections so made have been withdrawn.
- (6) Subsection (5) shall not apply to an order under this section which is made with respect only to an area falling within, or forming the whole of, a district defined in an order under section 11 of the *Wild Birds Protection Act 1932* (bird sanctuaries) where the latter order was in operation immediately before the commencement of the order made under this section.⁷

4 Exceptions to ss 1 and 3

[P1981/69/4]

- (1) Nothing in section 1 or in any order made under section 3 shall make unlawful —
- (a) anything done in pursuance of a requirement by the Department under section 1 of the *Prevention of Damage by Agricultural Pests Act 1956*;
 - (b) anything done under section 3 of the *Forestry Act 1984*; or
 - (c) anything done under, or in pursuance of an order made under, any provision of the Diseases of Animals (Prevention) Acts 1948 to 1975.
- (2) Notwithstanding anything in section 1 or any order made under section 3, a person shall not be guilty of an offence by reason of —
- (a) the taking of any wild bird if he shows that the bird had been disabled otherwise than by his unlawful act and was taken solely for the purpose of tending it and releasing it when no longer disabled;
 - (b) the killing of any wild bird if he shows that the bird had been so seriously disabled otherwise than by his unlawful act that there was no reasonable chance of its recovering; or
 - (c) any act made unlawful by those provisions if he shows that the act was the incidental result of a lawful operation and could not reasonably have been avoided.
- (3) Notwithstanding anything in section 1 or any order made under section 3, an authorised person shall not be guilty of an offence by reason

of the killing or injuring of any wild bird, other than a bird included in Schedule 1, if he shows that his action was necessary for the purpose of —

- (a) preserving public health, public safety or air safety;
 - (b) preventing the spread of disease; or
 - (c) preventing serious damage to livestock, foodstuffs for livestock, crops, vegetables, fruit, growing timber, or fisheries.
- (4) An authorised person shall not be entitled to rely on the defence provided by subsection (3)(c) as respects any action taken at any time if it had become apparent, before that time, that the action would prove necessary for the purpose mentioned in that subsection and either —
- (a) a licence under section 16 authorising that action had not been applied for as soon as reasonably practicable after that fact had become apparent; or
 - (b) an application for such a licence had been determined.

5 Prohibition of certain methods of killing or taking wild birds

[P1981/69/5]

- (1) Subject to the provisions of this Part, if any person —
- (a) sets in position any of the following articles, being an article which is of such a nature and is so placed as to be calculated to cause bodily injury to any wild bird coming in to contact therewith, that is to say, any spring, trap, gin, snare, hook and line, any electrical device for killing, stunning or frightening, or any poisonous, poisoned or stupefying substance;⁸
 - (b) uses, or causes to be used, for the purpose of killing or taking any wild bird any such article as aforesaid, whether or not of such a nature and so placed as aforesaid, or any net, baited board, bird-lime or substance of a like nature to bird-lime;
 - (c) uses, for the purpose of killing or taking any wild bird —
 - (i) any bow, crossbow or catapult;
 - (ii) any explosive other than ammunition for a firearm;
 - (iii) any automatic or semi-automatic weapon;
 - (iv) any shot-gun of which the barrel has an internal diameter at the muzzle of more than one and three-quarter inches;
 - (v) any device for illuminating a target or any sighting for night shooting;
 - (vi) any form of artificial lighting or any mirror or other dazzling device;
 - (vii) any gas or smoke not falling within paragraphs (a) and (b); or

- (viii) any chemical wetting agent;⁹
- (d) uses, as a decoy, for the purpose of killing or taking any wild bird, any sound recording or any live bird or other animal whatever which is tethered, or which is secured by means of braces or other similar appliance, or which is blind, maimed or injured; or¹⁰
- (e) uses any mechanically propelled vehicle in immediate pursuit of a wild bird for the purpose of killing or taking that bird; or¹¹
- (f) knowingly causes or permits to be done an act which is mentioned in paragraphs (a) to (e),¹²

he shall be guilty of an offence and be liable to a special penalty.

- (2) Subject to subsection (3), the Department may by order, either generally or in relation to any kind of wild bird specified in the order, amend subsection (1) by adding any method of killing or taking wild birds or by omitting any such method which is mentioned in that subsection.
- (3) The power conferred by subsection (2) shall not be exerciseable, except for the purpose of complying with an international obligation, in relation to any method of killing or taking wild birds which involves the use of a firearm.
- (4) In any proceedings under subsection (1)(a) or under subsection (1)(f) relating to an act which is mentioned in subsection (1)(a) it shall be a defence to show that the article was set in position for the purpose of killing or taking, in the interests of public health, agriculture, forestry, fisheries, or nature conservation, any wild animals which could be lawfully killed or taken by those means and that he took or caused to be taken all reasonable precautions to prevent injury thereby to wild birds.¹³
- (5) Nothing in subsection (1) shall make unlawful —
 - (a) the use of a cage-trap or net by an authorised person for the purpose of taking a bird included in Part II of Schedule 2; or
 - (b) the use of a cage-trap or net for the purpose of taking any game bird if it is shown that the taking of the bird is solely for the purpose of breeding;

but nothing in this subsection shall make lawful the use of any net for taking birds in flight or the use for taking birds on the ground of any net which is projected or propelled otherwise than by hand.

- (6) In any proceedings under subsection (1)(c)(iii) it shall be a defence to show that the automatic or semi-automatic weapon was used by an authorised person for the purpose of preventing serious damage to livestock, foodstuffs for livestock, crops, vegetables, fruit, growing timber or fisheries by any bird included in Part II of Schedule 2.

6 Sale etc of live or dead wild birds, eggs etc

[P1981/69/6]

- (1) Subject to the provisions of this Part, if any person —
- (a) sells, offers or exposes for sale, or has in his possession or transports for the purpose of sale, any live wild bird other than a bird included in Part I of Schedule 3, or an egg of a wild bird or any part of such an egg; or
 - (b) publishes or causes to be published any advertisement likely to be understood as conveying that he buys or sells, or intends to sell, any of those things,
- he shall be guilty of an offence.
- (2) Subject to the provisions of this Part, if any person who is not for the time being registered in accordance with regulations made by the Department —
- (a) sells, offers or exposes for sale, or has in his possession or transports for the purpose of sale, any dead wild bird other than a bird included in Part II or III of Schedule 3, or any part of, or anything derived from such a wild bird; or
 - (b) publishes or causes to be published any advertisement likely to be understood as conveying that he buys or sells, or intends to buy or sell, any of those things,
- he shall be guilty of an offence.
- (3) Subject to the provisions of this Part, if any person sells, offers or exposes for sale, or has in his possession or transports for the purpose of sale, any dead wild bird which has been killed or taken in contravention of this Part he shall be guilty of an offence.
- (4) Subject to the provisions of this Part, if any person shows or causes or permits to be shown for the purposes of any competition or in any premises in which a competition is being held —
- (a) any live wild bird other than a bird included in Part I of Schedule 3; or
 - (b) any live bird one of whose parents was such a wild bird,
- he shall be guilty of an offence.
- (5) Any person convicted of an offence under this section in respect of —
- (a) a bird included in Schedule 1 or any part of, or anything derived from, such a bird; or
 - (b) an egg of such bird or any part of such an egg,
- shall be liable to a special penalty.
- (6) Any reference in this section to any bird included in Part I of Schedule 3 is a reference to any bird included in that Part which was bred in

captivity and has been ringed or marked in accordance with regulations made by the Department.

- (7) Any reference in this section to any bird included in Part II or III of Schedule 3 is a reference to any bird included in Part II and, during the period commencing with 1st September in any year and ending with 28th February of the following year, any bird included in Part III of that Schedule.
- (8) The power of the Department to make regulations under subsection (2) shall include power —
 - (a) to impose requirements as to the carrying out by a person registered in accordance with the regulations of any act which, apart from the registration, would constitute an offence under this section; and
 - (b) to provide that any contravention of the regulations shall constitute such an offence.
- (9) Regulations under subsection (2) shall secure that no person shall become or remain registered —
 - (a) within five years of his having been convicted of an offence under this Part for which a special penalty is provided; or
 - (b) within three years of his having been convicted of any other offence under this Part so far as it relates to the protection of birds or other animals or any offence involving their ill-treatment,no account being taken for this purpose of a conviction which has become spent for the purposes of the *Rehabilitation of Offenders Act 2001*.¹⁴
- (10) Any person authorised in writing by the Department may, at any reasonable time and (if required to do so) upon producing evidence that he is authorised, enter and inspect any premises where a registered person keeps any wild birds for the purpose of ascertaining whether an offence under this section is being, or has been, committed on those premises.
- (11) Any person who intentionally obstructs a person acting in the exercise of the power conferred by subsection (10) shall be guilty of an offence.

7 Registration etc of persons in possession of certain captive birds

[P1981/69/7]

- (1) If any person who is not for the time being registered in accordance with regulations made by the Department keeps or has in his possession or under his control any bird included in Schedule 4, he shall be guilty of an offence and shall be liable to a special penalty.
- (2) The power of the Department to make regulations under this section shall include power —

- (a) to impose requirements which must be satisfied before a person can be registered in accordance with the regulations;
 - (b) to require such birds to be ringed and marked in accordance with the regulations;
 - (c) to impose requirements as to the registration of such birds.
- (2A) In respect of the registration of a bird under this section, the Department shall charge such fees as are prescribed under the *Fees and Duties Act 1989*.¹⁵
- (3) If any person fails to comply with regulations under this section he shall be guilty of an offence.
- (4) If any person keeps or has in his possession or under his control any bird included in Schedule 4 —
 - (a) within five years of his having been convicted of an offence under this Part for which a special penalty is provided; or
 - (b) within three years of his having been convicted of any other offence under this Part so far as it relates to the protection of birds or other animals or any offence involving their ill-treatment,he shall be guilty of an offence.
- (5) If any person knowingly disposes of or offers to dispose of any bird included in Schedule 4 to another person —
 - (a) within five years of the conviction of that other person for such an offence as is mentioned in subsection (4)(a);
 - (b) within three years of the conviction of that other person for such an offence as is mentioned in subsection (4)(b),he shall be guilty of an offence.
- (5A) No account shall be taken for the purposes of subsections (4) and (5) of any conviction which has become spent for the purposes of the *Rehabilitation of Offenders Act 2001*.¹⁶
- (6) Any person authorised in writing by the Department may, at any reasonable time and (if required to do so) upon producing evidence that he is authorised, enter and inspect any premises where any birds included in Schedule 4 are kept for the purpose of ascertaining whether an offence under this section is being, or has been, committed on those premises.
- (7) Any person who intentionally obstructs a person acting in the exercise of the power conferred by subsection (6) shall be guilty of an offence.
- (8) Admission to a dwelling shall not be demanded as of right unless a warrant has been issued by a justice of the peace under subsection (9).
- (9) If it is shown to the satisfaction of a justice of the peace on information on oath that there are reasonable grounds for entry of the dwelling, the

justice may by warrant under his hand authorise an authorised person to enter the dwelling.

- (10) A warrant issued under subsection (9) shall continue in force for 5 days.

8 Protection of captive birds

[P1981/69/8]

- (1) If any person keeps or confines any bird whatever in any cage or other receptacle which is not sufficient in height, length or breadth to permit the bird to stretch its wings freely, he shall be guilty of an offence and be liable to a special penalty.
- (2) Subsection (1) does not apply to poultry, or to the keeping or confining of any bird —
- (a) while that bird is in the course of conveyance, by whatever means;
 - (b) while that bird is being shown for the purposes of any public exhibition or competition if the time during which the bird is kept or confined for those purposes does not in the aggregate exceed 72 hours;
 - (c) while that bird is undergoing examination or treatment by a veterinary surgeon; or¹⁷
 - (d) while that bird is being kept in a birdbag or keeping cage as part of a ringing exercise carried out under and in accordance with the terms of a licence granted under section 16(1)(b).
- (3) Every person who —
- (a) promotes, arranges, conducts, assists in, receives money for, or takes part in, any event whatever at or in the course of which captive birds are liberated by hand or by any other means whatever —
 - (i) for the purpose of being shot; or
 - (ii) for the purpose of being hunted by trained birds of prey, immediately after their liberation; or
 - (b) being the owner or occupier of any land, permits that land to be used for the purposes of such an event,
- shall be guilty of an offence and be liable to a special penalty.

*Protection of other animals***9 Protection of certain wild animals**

[P1981/69/9]

- (1) Subject to the provisions of this Part, if any person intentionally or recklessly kills, injures or takes any wild animal included in Schedule 5 without reasonable excuse, he shall be guilty of an offence.¹⁸
- (2) Subject to the provisions of this Part, if any person has in his possession or control any live or dead wild animal included in Schedule 5 or any part of, or anything derived from, such an animal, he shall be guilty of an offence.
- (3) A person shall not be guilty of an offence under subsection (2) if he shows that —
 - (a) the animal had not been killed or taken, or had been killed or taken otherwise than in contravention of the relevant provisions; or
 - (b) the animal or other thing in his possession or control had been sold (whether to him or any other person) otherwise than in contravention of those provisions;and in this subsection “the relevant provisions” means the provisions of this Part.
- (4) Subject to the provisions of this Part, if any person intentionally or recklessly —
 - (a) damages or destroys, or obstructs access to, any structure or place which any wild animal included in Schedule 5 uses for shelter or protection; or
 - (b) disturbs any such animal while it is occupying a structure or place which it uses for that purpose,he shall be guilty of an offence.¹⁹
- (4A) Subject to the provisions of this Part, any person who intentionally or recklessly disturbs any wild animal included in Schedule 5 or —
 - (a) a dolphin or whale (cetacean);
 - (b) a basking shark (*cetorhinus maximus*);
 - (c) a seal (all species) (pinnepedia); or
 - (d) a turtle (marine) (all species) (dermochelydiae and cheloniidae),shall be guilty of an offence.²⁰
- (5) Subject to the provisions of this Part, if any person —
 - (a) sells, offers or exposes for sale, or has in his possession or transports for the purpose of sale, any live or dead wild animal

included in Schedule 5, or any part of, or anything derived from, such an animal; or

- (b) publishes or causes to be published any advertisement likely to be understood as conveying that he buys or sells, or intends to buy or sell, any of those things,

he shall be guilty of an offence.

- (6) In any proceedings for an offence under subsection (1), (2) or (5)(a), the animal in question shall be presumed to have been a wild animal unless the contrary is shown.

10 Exceptions to s 9

[P1981/69/10]

- (1) Nothing in section 9 shall make unlawful —
 - (a) anything done in pursuance of a requirement by the Department under section 1 of the *Prevention of Damage by Agricultural Pests Act 1956*;
 - (b) anything done under section 3 of the *Forestry Act 1984*; or
 - (c) anything done under, or in pursuance of an order made under, the Diseases of Animals (Prevention) Acts 1948 to 1975.
- (2) Nothing in section 9(4) shall make unlawful anything done within a dwelling-house.
- (3) Notwithstanding section 9, a person shall not be guilty of an offence by reason of —
 - (a) the taking of any such animal if he shows that the animal had been disabled otherwise than by his unlawful act and was taken solely for the purpose of tending it and releasing it when no longer disabled;
 - (b) the killing of any such animal if he shows that the animal had been so seriously disabled otherwise than by his unlawful act that there was no reasonable chance of its recovering; or
 - (c) any act made unlawful by that section if he shows that the act was the incidental result of a lawful operation and could not reasonably have been avoided.
- (4) Notwithstanding section 9, an authorised person shall not be guilty of an offence by reason of the killing or injuring of a wild animal included in Schedule 5 if he shows that his action was necessary for the purpose of preventing serious damage to livestock, foodstuffs for livestock, crops, vegetables, fruit, growing timber or any other form of property or to fisheries.
- (5) A person shall not be entitled to rely on the defence provided by subsection (2) or (3)(c) as respects anything done in relation to a bat

otherwise than in the living area of a dwelling house unless he had notified the Department of the proposed action or operation and allowed it a reasonable time to advise him as to whether it should be carried out and, if so, the method to be used.

- (6) An authorised person shall not be entitled to rely on the defence provided by subsection (4) as respects any action taken at any time if it had become apparent, before that time, that the action would prove necessary for the purpose mentioned in that subsection and either —
- (a) a licence under section 16 authorising that action had not been applied for as soon as reasonably practicable after that fact had become apparent; or
 - (b) an application for such a licence had been determined.

11 Prohibition of certain methods of killing or taking wild animals

[P1981/69/11]

- (1) Subject to the provisions of this Part, if any person —
- (a) sets in position any self-locking snare which is of such a nature and so placed as to be calculated to cause bodily injury to any wild animal coming into contact therewith;
 - (b) uses for the purpose of killing or taking any wild animal any self-locking snare, whether or not of such a nature or so placed as aforesaid, any bow, crossbow or caterpult, or any explosive other than ammunition for a firearm; or
 - (c) uses as a decoy, for the purpose of killing or taking any wild animal, any live mammal or bird whatever; or
 - (d) knowingly causes or permits to be done an act which is mentioned in paragraphs (a) or (b)²¹

he shall be guilty of an offence.

- (2) Subject to the provisions of this Part if any person —
- (a) sets in position any of the following articles, being an article which is of such a nature and so placed as to be calculated to cause bodily injury to any wild animal included in Schedule 6 which comes into contact therewith, that is to say, any trap or snare, any electrical device for killing or stunning, or any poisonous, poisoned or stupefying substance;
 - (b) uses for the purpose of killing or taking any such wild animal any such article as aforesaid, whether or not of such a nature and so placed as aforesaid, or any net;
 - (c) uses for the purpose of killing or taking any such wild animal —
 - (i) any automatic or semi-automatic weapon;

- (ii) any device for illuminating a target or sighting device for night shooting;
- (iii) any form of artificial light or any mirror or other dazzling device; or
- (iv) any gas or smoke not falling within paragraphs (a) and (b);
- (d) uses as a decoy, for the purpose of killing or taking any such wild animal, any sound recording; or
- (e) uses any mechanically propelled vehicle in immediate pursuit of any such wild animal for the purpose of driving, killing or taking that animal; or
- (f) knowingly causes or permits to be done an act which is mentioned in paragraphs (a) to (e),²²

he shall be guilty of an offence.

- (3) Subject to the provisions of this Part, if any person —
 - (a) sets in position or knowingly causes or permits to be set in position any snare which is of such a nature and so placed as to be calculated to cause bodily injury to any wild animal coming into contact therewith; and²³
 - (b) while the snare remains in position fails without reasonable excuse, to inspect it, or cause it to be inspected, at least once every day

he shall be guilty of an offence.

- (4) The Department may, for the purpose of complying with an international obligation, by order, either generally or in relation to any kind of wild animal specified in the order, amend subsection (1) or (2) by adding any method of killing or taking wild animals or by omitting any such method as is mentioned in that subsection.
- (5) In any proceedings for an offence under subsection (1)(b) or (c) or (2) (b), (c), (d) or (e) and in any proceedings for an offence under subsection (1)(d) or (2)(f) relating to an act which is mentioned in any of those paragraphs, the animal in question shall be presumed to have been a wild animal unless the contrary is shown.²⁴
- (6) In any proceedings for an offence under subsection (2)(a) or under subsection (2)(f) relating to an act which is mentioned in subsection (2)(a) it shall be a defence to show that the article was set in position by the accused for the purpose of killing or taking, in the interests of public health, agriculture, forestry, fisheries or nature conservation, any wild animals which could be lawfully killed or taken by those means and that he took or caused to be taken all reasonable precautions to prevent injury thereby to any wild animals included in Schedule 6.²⁵

*Protection of plants***12 Protection of wild plants**

[P1981/69/13]

- (1) Subject to the provisions of this Part, if any person —
- (a) intentionally or recklessly picks, uproots or destroys any wild plant included in Schedule 7; or²⁶
 - (b) not being an authorised person, intentionally or recklessly uproots any wild plant not included in that Schedule,²⁷
- he shall be guilty of an offence.
- (2) Subject to the provisions of this Part, if any person —
- (a) sells, offers or exposes for sale, or has in his possession or transports for the purpose of sale, any live or dead wild plant included in Schedule 7, or any part of, or anything derived from, such a plant; or
 - (b) publishes or causes to be published any advertisement likely to be understood as conveying that he buys or sells, or intends to buy or sell, any of those things,
- he shall be guilty of an offence.
- (3) Notwithstanding anything in subsection (1), a person shall not be guilty of an offence by reason of any act made unlawful by that subsection if he shows that the act was an incidental result of a lawful operation and could not reasonably have been avoided.
- (4) In any proceedings for an offence under subsection (2)(a), the plant in question shall be presumed to have been a wild plant unless the contrary is shown.
- (5) Any person authorised in writing by the Department may, at any reasonable time and (if required to do so) upon producing evidence that he is authorised, enter any land for the purpose of ascertaining whether an offence under subsection (1) or (2) is being, or has been, committed on that land; but nothing in this subsection shall authorise any person to enter a dwelling.
- (6) Any person who intentionally obstructs a person acting in the exercise of the power conferred by subsection (5) shall be guilty of an offence.

*Miscellaneous***13 Areas of special protection: animals and plants**

- (1) The Department may by order make provision with respect to any area specified in the order providing for all or any of the following matters, that is to say —
- (a) that any person who, within that area or any part of it specified in the order, at any time or during any period so specified intentionally —
 - (i) kills, injures or takes any wild animal;
 - (ii) takes, damages or destroys any structure or place which such animal uses for shelter or protection while that structure is in use or being built or while that place is in use;
 - (iii) takes, damages or destroys anything which conceals or protects any such structure or place;
 - (iv) takes or destroys the egg, larva, pupa or any such immature state of such an animal;
 - (v) disturbs such animal while it is building a structure for shelter or protection or while it is in, on or near such structure containing the egg, larva, pupa or any such immature stage of such an animal;
 - (vi) disturbs the dependent young of such animal; or
 - (vii) picks, removes, uproots or destroys any wild plant;shall be guilty of an offence under this section;
 - (b) that any person who, except as may be provided in the order, enters into that area or any part of it specified in the order at any time or during any period so specified shall be guilty of an offence under this section;
 - (c) that where any offence under this Part, or any such offence under this Part as may be specified in the order, is committed within that area, the offender shall be liable to a special penalty.
- (2) If any authorised person suspects with reasonable cause that any person is committing or has committed an offence under an order made under this section in relation to any area, he may require that person —
- (a) to give his full name and address; and
 - (b) to quit that area forthwith;
- and any person who fails to comply with a requirement under this paragraph shall be guilty of an offence.
- (3) The making of any order under this section with respect to any area shall not affect the exercise by any person of any right vested in him, whether

as owner, lessee or occupier of any land in that area or by virtue of a licence or agreement.

- (4) Before making any order under this section the Department —
 - (a) shall consult with any local authority within whose district the area with respect to which the order is to be made or any part thereof is situated; and
 - (b) shall give particulars of the intended order either by notice in writing to every owner and every occupier of any land included in the area with respect to which the order is to be made or, where the giving of such a notice is in the Department's opinion impracticable, by advertisement in a newspaper published and circulating in the Island.
- (5) The Department shall not make an order under this section unless —
 - (a) all the owners and occupiers aforesaid have consented thereto; or
 - (b) no objections thereto have been made by any of those owners or occupiers before the expiration of a period of 3 months from the date of the giving of the notice or the publication of the advertisement; or
 - (c) any such objections so made have been withdrawn.

14 Introduction of new species etc

[P1981/69/14]

- (1) Subject to the provisions of this Part, if any person releases or allows to escape into the wild any animal which —
 - (a) is of a kind which is not ordinarily resident in and is not a regular visitor to the Island in a wild state; or
 - (b) is included in Part I of Schedule 8,he shall be guilty of an offence.
- (2) Subject to the provisions of this Part, if any person plants or otherwise causes to grow in the wild any plant which is included in Part II of Schedule 8 he shall be guilty of an offence.
- (3) Subject to subsection (4), it shall be a defence to a charge of committing an offence under subsection (1) or (2) to prove that the accused took all reasonable steps and exercised all due diligence to avoid committing the offence.
- (4) Where the defence provided by subsection (3) involves an allegation that the commission of the offence was due to the act or default of another person, the person charged shall not, without leave of the court, be entitled to rely on the defence unless, within a period ending seven clear days before the hearing, he has served on the prosecutor a notice giving

such information identifying or assisting in the identification of the other person as was then in his possession.

- (5) Any person authorised in writing by the Department may, at any reasonable time and (if required to do so) upon producing evidence that he is authorised, enter any land for the purpose of ascertaining whether an offence under subsection (1) or (2) is being, or has been, committed on that land; but nothing in this subsection shall authorise any person to enter a dwelling.
- (6) Any person who intentionally obstructs a person acting in the exercise of the power conferred by subsection (5) shall be guilty of an offence.

15 [Repealed]²⁸

Supplemental

16 Power to grant licences

[P1981/69/16]

- (1) Sections 1, 5, 6(3) and (4), 7 and 8 do not apply to anything done —
 - (a) for scientific or educational purposes;
 - (b) for the purpose of ringing or marking, or examining any ring or mark on, wild birds;
 - (c) for the purpose of conserving wild birds;
 - (d) for the purpose of protecting any collection of wild birds;
 - (e) for the purposes of falconry or aviculture;
 - (f) for the purposes of any public exhibition or competition;
 - (g) for the purpose of taxidermy;
 - (h) for the purpose of photography;
 - (i) for the purposes of preserving public health or public or air safety;
 - (j) for the purpose of preventing the spread of disease; or
 - (k) for the purposes of preventing serious damage to livestock, foodstuffs for livestock, crops, vegetables, fruit, growing timber or fisheries,

if it is done under and in accordance with the terms of a licence granted by the Department.²⁹

- (1A) Orders under section 3 do not apply to anything done by an owner or occupier, or any person authorised by the owner or occupier, of the land on which it is done, if it is done under and in accordance with the terms of a licence granted by the Department.³⁰

- (2) Section 1 and orders under section 3 do not apply to anything done for the purpose of providing food for human consumption in relation to a gull's egg if it is done under and in accordance with the terms of a licence granted by the Department.
- (3) Sections 9(1), (2) and (4), 11(1) and (2) and 12(1) and orders under section 13 do not apply to anything done —
- (a) for scientific or educational purposes;
 - (b) for the purpose of ringing or marking, or examining any ring or mark on, wild animals;
 - (c) for the purpose of conserving wild animals or wild plants or introducing them to particular areas;
 - (d) for the purpose of protecting any zoological or botanical collection;
 - (e) for the purpose of photography;
 - (f) for the purposes of preserving public health or public safety;
 - (g) for the purpose of preventing the spread of disease; or
 - (h) for the purposes of preventing serious damage to livestock, foodstuffs for livestock, crops, vegetables, fruit, growing timber or any other form of property or to fisheries,
- if it is done under and in accordance with the terms of a licence granted by the Department.
- (4) The following provisions, namely —
- (a) section 6(1) and (2);
 - (b) sections 9(5) and 12(2); and
 - (c) section 14,
- do not apply to anything done under and in accordance with the terms of a licence granted by the Department.
- (5) Subject to subsection (6), a licence under the foregoing provisions of this section —
- (a) may be, to any degree, general or specific;
 - (b) may be granted either to persons of a class or to a particular person;
 - (c) may be subject to compliance with any specified conditions;
 - (d) may be modified or revoked at any time by the Department; and
 - (e) subject to paragraph (d), shall be valid for the period stated in the licence;
- and the Department may charge therefor such sum (if any) as it may determine.

- (6) A licence under subsection (1), (1A), (2) or (3) which authorises any person to kill wild birds or wild animals —
 - (a) shall specify the area within which, and the methods by which the wild birds or wild animals may be killed; and
 - (b) subject to subsection (5)(d), shall be valid for the period, not exceeding two years, stated in the licence.³¹
- (7) It shall be a defence in proceedings for an offence under section 8(b) of the *Cruelty to Animals Act 1925* (which restrict the placing on land of poison and poisonous substances) to show that —
 - (a) the act alleged to constitute the offence was done under and in accordance with the terms of a licence issued under subsection (1) or (3); and
 - (b) any conditions specified in the licence were complied with.
- (8) For the purposes of a licence granted under this section, the definition of a class of persons may be framed by reference to any circumstances whatever including, in particular, their being authorised by any other person.
- (9) The Department —
 - (a) shall from time to time consult with the Wildlife Committee as to the exercise of its functions under this section; and
 - (b) shall not grant a licence of any description unless it has been advised by the Committee as to the circumstances in which, in their opinion, licences of that description should be granted.

17 False statements made for obtaining registration or licence etc

[P1981/69/17]

A person who, for the purposes of obtaining, whether for himself or another, a registration in accordance with regulations made under section 6 (2) or 7(1) or the grant of a licence under section 16 —

- (a) makes a statement or representation, or furnishes a document or information, which he knows to be false in a material particular; or
 - (b) recklessly makes a statement or representation, or furnishes a document or information, which is false in a material particular,
- shall be guilty of an offence.

18 Prohibition of hunting with dogs

- (1) If any person uses or causes to be used any dog or pack of dogs for the purpose of coursing, hunting or otherwise endeavouring to take or kill any bird or other animal he shall be guilty of an offence.

- (2) In any proceedings for an offence under subsection (1) it shall be a defence for an authorised person to show that the dog or dogs were being used for the purpose of hunting, taking or killing rabbits or rats.
- (3) In this section, the words “bird or other animal” do not include game within the meaning of the *Game Act 1882*.
- (4) This section is without prejudice to section 3 of the *Game (Hares) Act 1971* (prohibition of competitive hare coursing).

19 Possession of implements etc

[P1981/69/18]

Any person who for the purposes of committing an offence under the foregoing provisions of this Part, has in his possession anything capable of being used for committing the offence shall be guilty of an offence and shall be punishable in like manner as for the said offence.

20 Enforcement

[P1981/69/19]

- (1) If a constable has reasonable cause to suspect that any person is committing or has committed an offence under this Part, the constable may without warrant —
 - (a) stop and search that person if the constable suspects with reasonable cause that evidence of the commission of the offence is to be found on that person;
 - (b) search or examine anything which that person may then be using or have in his possession if the constable suspects with reasonable cause that evidence of the commission of the offence is to be found on that thing;
 - (c) [Repealed]³²
 - (d) seize and detain for the purposes of proceedings under this Part anything which may be evidence of the commission of the offence or may be liable to be forfeited under section 22.
- (2) If a constable has reasonable cause to suspect that any person is committing an offence under this Part, he may, for the purpose of exercising the powers conferred by subsection (1) or arresting a person, under section 28 of the *Police Powers and Procedures Act 1998*, for such an offence, enter any land other than a dwelling-house.³³
- (3) If a justice of the peace is satisfied by information on oath that there are reasonable grounds for suspecting that —
 - (a) an offence under section 1, 3, 5, 7 or 8 in respect of which this Part or any order made under it provides for a special penalty; or
 - (b) an offence under section 6, 9, 11(1) or (2), 12, 13 or 14,

has been committed and that evidence of the offence may be found on any premises, he may grant a warrant to any constable (with or without other persons) to enter upon and search those premises for the purpose of obtaining that evidence.

21 Summary prosecutions

[P1981/69/20]

- (1) This section applies to —
 - (a) any offence under section 1(1) or 3(1) involving the killing or taking of any wild bird or the taking of an egg of such a bird;
 - (b) any offence under section 9(1) or 13(1) involving the killing or taking of any wild animal; and
 - (c) any offence under section 12(1) or 13(1) involving the picking, uprooting or destruction of any wild plant.
- (2) Summary proceedings for an offence to which this section applies may be brought within a period of 6 months from the date on which evidence sufficient in the opinion of the prosecutor to warrant the proceedings came to his knowledge; but no such proceedings shall be brought by virtue of this section more than two years after the commission of the offence.
- (3) For the purpose of this section a certificate signed by or on behalf of the prosecutor and stating the date on which such evidence as aforesaid came to his knowledge shall be conclusive evidence of that fact; and a certificate stating that matter and purporting to be so signed shall be deemed to be so signed unless the contrary is proved.

22 Penalties, forfeitures etc

[P1981/69/21]

- (1) Subject to subsection (6), a person guilty of an offence under section 1, 3, 5, 6, 7, 8 or 13 shall be liable on summary conviction —
 - (a) in a case where this Part or any order made under it provides that he shall be liable to a special penalty, to a fine not exceeding £5,000;
 - (b) in any other case, to a fine not exceeding £2,500.³⁴
- (2) Subject to subsection (6), a person guilty of an offence under section 9 or 11(1) or (2) shall be liable on summary conviction to a fine not exceeding £5,000.³⁵
- (3) Subject to subsection (6), a person guilty of an offence under section 11(3), 12 or 17 shall be liable on summary conviction to a fine not exceeding £2,500.³⁶

- (4) A person guilty of an offence under section 18 shall be liable on summary conviction to a fine not exceeding £5,000.
- (5) A person guilty of an offence under section 14 shall be liable —
 - (a) on summary conviction, to a fine not exceeding £5,000;
 - (b) on conviction on information, to a fine.
- (6) Where an offence to which subsection (1), (2) or (3) applies was committed in respect of more than one bird, nest, egg, other animal, plant or other thing, the maximum fine which may be imposed under that subsection shall be determined as if the person convicted had been convicted of a separate offence in respect of each bird, nest, egg, animal, plant or thing.
- (7) The court by which any person is convicted of an offence under this Part —
 - (a) shall order the forfeiture of any bird, nest, egg, other animal, plant or other thing in respect of which the offence was committed; and
 - (b) may order the forfeiture of any vehicle, animal, weapon or other thing which was used to commit the offence and, in the case of an offence under section 14, any animal or plant which is of the same kind as that in respect of which the offence was committed and was found in his possession.

23 Power to vary Schedules

[P1981/69/22]

- (1) The Department may by order add any bird to, or remove any bird from, any of or any Part of Schedules 1 to 4.
- (2) An order under subsection (1) adding any bird to Part II of Schedule 1 or Part I of Schedule 2 may prescribe a close season in the case of that bird for the purposes of sections 1 and 2; and any close season so prescribed shall commence on a date not later than 21st February and end on a date not earlier than 31st August.
- (3) The Department may, on a representation made to it by the Wildlife Committee, by order —
 - (a) add to Schedule 5 or Schedule 7 any animal or plant which, in its opinion, is in danger of extinction in the Island or is likely to become so endangered unless conservation measures are taken; and
 - (b) remove from Schedule 5 or Schedule 7 any animal or plant which, in its opinion, is no longer so endangered or likely to become so endangered.
- (4) The Department may, for the purpose of complying with an international obligation, by order —

- (a) add any animals to, or remove any animals from, Schedule 5 or Schedule 6; and
 - (b) add any plants to, or remove any plants from, Schedule 7.
- (5) The Department may by order —
 - (a) add any animals to, or remove any animals from, Part I of Schedule 8; and
 - (b) add any plants to, or remove any plants from, Part II of that Schedule.

24 Advisory bodies and their functions

[P1981/69/23]

- (1) [Repealed]³⁷
- (2) The Wildlife Committee shall exercise the functions conferred on it by this Act.
- (3) It shall be the duty of the Wildlife Committee to advise the Department on any question which the Department may refer to it or on which it considers it should offer its advice —
 - (a) in connection with the administration of this Act; or
 - (b) otherwise in connection with the protection of birds or other animals or plants.

25 Regulations, orders, notices etc

[P1981/69/26]

- (1) Any regulations, byelaws or order under a provision of this Act shall be laid before Tynwald as soon as may be after the making thereof, and if Tynwald at the sitting before which such regulations, byelaws or order are so laid or at the next sitting resolves that the regulations, byelaws or order shall be annulled, the regulations, byelaws or order shall cease to have effect.
- (2) Before making any order under this Part, the Department —
 - (a) except in the case of an order under section 2(6), shall give to any local authority affected and, except in the case of an order under section 3, any other person affected, by such means as it may think appropriate, an opportunity to submit objections or representations with respect to the subject matter of the order;
 - (b) shall consult with the Wildlife Committee as to whether the order should be made; and
 - (c) may, if it thinks fit, cause a public inquiry to be held.
- (3) The Department shall give consideration to any proposals for the making by it of an order under this Part with respect to any area which may be submitted to it by a local authority whose area includes that area.

26 Interpretation of Part I

[P1981/69/27]

(1) In this Part, unless the context otherwise requires —

“**advertisement**” includes a catalogue, a circular and a price list;

“**authorised person**” means —

- (a) the owner or occupier, or any person authorised by the owner or occupier, of the land on which the action authorised is taken;
- (b) as respects anything done in relation to wild birds, any person authorised in writing by any of the following bodies, that is to say, the Department or the Manx Utilities Authority;³⁸

so, however, that the authorisation of any person for the purposes of this definition shall not confer any right of entry upon any land;

“**automatic weapon**” and “**semi-automatic weapon**” do not include any weapon the magazine of which is incapable of holding more than two rounds;

“**aviculture**” means the breeding and rearing of birds in captivity;

“**destroy**”, in relation to an egg, includes doing anything to the egg which is calculated to prevent it from hatching, and “**destruction**” shall be construed accordingly;

“**domestic duck**” means any domestic form of duck;

“**domestic goose**” means any domestic form of goose;

“**firearm**” has the same meaning as in the *Firearms Act 1947*;

“**game bird**” means any pheasant, partridge, grouse (or moor game);

“**livestock**” includes any animal which is kept —

- (a) for the provision of food, wool, skins or fur;
- (b) for the purpose of its use in the carrying on of any agricultural activity; or
- (c) for the provision or improvement of shooting or fishing;

“**occupier**”, in relation to any land other than the foreshore, includes any person having any right of hunting, shooting, fishing or taking game or fish;

“**pick**”, in relation to a plant, means gather or pluck any part of the plant without uprooting it;

“**poultry**” means domestic fowls, geese, ducks, guinea-fowls, pigeons and quails, and turkeys;

“**sale**” includes hire, barter and exchange and cognate expressions shall be construed accordingly;

“**uproot**”, in relation to a plant, means dig up or otherwise remove the plant from the land on which it is growing;

“**vehicle**” includes aircraft, hovercraft and boat;

“**wild animal**” means any animal (other than a bird) which is or (before it was killed or taken) was living wild;

“**wild bird**” means any bird of a kind which is ordinarily resident in or is a visitor to the Island in a wild state but does not include poultry or, except in sections 5 and 16, any game bird;

“**Wildlife Committee**” has the meaning given by section 24;

“**wild plant**” means any plant which is or (before it was picked, uprooted or destroyed) was growing wild and is of a kind which ordinarily grows in the Island in a wild state.

- (2) A bird shall not be treated as bred in captivity for the purposes of this Part unless its parents were lawfully in captivity when the egg was laid.
- (3) Any reference in this Part to an animal of any kind includes, unless the context otherwise requires, a reference to an egg, larva, pupa, or other immature stage of an animal of that kind.
- (4) This Part extends to the territorial waters adjacent to the Island.

26A Exemption for certain flood risk management functions

- (1) This section applies for the performance of a function (the “activity”) by the Manx Utilities Authority (“the Authority”) under the *Flood Risk Management Act 2013*.³⁹
- (2) The Authority does not commit an offence against, or contravene, this Part if —
 - (a) the activity is carried out with the Department’s consent; or
 - (b) in an emergency and the Authority informs the Department of the carrying out of the activity as soon as practicable after it happened.
- (3) Reasonable conditions may be imposed on the consent, but it must not be unreasonably withheld or delayed.
- (4) Any question as to the following must be referred to an arbitrator to be agreed between the parties or, failing agreement, appointed by the Governor in Council —
 - (a) whether the carrying out of the activity constitutes, or will constitute, an offence against, or a contravention of, this Part;
 - (b) whether consent is being unreasonably withheld or delayed; or
 - (c) whether a condition imposed on any consent given is reasonable.⁴⁰

PART II – NATURE CONSERVATION

27 Areas of special scientific interest

[P1981/69/28]

- (1) Where the Department, after consultation with the Wildlife Committee is of the opinion that any area of land is of special interest by reason of any of its flora, fauna, or geological or physiographical features, it may notify that fact to —
 - (a) [Repealed]⁴¹
 - (b) to every owner and occupier of any of that land.
- (2) A notification under subsection (1) shall have effect immediately it is served on all the persons mentioned in subsection (1) and shall continue in force until rescinded.
- (3) A notification under subsection (1) shall specify —
 - (a) the flora, fauna or geological or physiographical features by reason of which the land is of special interest;⁴²
 - (b) any operations or activity of a type appearing to the Department to be likely to damage that flora or fauna or those features; and
 - (c) the time (not being less than 3 months from the date of the notification) within which and the manner in which representations or objections with respect to the notification may be made.
- (4) If any representations or objections are received by the Department from any of the persons mentioned in subsection (1), the Department shall consider those representations or objections and, within 2 months of the expiry of the period referred to in subsection (3)(c), shall either confirm, vary or rescind the notification and shall notify the persons mentioned in subsection (1) of its decision and the reasons for it.
- (5) The owner or occupier of any land which is the subject of a notification under subsection (1) shall not carry out, nor cause or permit to be carried out on that land any operation or activity specified in the notification unless —
 - (a) one of them has given the Department written notice by recorded delivery of a proposal to carry out the operation or activity specifying its nature and the land on which it is proposed to carry it out; and
 - (b) one of the following conditions is fulfilled —
 - (i) that the operation or activity is carried out with the Department's written consent;

- (ii) that the operation or activity is carried out in accordance with the terms of an agreement entered into under section 30;
 - (iii) subject to subsection (6), that a period of 4 months has expired from the giving of notice under paragraph (a).
- (6) If, before the expiration of the said period of 4 months the Department offers to enter into an agreement under section 30, sub-paragraph (iii) of subsection (5)(b) shall have effect as if for the period of 4 months there were substituted a period of 12 months and if, within the latter period, the offeree does not enter into such an agreement, the matter shall be referred to an arbitrator to be agreed between the offeree and the Department, or in the absence of such agreement by the Clerk of the Rolls, and the findings of the arbitrator shall be binding on both parties.⁴³
- (7) An agreement offered by the Department in accordance with subsection (6) shall provide for the making by the Department of payments determined in accordance with financial guidelines which shall be prepared by the Department after consultation with bodies appearing to it to be representative of the interests likely to be affected.
- (8) Guidelines prepared under subsection (7) shall not have effect unless approved by Tynwald.
- (9) A person who, without reasonable excuse, contravenes subsection (5) shall be liable on summary conviction to a fine not exceeding £5,000 or on conviction on information to a fine.
- (10) It is a reasonable excuse in any event for a person to carry out an operation if the operation was an emergency operation particulars of which (including details of the emergency) were notified to the Department as soon as practicable after the commencement of the operation.
- (11) Proceedings for an offence under subsection (9) shall not be taken without the consent of the Attorney General.
- (12) In this section “operation” includes the removal or disturbance of any rock, plant, animal or other thing.
- (13) In this section “land” includes land covered (whether constantly or not) by water.⁴⁴

28 Restoration where s 27 contravened

[P1981/69/31]

- (1) Where the operation or activity in respect of which a person is convicted of an offence under section 27 has destroyed or damaged any of the flora, fauna, or geological or physiographical features by reason of which the land on which it was carried out is of special interest, the court of summary jurisdiction by which he is convicted, in addition to dealing

with him in any other way, may make an order requiring him to carry out, within such period as may be specified in the order, such operations for the purpose of restoring the land to its former condition as may be so specified.

- (2) Where an order is made under this section by a court of summary jurisdiction, the period specified in the order shall not begin to run —
 - (a) in any case until the expiration of the period for the time being prescribed by law for the giving of notice of appeal against a decision of a court of summary jurisdiction;
 - (b) where notice of appeal is given within the period so prescribed, until determination of the appeal.
- (3) At any time before an order under this section has been complied with or fully complied with, a court of summary jurisdiction may, on the application of the person against whom it was made, discharge or vary the order if it appears to the court that a change in circumstances has made compliance or full compliance with the order impracticable or unnecessary.
- (4) If, within the period so specified in an order under this section, the person against whom it was made fails, without reasonable excuse, to comply with it, he shall be liable on summary conviction —
 - (a) to a fine not exceeding £5,000; and
 - (b) in the case of a continuing offence, to a further fine not exceeding £100 for each day during which the offence continues after conviction.
- (5) If, within the period specified in an order under this section, any operations specified in the order have not been carried out, the Department may by its agents and employees enter the land and carry out those operations and recover from the person against whom the order was made any expenses reasonably incurred in doing so.

29 Compensation

- (1) Where notification is made under section 27, the Department may pay compensation to any person having at the time of the making of the notification an interest in land to which the notification relates who, on a claim made to the Department within the time and in the manner prescribed by regulations under this section, shows that the value of his interest is less than it would have been if the notification had not been made.⁴⁵
- (2) The amount of the compensation paid under subsection (1) shall be equal to the difference between the two values mentioned in that subsection.
- (3) For the purposes of subsection (1) —

- (a) an interest in land shall be valued as at the time when the notification was made;
 - (b) where a person, by reason of his having more than one interest in land, makes more than one claim under that subsection in respect of the same notification, his various interests shall be valued together.
- (4) For the purpose of assessing any compensation payable under this section, the rules set out in section 5 of the *Acquisition of Land Act 1984* (rules for assessing compensation) shall, so far as applicable and subject to any necessary modifications, have effect as they have effect for the purpose of assessing compensation for the compulsory acquisition of an interest in land.
- (5) No claim shall be made under this section in respect of any notification under section 27 unless the Department has made a decision under subsection (4) of that section and given notice in accordance with that subsection.
- (6) Except insofar as may be provided by regulations under this section, any question of disputed compensation under this section shall be referred to and determined by an arbitrator appointed under, and selected in accordance with, section 3 of that Act (tribunal for assessing compensation).
- (7) In relation to the determination of any such question, the provisions of sections 9 and 10 of that Act (procedure and costs) shall apply, subject to any necessary modifications and to the provisions of regulations under this section.

30 Management agreements with owners etc of land

- (1) The Department may for the purpose of —
 - (a) conserving or enhancing the natural beauty or amenity of any land;
 - (b) conserving the flora, fauna or geological or physiographical features of any land; or
 - (c) promoting its enjoyment by the public,make an agreement (in this section referred to as a “management agreement”) with any person having an interest in the land with respect to the management of the land during a specified term or without limitation of the duration of the agreement.
- (2) Without prejudice to the generality of subsection (1), a management agreement may —
 - (a) impose on the person having an interest in the land restrictions as respects the method of cultivating the land, its use for agricultural purposes or the exercise of rights over the land and may impose

- obligations on that person to carry out works or agricultural or forestry operations or do other things on the land; and
- (b) contain such further provisions (including provisions for the making of payments by either party to the other) as appear to the Department to be necessary or expedient for the purposes of the agreement.
- (3) The provisions of a management agreement with any person interested in the land shall, unless the agreement otherwise provides, be binding on persons deriving title under or from that person and be enforceable by the Department against those persons accordingly.
 - (4) Schedule 2 to the *Forestry Act 1984* (power for tenant for life and others to enter into forestry covenants) shall apply to management agreements as it applies to forestry covenants.
 - (5) The powers conferred by this section on the Department shall be in addition to and not in derogation of any powers of the Department conferred by or under any statutory provision.

31 National nature reserves

[P1981/69/35]

- (1) Where the Department is satisfied that any land which is being managed as a nature reserve by the owner or occupier is of national importance, it may, subject to section 37(3), declare that land to be a national nature reserve.
- (2) A declaration by the Department that any land is a national nature reserve shall be conclusive of the matters declared.
- (3) On the application of the owner and occupier concerned, the Department may, subject to section 37(3), as respects any land which is declared to be a national nature reserve under subsection (1), make byelaws for the protection of the reserve.
- (4) Without prejudice to the generality of subsection (3) and subject to subsection (6), byelaws under this section —
 - (a) may provide for prohibiting or restricting the entry into, or movement within, nature reserves of persons, vehicles, boats and animals;
 - (b) may prohibit or restrict the killing, taking, molesting or disturbance of living creatures of any description in a nature reserve, the taking, destruction or disturbance of eggs of any such creature, the taking of, or interference with, vegetation of any description in a nature reserve, or the doing of anything therein which will interfere with the soil or damage any object in the reserve;

- (c) may prohibit or restrict the shooting of birds or of birds of any description within such area surrounding or adjoining a nature reserve (whether the area be of land or of sea) as appears to the Department requisite for the protection of the reserve;
 - (d) may contain provisions prohibiting the depositing of rubbish and the leaving of litter in a nature reserve;
 - (e) may prohibit or restrict, or provide for prohibiting or restricting, the lighting of fires in a nature reserve, or the doing of anything likely to cause a fire in a nature reserve;
 - (f) may provide for the issue, on such terms and subject to such conditions as may be specified in the byelaws, of permits authorising entry into a nature reserve or the doing of anything therein which would otherwise be unlawful, whether under the byelaws or otherwise;
 - (g) may be made so as to relate either to the whole or to any part of the reserve or, in the case of byelaws made under paragraph (c) of any such surrounding or adjoining area as is mentioned in that paragraph, and may make different provisions for different parts thereof.
- (5) Before making any byelaws under subsection (4)(c), the Department shall consult with the occupier of the land to which the byelaws shall apply.
- (6) Byelaws under this section shall not interfere with the exercise by any person of a right vested in him as owner, lessee or occupier of land in a nature reserve, or with the exercise of any public right of way or of any functions of statutory undertakings, of the Department of Infrastructure, the Department of Environment, Food and Agriculture or any drainage authority.⁴⁶
- (7) In this section “nature reserve” means land managed for the purpose —
 - (a) of providing, under suitable conditions and control, special opportunities for the study of, and research into, matters relating to the fauna and flora of the Island and the physical conditions in which they live, and for the study of geological and physiographical features of special interest in the area, or
 - (b) of preserving flora, fauna or geological or physiographical features of special interest in the area,or for both those purposes.
- (8) If any person contravenes any byelaw made under this section he shall be guilty of an offence and, on summary conviction, shall be liable to a fine not exceeding £2,500.

32 Marine nature reserves

[P1981/69/36]

- (1) Where, in the case of any land covered (continuously or intermittently) by tidal waters or parts of the sea in or adjacent to the Island up to the seaward limits of territorial waters it appears to the Department expedient, on an application made by the Wildlife Committee that the land and waters covering it should be managed by the Department for the purpose of —

- (a) conserving marine flora or fauna or geological or physiographical features of special interest in the area; or
- (b) providing, under suitable conditions and control, special opportunities for the study of, and research into, matters relating to marine flora and fauna and the physical conditions in which they live, or for the study of geological and physiographical features of special interest in the area,

the Department may, subject to section 37(3), by order designate the area comprising that land and those waters as a marine nature reserve; and the Department shall manage any area so designated for either or both of those purposes.

- (2) An application for an order under this section shall be accompanied by a copy of the byelaws which, if an order is made, the Wildlife Committee recommend that the Department make under section 33 for the protection of the area specified in the application.
- (3) The powers exercisable by the Department for the purpose of managing an area designated as a marine nature reserve under this section shall include power to install markers indicating the existence and extent of the reserve.
- (4) Nothing in this section or in byelaws made under section 33 shall interfere with the exercise of any functions of a relevant authority, any functions conferred by or under an enactment (whenever passed) or any right of any person (whenever vested).
- (5) In this section “relevant authority” means the Manx Utilities Authority, the Department of Environment, Food and Agriculture, the Department of Infrastructure, a lighthouse authority or a drainage authority.⁴⁷

33 Byelaws for protection of marine nature reserves

[P1981/69/37]

- (1) The Department may, subject to section 37(3), make byelaws for the protection of any area designated as a marine nature reserve under section 32.
- (2) Without prejudice to the generality of subsection (1), byelaws made under this section as respects a marine nature reserve —

- (a) may provide for prohibiting or restricting, either absolutely or subject to any exceptions —
 - (i) the entry into, or movement within, the reserve of persons and vessels;
 - (ii) the killing, taking, destruction, molestation or disturbance of animals or plants of any description in the reserve, or the doing of anything therein which will interfere with the sea bed or damage or disturb any object in the reserve; or
 - (iii) the depositing of rubbish in the reserve;
 - (b) may provide for the issue, on such terms and subject to such conditions as may be specified in the byelaws, of permits authorising entry into the reserve or the doing of anything which would otherwise be unlawful under the byelaws; and
 - (c) may be so made as to apply either generally or with respect to particular parts of the reserve or particular times of the year.
- (3) Nothing in byelaws made under this section shall —
- (a) prohibit or restrict the exercise of any right of passage by a vessel other than a pleasure boat; or
 - (b) prohibit, except with respect to particular parts of the reserve at particular times of the year, the exercise of any such right by a pleasure boat.
- (4) Nothing in byelaws so made shall make unlawful —
- (a) anything done for the purpose of securing the safety of any vessel, or of preventing damage to any vessel or cargo, or of saving life;
 - (b) the discharge of any substance from a vessel; or
 - (c) anything done more than 30 metres below the sea bed.
- (5) If any person contravenes any byelaws made under this section he shall be guilty of an offence and, on summary conviction, shall be liable to a fine not exceeding £2,500.
- (6) Proceedings for an offence under byelaws made under this section shall not be taken without the consent of the Attorney General.
- (6A) A byelaw made under this section which relates to —
- (a) fishing for, or the taking or killing of, any fish, or
 - (b) the use or possession of any fishing engine,
- shall be treated as sea fishery legislation for the purposes of Part 8 of the *Fisheries Act 2012*.
- In this subsection “fish” and “fishing engine” have the same meanings as in that Act.⁴⁸

- (6B) An offence under a byelaw referred to in subsection (6A) shall be treated as an offence under the *Fisheries Act 2012* for the purposes of section 78 (forfeiture) of that Act.⁴⁹
- (7) In this section “vessel” includes a hovercraft and any aircraft capable of landing on water and “pleasure boat” shall be construed accordingly.
- (8) References in this section to animals or plants of any description include references to eggs, seeds, spores, larvae or other immature stages of animals or plants of that description.

PART III – MISCELLANEOUS AND GENERAL

34 Offences by bodies corporate etc

[P1981/69/69]

- (1) Where a body corporate is guilty of an offence under this Act and that offence is proved to have been committed with the consent or connivance of, or to be attributable to any neglect on the part of, any director, manager, secretary or other similar officer of the body corporate or any person who was purporting to act in any such capacity he, as well as the body corporate, shall be guilty of that offence and shall be liable to be proceeded against and punished accordingly.
- (2) Where the affairs of a body corporate are managed by its members subsection (1) shall apply in relation to the acts and defaults of a member in connection with his functions of management as if he were a director of the body corporate.

35 Publication of information

The Department shall publish information or arrange for the publication of information in such form and such manner as it considers appropriate with respect to —

- (a) the operation of this Act and any public document under it;
- (b) any matters relating to the functions of the Department under this Act;
- (c) any other matters about which it appears to it to be desirable to publish information for the guidance of persons it considers likely to be affected by this Act.

36 Duty to have regard to environment etc

- (1) In regard to any functions of the Department which may affect the physical environment, the Department shall, so far as may be consistent with the proper discharge of such functions, endeavour to secure a reasonable balance between —

- (a) the promotion and maintenance of a stable and efficient agricultural industry; and
 - (b) the conservation and enhancement of the natural beauty and amenity of the countryside, the protection of wildlife habitat, and the conservation of flora and fauna and geological or physiographical features of interest.
- (2) Without prejudice to subsection (1), in the exercise of any functions which may affect the physical environment, a department, statutory board or local authority shall, so far as may be consistent with the proper discharge of those functions, have regard to the matters specified in subsection (1)(b).

37 Financial provisions

[P1981/69/70]

- (1) There shall be defrayed out of money provided by Tynwald any expenses incurred by the Department under this Act.
- (2) Any sums received by the Department under this Act shall form part of the General Revenue of the Island.
- (3) The Department shall obtain the concurrence of the Treasury before making any order, bye-laws or declaration under sections 31, 32 and 33.

38 General interpretation

[P1981/69/71]

In this Act —

“**the commencement date**”, in relation to any provision of this Act, means the date of the coming into force of that provision;

“**the Department**” has the meaning given by section 2(3);

“**modifications**” includes additions, alterations and omissions and cognate expressions shall be construed accordingly;

“**the Wildlife Committee**” means the Wildlife Committee established under section 1(2) of the *Endangered Species Act 2010*.⁵⁰

39 Consequential amendments

- (1) [Amends section 2 and substitutes section 9 of the *Game Act 1882*.]
- (2) [Amends section 9 of the *Curraghs Acquisition Act 1963*.]
- (3) [Amends sections 2, 4 and 7 of the *Wild Animals (Restriction on Importation, Etc.) Act 1980*.]

40 Repeals

The enactments mentioned in Schedule 10 are repealed to the extent specified in the third column of that Schedule.

41 Short title and commencement

- (1) This Act may be cited as the Wildlife Act 1990.
- (2) This Act shall come into operation on such day as the Department may by order appoint and different days may be appointed under this subsection for different provisions, different purposes or different areas.
- (3) An order under subsection (2) may make such transitional provision as appears to the Department to be necessary or expedient in connection with the provisions thereby brought into force.⁵¹

SCHEDULE 1

BIRDS WHICH ARE PROTECTED BY SPECIAL PENALTIES

Sections 1, 2, 4, 6 and 23[Sch 1 substituted by SD129/04.]

PART I – AT ALL TIMES

<i>Common name</i>	<i>Scientific name</i>
Avocet	Recurvirostra avosetta
Bittern	Botaurus stellaris
Bullfinch	Pyrrhula pyrrhula
Bunting, Corn	Miliaria calandra
Bunting, Reed	Emberiza schoeniclus
Buzzard	Buteo buteo
Buzzard, Honey	Pernis apivorus
Buzzard, Rough-legged	Buteo lagopus
Chough	Pyrrhocorax pyrrhocorax
Corncrake	Crex crex
Crake, Spotted	Porzana porzana
Crane, Common	Grus grus
Crossbills (all species)	Loxia
Curlew	Numenius arquata
Dipper	Cinclus cinclus
Dove, Turtle	Streptopelia turtur
Eagle, Golden	Aquila chrysaetos
Eagle, White-tailed	Haliaeetus albicilla
Egret, Little	Egretta garzetta
Falcons (all species)	Falconidae
Flycatcher, Spotted	Muscicapa striata
Gadwall	Anas strepera
Garganey	Anas querquedula
Geese (all species)	Anser and Branta
Goshawk	Accipiter gentilis
Grebe, Little	Tachybaptus ruficollis
Harriers (all species)	Circus
Heron, Purple	Ardea purpurea
Kingfisher	Alcedo atthis
Kite, Black	Milvus migrans
Kite, Red	Milvus milvus
Lapwing	Vanellus vanellus
Martin, Sand	Riparia riparia
Nightjar	Caprimulgus europaeus
Osprey	Pandion haliaetus
Ouzel, Ring	Turdus torquatus
Owl (all species)	Strigiformes

<i>Common name</i>	<i>Scientific name</i>
Petrel, Storm	Hydrobates pelagicus
Pintail	Anas acuta
Pipit, Tree	Anthus trivialis
Plover, Golden	Pluvialis apricaria
Plover, Little Ringed	Charadrius dubius
Plover, Ringed	Charadrius hiaticula
Quail, Common	Coturnix coturnix
Redstart, Black	Phoenicurus ochruros
Redstart, Common	Phoenicurus phoenicurus
Sandpiper, Common	Actitis hypoleucos
Scaup	Aythya marila
Shag	Phalacrocorax aristotelis
Shearwater, Manx	Puffinus puffinus
Shelduck	Tadorna tadorna
Shoveler	Anas clypeata
Skylark	Alauda arvensis
Snipe, Jack	Lymnocyptes minimus
Sparrow, House	Passer domesticus
Sparrow, Tree	Passer montanus
Sparrowhawk	Accipiter nisus
Spoonbill	Platalea eucorodia
Starling	Sturnus vulgaris
Swan, Bewick's	Cygnus bewickii
Swan, Whooper	Cygnus cygnus
Swift	Apus apus
Tern (all species)	Sternidae
Thrush, Song	Turdus philomelos
Tit, Bearded	Panurus biarmicus
Twite	Carduelis flavirostris
Warbler, Grasshopper	Locustella naevia
Water Rail	Rallus aquaticus
Whinchat	Saxicola rubetra
Yellowhammer	Emberiza citrinella

PART II – DURING THE CLOSE SEASON

<i>Common name</i>	<i>Scientific name</i>
Teal	Anas crecca
Snipe	Gallinago gallinago
Woodcock	Scolopax rusticola

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 2**BIRDS WHICH MAY BE KILLED OR TAKEN**Section 2, 3, 5 and 23⁵²**PART I – OUTSIDE THE CLOSE SEASON**

<i>Common name</i>	<i>Scientific name</i>
Mallard	<i>Anas platyrhynchos</i>
Snipe	<i>Gallinago gallinago</i>
Teal	<i>Anas crecca</i>
Wigeon	<i>Anas penelope</i>
Woodcock	<i>Scolopax rusticola</i>

PART II – BY AUTHORISED PERSONS AT ALL TIMES

<i>Common name</i>	<i>Scientific name</i>
—	—

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 3**BIRDS WHICH MAY BE SOLD**

Sections 6 and 23

**PART I – ALIVE AT ALL TIMES IF RINGED AND BRED IN
CAPTIVITY**

<i>Common name</i>	<i>Scientific name</i>
Blackbird	<i>Turdus merula</i>
Brambling	<i>Fringilla montifringilla</i>
Bullfinch	<i>Pyrrhula pyrrhula</i>
Bunting, Reed	<i>Emberiza schoeniclus</i>
Chaffinch	<i>Fringilla coelebs</i>
Dunnock	<i>Prunella modularis</i>
Goldfinch	<i>Carduelis carduelis</i>
Greenfinch	<i>Carduelis chloris</i>
Jackdaw	<i>Corvus monedula</i>
Jay	<i>Garrulus glandarius</i>
Linnet	<i>Carduelis cannabina</i>
Magpie	<i>Pica pica</i>
Redpoll	<i>Carduelis flammea</i>
Siskin	<i>Carduelis spinus</i>
Starling	<i>Sturnus vulgaris</i>
Thrush, Song	<i>Turdus philomelos</i>
Twite	<i>Carduelis flavirostris</i>
Yellowhammer	<i>Emberiza citrinella</i>

PART II – DEAD AT ALL TIMES

<i>Common name</i>	<i>Scientific name</i>
Pigeon, Feral	<i>Columba livia</i>
Woodpigeon	<i>Columba palumbus</i>

PART III – DEAD FROM 1ST SEPTEMBER TO 28TH FEBRUARY

<i>Common name</i>	<i>Scientific name</i>
Mallard	<i>Anas platyrhynchos</i>
Snipe, Common	<i>Gallinago gallinago</i>
Teal	<i>Anas crecca</i>
Wigeon	<i>Anas penelope</i>
Woodcock	<i>Scolopax rusticola</i>

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 4

CAPTIVE BIRDS WHICH MAY BE POSSESSED BY REGISTERED PERSONS ONLY

Sections 7 and 23

<i>Common name</i>	<i>Scientific name</i>
Avocet	<i>Recurvirostra avosetta</i>
Bee-eater	<i>Merops apiaster</i>
Bittern	<i>Botaurus stellaris</i>
Bittern, Little	<i>Ixobrychus minutus</i>
Bluethroat	<i>Luscinia svecica</i>
Bunting, Cirl	<i>Emberiza cirlus</i>
Bunting, Lapland	<i>Calcarius lapponicus</i>
Bunting, Snow	<i>Plectrophenax nivalis</i>
Chough	<i>Pyrrhocorax pyrrhocorax</i>
Corncrake	<i>Crex crex</i>
Crake, Spotted	<i>Porzana porzana</i>
Crossbills (all species)	<i>Loxia</i>
Curlew, Stone	<i>Burhinus oedicephalus</i>
Divers (all species)	<i>Gavia</i>
Dotterel	<i>Charadrius morinellus</i>
Duck, Long-tailed	<i>Clangula hyemalis</i>
Falcons (all species)	<i>Falconidae</i>
Fieldfare	<i>Turdus pilaris</i>
Firecrest	<i>Regulus ignicapillus</i>
Godwit, Black-tailed	<i>Limosa limosa</i>
Grebe, Black-necked	<i>Podiceps nigricollis</i>
Grebe, Slavonian	<i>Podiceps auritus</i>
Greenshank	<i>Tringa nebularia</i>
Hawks, True (except old)	<i>Accipitridae</i> (except world vultures) that the genera <i>Aegypius</i> , is to say, Buzzards, <i>Gypaetus</i> , <i>Gypohierax</i> , Eagles, Harriers, Gyps, <i>Neophron</i> , Hawks and Kites (all <i>Sarcogyps</i> and species in each case)
	<i>Trigonoceps</i>
Hoopoe	<i>Upupa epops</i>
Kingfisher	<i>Alcedo atthis</i>
Oriole, Golden	<i>Oriolus oriolus</i>
Osprey	<i>Pandion haliaetus</i>
Owl (all species)	<i>Strigiformes</i>

<i>Common name</i>	<i>Scientific name</i>
Petrel, Leach's	<i>Oceanodroma leucorhoa</i>
Phalarope, Red-necked	<i>Phalaropus lobatus</i>
Plover, Kentish	<i>Charadrius alexandrinus</i>
Plover, Little ringed	<i>Charadrius dubius</i>
Quail, Common	<i>Coturnix coturnix</i>
Redstart, Black	<i>Phoenicurus ochruros</i>
Redwing	<i>Turdus iliacus</i>
Rosefinch, Scarlet	<i>Carpodacus erythrinus</i>
Ruff	<i>Philomachus pugnax</i>
Sandpiper, Green	<i>Tringa ochropus</i>
Sandpiper, Purple	<i>Calidris maritima</i>
Sandpiper, Wood	<i>Tringa glareola</i>
Scoter, Common	<i>Melanitta nigra</i>
Scoter, Velvet	<i>Melanitta fusca</i>
Serín	<i>Serinus serinus</i>
Shorelark	<i>Eremophila alpestris</i>
Shrike, Red-backed	<i>Lanius collurio</i>
Spoonbill	<i>Platalea leucorodia</i>
Stilt, Black-winged	<i>Himantopus himantopus</i>
Stint, Temminck's	<i>Calidris temminckii</i>
Tern, Black	<i>Chlidonias niger</i>
Tern, Little	<i>Sterna albifrons</i>
Tern, Roseate	<i>Sterna dougallii</i>
Tit, Bearded	<i>Panurus biarmicus</i>
Tit, Crested	<i>Parus cristatus</i>
Treecreeper, Short-toed	<i>Certhia brachydactyla</i>
Warbler, Cetti's	<i>Cettia cetti</i>
Warbler, Dartford	<i>Sylvia undata</i>
Warbler, Marsh	<i>Acrocephalus palustris</i>
Warbler, Savi's	<i>Locustella luscinioides</i>
Whimbrel	<i>Numenius phaeopus</i>
Woodlark	<i>Lullulla arborea</i>
Wryneck	<i>Jynx torquilla</i>

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 5

ANIMALS WHICH ARE PROTECTED

Sections 9, 10 and 23⁵³

<i>Common name</i>	<i>Scientific name</i>
Bats, Horseshoe (all species)	Rhinolophidae
Bats, Typical (all species)	Vespertilionidae
Beefly, Lesser	Bombylius minor
Cricket, Dotted bush	Leptophyes punctatissima
Cricket, Dark bush	Pholidoptera griseoptera
Frog, Common	Rana temporaria
Grasshopper, Lesser mottled	Stenobothrus stigmaticus
Lizard, Viviparous	Lacerta vivipara
Moth, Grey	Hadena caesia mananii
Moth, Scarce crimson and gold	Pyrausta sanguinalis
Seals (all species)	Pinnepedia
Shark, Basking	Cetorhinus maximus
Shrimp, Fairy	Chirocephalus diaphanus
Turtle (marine) (all species)	Dermochelyidae and Cheloniidae
Whales (all species)	Cetacea

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 6

ANIMALS WHICH MAY NOT BE KILLED OR TAKEN BY CERTAIN METHODS

Sections 11 and 23

<i>Common name</i>	<i>Scientific name</i>
Bats, Horseshoe (all species)	Rhinolophidae
Bats, Typical (all species)	Vespertilionidae
Dolphin, Bottle-nosed	Tursiops truncatus (otherwise known as Tursiops tursio)
Dolphin, Common	Delphinus delphis
Hedgehog	Erinaceus europaeus
Otter, Common	Lutra lutra
Porpoise, Harbour (otherwise known as Common porpoise)	Phocaena phocaena
Stoat ("Weasel")	Mustela erminea

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 7

PLANTS WHICH ARE PROTECTED

Sections 12 and 23⁵⁴

<i>Common name</i>	<i>Scientific name</i>
Agrimony	<i>Agrimonia eupatoria</i>
Alpine clubmoss	<i>Diphasiastrum alpinum</i>
Beech fern	<i>Phegopteris connectilis</i>
Blunt-leaved pondweed	<i>Potamogeton obtusifolius</i>
Brackish water crowfoot	<i>Ranunculus baudotii</i>
Burnet saxifrage	<i>Pimpinella saxifraga</i>
Carline thistle	<i>Carlina vulgaris</i>
Celery-leaved buttercup	<i>Ranunculus sceleratus</i>
Common bladderwort	<i>Utricularia vulgaris</i>
Common cow-wheat	<i>Melampyrum pratense</i>
Common sea lavender	<i>Limonium vulgare</i>
Common wintergreen	<i>Pyrola minor</i>
Cranberry	<i>Vaccinium oxycoccus</i>
Dioecious sedge	<i>Carex dioica</i>
Dodder	<i>Cuscuta epithymum</i>
Dune fescue	<i>Vulpia fasciculata</i>
Eelgrass	<i>Zostera marina</i>
Fennel pondweed	<i>Potamogeton pectinatus</i>
Few-flowered spike rush	<i>Eleocharis quinqueflora</i>
Field gentian	<i>Gentianella campestris</i>
Floating burr-reed	<i>Sparganium natans</i>
Floating club-rush	<i>Eleogiton fluitans</i>
Grass-leaved orache	<i>Atriplex littoralis</i>
Greater broomrape	<i>Orobanche rapum-genistae</i>
Greater spearwort	<i>Ranunculus lingua</i>
Hare's-foot clover	<i>Trifolium arvense</i>
Hay-scented buckler-fern	<i>Dryopteris aemula</i>
Horned pondweed	<i>Zannichellia palustris</i>
Iceland cress	<i>Rorippa islandica</i>
Isle of Man cabbage	<i>Coicya monensis monensis</i>
Ivy-leaved bellflower	<i>Wahlenbergia hederacea</i>
Juniper	<i>Juniperus communis</i>
Killarney filmy fern	<i>Trichomanes speciosum</i>
Least willow	<i>Salix herbacea</i>
Lesser clubmoss	<i>Selaginella selaginoides</i>

<i>Common name</i>	<i>Scientific name</i>
Lesser tussock-sedge	<i>Carex diandra</i>
Lesser water-plantain	<i>Baldellia ranunculoides</i>
Maidenhair fern	<i>Adiantum capillus veneris</i>
Marsh hawk's-beard	<i>Crepis paludosa</i>
Marsh stitchwort	<i>Stellaria palustris</i>
Mountain pansy	<i>Viola lutea</i>
Narrow buckler-fern	<i>Dryopteris carthusiana</i>
Narrow-leaved water-plantain	<i>Alisma lanceolatum</i>
Nodding bur-marigold	<i>Bidens cernua</i>
Orchid (all species)	<i>Orchidaceae</i>
Oyster plant	<i>Mertensia maritima</i>
Pale butterwort	<i>Pinguicula lusitanica</i>
Pale sedge	<i>Carex pallescens</i>
Parsley fern	<i>Cryptogramma crispa</i>
Parsley water-dropwort	<i>Oenanthe lachenalii</i>
Pennyroyal	<i>Mentha pulegium</i>
Pillwort	<i>Pilularia globulifera</i>
Pink water speedwell	<i>Veronica catenata</i>
Portland spurge	<i>Euphorbia portlandica</i>
Rock sea lavenders	<i>Limonium binervosum</i> agg.
Saltmarsh flat-sedge	<i>Blysmus rufus</i>
Sea-purslane	<i>Atriplex portulacoides</i>
Sea wormwood	<i>Seriphidium maritimum</i>
Shepherd's cress	<i>Teesdalia nudicaulis</i>
Slender spike-rush	<i>Eleocharis uniglumis</i>
Smooth cat's-ear	<i>Hypochaeris glabra</i>
Spring sandwort	<i>Minuartia verna</i>
Stagshorn club moss	<i>Lycopodium clavatum</i>
Strawberry clover	<i>Trifolium fragiferum</i>
Tubular water-dropwort	<i>Oenanthe fistulosa</i>
Viper's bugloss	<i>Echium vulgare</i>
Western clover	<i>Trifolium occidentale</i>
White sedge	<i>Carex curta</i>
Wilson's filmy fern	<i>Hymenophyllum wilsonii</i>
Wood fescue	<i>Festuca altissima</i>
Wood melick	<i>Melica uniflora</i>
Wood small-reed	<i>Calamagrostis epigejos</i>
Wood speedwell	<i>Veronica montana</i>
Wood vetch	<i>Vicia sylvatica</i>
Yellow waterlily	<i>Nuphar lutea</i>

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.

SCHEDULE 8

ANIMALS AND PLANTS TO WHICH SECTION 14 APPLIES

Sections 14 and 23

PART I – ANIMALS WHICH ARE ESTABLISHED IN THE WILD

<i>Common name</i>	<i>Scientific name</i>
Common carp/Mirror carp	<i>Cyprinus carpio</i>
Goose, Canada	<i>Branta canadensis</i>
Lizard, Common Wall	<i>Podarcis muralis</i>
Newt, Alpine	<i>Triturus alpestris</i>
Perch	<i>Perca fluviatilis</i>
Rat, Black	<i>Rattus rattus</i>
Roach	<i>Rutilus rutilus</i>
Roach x Rudd hybrids	<i>Scardinius spp.</i>
Rudd	<i>Scardinius erythrophthalmus</i>
Tench	<i>Tinca tinca</i>
Wallaby, Red-necked	<i>Macropus rufogriseus</i> ⁵⁵

PART II – PLANTS

<i>Common name</i>	<i>Scientific name</i>
American skunk cabbage	<i>Lysichiton americanum</i>
Canadian waterweed	<i>Elodea canadensis</i>
Common rhododendron	<i>Rhododendron ponticum</i>
Cord grass	<i>Spartina anglica</i>
Cotoneaster spp.	<i>Cotoneaster spp including C.horizontalis, C. integrifolius, C. simonsii</i>
Curly waterweed	<i>Lagarosiphon major</i>
Floating pennywort	<i>Hydrocotyle ranunculoides</i>
Fringed water lily	<i>Nymphoides peltata</i>
Giant knotweed	<i>Fallopia sachalinensis</i>
Giant rhubarb	<i>Gunnera tinctoria</i>
Giant salvinia	<i>Salvinia molesta</i>
Himalayan Balsam	<i>Impatiens glandulifera</i>
Hogweed, Giant	<i>Heracleum mantegazzianum</i>
Hottentot fig	<i>Carpobrotus edulis</i>
Keeled garlic	<i>Allium carinatum</i>
Kelp, Giant	<i>Macrocystis pyrifera</i>
Knotweed, Japanese	<i>Polygonum cuspidatum</i>
Mind-your-own business	<i>Soleirola solerolii</i>
Montbretia	<i>Crocasmia x crocosmiliflora</i>

<i>Common name</i>	<i>Scientific name</i>
New Zealand pygmyweed (also known as Australian stonecrop)	<i>Crassula helmsil</i> (also sold as <i>Crassula recurva</i> , <i>Tillaea recurva</i> , <i>Tillaea helmsil</i>)
Nuttall's water-weed	<i>Elodea nutallii</i>
NZ privet	<i>Griselinia littoralis</i>
Parrots feather	<i>Myriophyllum aquaticum</i>
Pirri-pirri burr	<i>Aceana novae-zelandiae</i> and <i>Aceana ovalifolia</i>
Sea buckthorn	<i>Hippophae rhamnoides</i>
Seaweed, Japanese	<i>Sargassum muticum</i>
Spanish bluebell and hybrids	<i>Hyacinthoides hispanica</i> and <i>Hyacinthoides x massartiana</i>
Three-cornered leek	<i>Allium triquetrum</i>
Water fern	<i>Azolla filiculoides</i>
Water hyacinth	<i>Eichhornia crassipes</i>
Water lettuce	<i>Pistia stratiotes</i>

NOTE: The common name or names given in the first column of this Schedule are included by way of guidance only; in the event of any dispute or proceedings, the common name or names shall not be taken into account.⁵⁶

SCHEDULE 9⁵⁷

SCHEDULE 10**REPEAL OF ENACTMENTS**

Section 40

PART I - REPEAL OF ACTS

Part I repeals the following Acts wholly —

Sea Gull Preservation Act 1867
Salmon and Fresh-water Fishery Act 1882
Game Act 1928
Sea Gull Preservation Amendment Act 1930
Wild Birds Protection Act 1932
Protection of Birds Act 1955
Wild Birds Protection (Amendment) Act 1975

and the following Acts in part —

Prevention of Damage by Agricultural Pests Act 1956
Game (Hares) Act 1971
Endangered Species (Import and Export) Act 1981
Treasury Act 1985
Fines Act 1986.

The repeal of the Wild Birds Protection Acts 1932 to 1975 not in operation as respects bird sanctuaries defined in any order under s 11 of the Wild Birds Protection Act 1932 which was in force immediately before 1/1/1991.]

PART II – REPEAL OF PUBLIC DOCUMENTS

Part II repeals the following public documents in part —

Customs and Excise Acts (Application) Order 1979 (GC38/79)
Customs and Excise Acts (Application) (Amendment) Order 1979
(GC249/79)
Customs and Excise (Transfer of Functions) Order 1980 (GC29/80)
Department of Agriculture, Fisheries and Forestry Order 1986
(GC121/86).]

ENDNOTES

Table of Legislation History

Legislation	Year and No	Commencement

Table of Renumbered Provisions

Original	Current

Table of Endnote References

-
- ¹ Subs (1) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a).
- ² Subs (5) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a).
- ³ Subs (3) amended by SD155/10 Sch 3.
- ⁴ Subpara (i) amended by Agriculture (Miscellaneous Provisions) Act 2000 s 2.
- ⁵ Para (a) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a).
- ⁶ Subs (2) amended by Agriculture (Miscellaneous Provisions) Act 2000 s 2, by SD155/10 Sch 9 and by SD2014/06.
- ⁷ Subs (6) added by Agriculture (Miscellaneous Provisions) Act 2000 s 2.
- ⁸ Para (a) amended by Criminal Justice Act 1996 Sch 2.
- ⁹ Para (c) amended by Criminal Justice Act 1996 Sch 2.
- ¹⁰ Para (d) amended by Criminal Justice Act 1996 Sch 2.
- ¹¹ Para (e) amended by Criminal Justice Act 1996 Sch 2.
- ¹² Para (f) inserted by Criminal Justice Act 1996 Sch 2.
- ¹³ Subs (4) amended by Criminal Justice Act 1996 Sch 2.
- ¹⁴ Subs (9) amended by Rehabilitation of Offenders Act 2001 Sch 2.
- ¹⁵ Subs (2A) inserted by Agriculture (Miscellaneous Provisions) Act 2000 s 2.
- ¹⁶ Subs (5A) inserted by Rehabilitation of Offenders Act 2001 Sch 2.
- ¹⁷ Para (c) amended by Veterinary Surgeons Act 2005 Schs 2 and 3.
- ¹⁸ Subs (1) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a) and (b)(i).
- ¹⁹ Subs (4) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a).
- ²⁰ Subs (4A) inserted by Agricultural (Miscellaneous Provisions) Act 2008 s 3(b)(ii).
- ²¹ Para (d) inserted by Criminal Justice Act 1996 Sch 2.

-
- ²² Para (f) inserted by Criminal Justice Act 1996 Sch 2.
- ²³ Para (a) amended by Criminal Justice Act 1996 Sch 2.
- ²⁴ Subs (5) amended by Criminal Justice Act 1996 Sch 2.
- ²⁵ Subs (6) amended by Criminal Justice Act 1996 Sch 2.
- ²⁶ Para (a) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a).
- ²⁷ Para (b) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(a).
- ²⁸ S 15 repealed by Endangered Species Act 2010 Sch 5.
- ²⁹ Subs (1) amended by Agriculture (Miscellaneous Provisions) Act 2000 s 2.
- ³⁰ Subs (1A) inserted by Agriculture (Miscellaneous Provisions) Act 2000 s 2.
- ³¹ Subs (6) amended by Agriculture (Miscellaneous Provisions) Act 2000 s 2.
- ³² Para (c) repealed by Police Powers and Procedures Act 1998 Sch 5.
- ³³ Subs (2) amended by Police Powers and Procedures Act 1998 Sch 4.
- ³⁴ Subs (1) amended by Statute Law Revision Act 1997 Sch 1.
- ³⁵ Subs (2) amended by Statute Law Revision Act 1997 Sch 1.
- ³⁶ Subs (3) amended by Statute Law Revision Act 1997 Sch 1.
- ³⁷ Subs (1) repealed by Endangered Species Act 2010 Sch 5.
- ³⁸ Para (b) amended by SD155/10 Sch 9 and by SD2014/06.
- ³⁹ Subs (1) amended by SD2014/06.
- ⁴⁰ S 26A inserted by Flood Risk Management Act 2013 s 88.
- ⁴¹ Para (a) amended by SD155/10 Schs 3 and 11.
- ⁴² Para (a) amended by Statute Law Revision Act 1992 Sch 1.
- ⁴³ Subs (6) amended by Transfer of Governor's Functions Act 1992 Sch 1.
- ⁴⁴ Subs (13) added by Statute Law Revision Act 1997 Sch 1.
- ⁴⁵ Subs (1) amended by Agricultural (Miscellaneous Provisions) Act 2008 s 3(c).
- ⁴⁶ Subs (6) amended by SD486/94 and by SD155/10 Schs 3, 5 and 11.
- ⁴⁷ Subs (5) amended by SD486/94, by SD155/10 Schs 3, 5, 9 and 11 and by SD2014/06.
- ⁴⁸ Subs (6A) inserted by Fisheries Act 2012 s 76.
- ⁴⁹ Subs (6B) inserted by Fisheries Act 2012 s 76.
- ⁵⁰ Definition of "the Wildlife Committee" added by Endangered Species Act 2010 Sch 4.
- ⁵¹ Majority of provisions in force by 1/1/1991 by GC9/91. ADO (s 15, Sch 9 and the entry in Sch 10 as they amend or repeal the Endangered Species (Import and Export) Act 1981) 9/12/2003 (SD824/03). The whole Act in force except to the extent that ss 1 to 8 do not apply to bird sanctuaries controlled by Wild Birds Protection Acts 1932 to 1975 [see Art. 2(2) and (3) GC9/91].
- ⁵² Sch 2 substituted by SD129/04.
- ⁵³ Sch 5 substituted by SD457/04.
- ⁵⁴ Sch 7 substituted by SD129/04.
- ⁵⁵ Part I amended by SD135/11.
- ⁵⁶ Part II amended by SD135/11.
- ⁵⁷ Sch 9 repealed by Endangered Species Act 2010 Sch 5.